arXiv:math/0105117v1 [math.QA] 15 May 2001

A locally compact quantum group analogue of the normalizer of $\mathbf{SU(1,1)}$ in $\mathbf{SL(2,\mathbb{C})}$

Erik Koelink and Johan Kustermans

Technische Universiteit Delft
Faculteit ITS
Afdeling Toegepaste Wiskundige Analyse
Mekelweg 4
2628CD Delft
The Netherlands

`h.t.koelink@its.tudelft.nl` & `j.kustermans@its.tudelft.nl`

Abstract

S.L. Woronowicz proved in 1991 that quantum SU(1,1) does not exist as a locally compact quantum group. Results by L.I. Korogodsky in 1994 and more recently by Woronowicz gave strong indications that the normalizer $\widetilde{SU}(1,1)$ of $SU(1,1)$ in $SL(2,\mathbb{C})$ is a much better quantization candidate than $SU(1,1)$ itself. In this paper we show that this is indeed the case by constructing $\widetilde{SU}_q(1,1)$, a new example of a unimodular locally compact quantum group (depending on a parameter $0 < q < 1$) that is a deformation of $\widetilde{SU}(1,1)$. After defining the underlying von Neumann algebra of $\widetilde{SU}_q(1,1)$ we use a certain class of q-hypergeometric functions and their orthogonality relations to construct the comultiplication. The coassociativity of this comultiplication is the hardest result to establish. We define the Haar weight and obtain simple formulas for the antipode and its polar decomposition. As a final result we produce the underlying C*-algebra of $\widetilde{SU}_q(1,1)$. The proofs of all these results depend on various properties of q-hypergeometric ${}_1\varphi_1$ functions.

Mathematics subject classification 2000: 33D80, 46L89

Introduction

Arguably one of the most important and simplest non-compact Lie groups is the $SU(1,1)$ group, which is isomorphic to $SL(2,\mathbb{R})$. In 1990, one of the first attempts to construct a quantum version of $SU(1,1)$ was made in [20] and [21] by T. Masuda, K. Mimachi, Y. Nakagami, M. Noumi, Y. Saburi and K. Ueno and independently, in [10] by L. Vaksman and L. Korogodsky. We follow [20] and [21] because these expositions are more elaborate. Their starting point is a real form $\mathcal{U}_q(\mathfrak{su}(1,1))$ of the quantum universal enveloping algebra $\mathcal{U}_q(\mathfrak{sl}(2,\mathbb{C}))$ (defined in [20, Eq. (1.9)]). Intuitively, one should view upon $\mathcal{U}_q(\mathfrak{su}(1,1))$ as a quantum universal enveloping algebra of the 'quantum Lie algebra' of the still to be constructed locally compact quantum group $SU_q(1,1)$. The dual $\mathcal{A}$ of $\mathcal{U}_q(\mathfrak{su}(1,1))$ is turned into some sort of topological Hopf*-algebra ([20, Sec. 2] and [21, Eq. (2.6)]) which is naively referred to as a topological quantum group. In a next step the coordinate Hopf*-algebra $A_q(SU(1,1))$ is given as a *-subalgebra of $\mathcal{A}$ that also inherits the comultiplication, counit and antipode from $\mathcal{A}$ ([20, Eq. (2.7)] and [21, Eq. (0.9)]).
In this philosophy, they first introduce infinite dimensional infinitesimal representations of quantum $SU_q(1,1)$ (see [21, Eq. (1.1)]) which they then exponentiate to infinite dimensional unitary corepresentations of $\mathcal{A}$ (see [21, Eq. (1.2)]) providing hereby the quantum analogues of the discrete, continuous and complementary series of $SU(1,1)$ but also a new strange series of corepresentations. In an attempt to get hold of the C*-algebra $\mathcal{B}$ that should be viewed as the quantum analogue of the space $C_0(SU(1,1))$, they follow the strategy of first transforming the generators and relations of $A_q(SU(1,1))$ formally into other elements that produce a *-algebra B (not inside $A_q(SU(1,1))$!) definable by generators and relations that, unlike in the case of $A_q(SU(1,1))$, can be faithfully represented by bounded operators. The C*-algebra $\mathcal{B}$ is then defined to be the universal enveloping C*-algebra of B.

Other important contributions (other than the ones that will be mentioned later on) to the study of different aspects of quantum $SU(1,1)$ have been made by T. Kakehi, T. Masuda, K. Ueno in [8], by D. Shklyarov, S. Sinel'shchikov & L. Vaksman in [24].

The authors of [21] seem to get a little bit overenthusiastic by claiming the existence of a comultiplication on $\mathcal{B}$ turning $\mathcal{B}$ into a Hopf*-algebra (see [21, Prop. 7]). This is in fact in stark contrast with the result, proven by S.L. Woronowicz in 1991, that showed that quantum $SU(1,1)$ does not exist as a locally compact quantum group (see [27, Thm. 4.1 and Sec. 4.C]). Not surprisingly, this was considered to be quite a setback for the theory of locally compact quantum groups in the operator algebra setting.

Recall that $SU(1,1)$ is the linear Lie group $\{\, X \in SL(2,\mathbb{C}) \mid X^*UX = U \,\}$, where $U = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. In 1994, a breakthrough in this stalemate was forced by L.I. Korogodsky (see [9]) who studied deformations of the linear Lie group $\widetilde{SU}(1,1)$ instead of that of $SU(1,1)$. Here, $\widetilde{SU}(1,1)$ denotes the Lie group $\{\, X \in SL(2,\mathbb{C}) \mid X^*UX = U \text{ or } X^*UX = -U \,\}$ which is in fact the normalizer of $SU(1,1)$ in $SL(2,\mathbb{C})$. In this paper, we will follow up on Korogodsky's and Woronowicz' ideas to introduce a locally compact quantum group $\widetilde{SU}_q(1,1)$, depending on a parameter $0 < q < 1$, that is the quantum version of this group $\widetilde{SU}(1,1)$. In doing so, we believe we also make a strong case for the use of q-hypergeometric functions in the operator algebra approach to quantum groups.

But what is actually meant by a locally compact quantum group? This question has kept a lot of people busy for the last 20 years and the most satisfying answer has been given by S. Vaes and the second author in [18] (see the introduction of this same paper for an extensive account of the history and the importance of different people in the development of the theory of locally compact quantum groups). The paper [18] is written in the C*-algebra setting but it will be easier to use the von Neumann algebraic approach as introduced in [19]. Both approaches are completely equivalent and the last result of this paper produces the generic C*-algebraic quantum group out of the von Neumann algebraic one. In order to see what we are aiming for, we formulate the definition of a von Neumann algebraic quantum group as defined in [19].

Definition 1. *Consider a von Neumann algebra M together with a unital normal *-homomorphism $\Delta : M \to M \otimes M$ such that $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$. Assume moreover the existence of*

1. *a normal semifinite faithful weight φ on M that is left invariant: $\varphi((\omega \otimes \iota)\Delta(x)) = \varphi(x)\omega(1)$ for all $\omega \in M_*^+$ and $x \in \mathcal{M}_\varphi^+$.*
2. *a normal semifinite faithful weight ψ on M that is right invariant: $\psi((\iota \otimes \omega)\Delta(x)) = \psi(x)\omega(1)$ for all $\omega \in M_*^+$ and $x \in \mathcal{M}_\psi^+$.*

Then we call the pair (M, Δ) a von Neumann algebraic quantum group.

For a discussion about the consequences of this definition and related notations we refer to [19, Sec. 1]. For quite a while, the major drawback of the theory of non-compact quantum groups was the lack of a whole array of different examples of non-compact locally compact quantum groups. So the importance of this paper lies mainly in the fact that we add an example to the rather short list of existing atomic non-compact quantum groups like quantum E(2), quantum $ax + b$ and quantum $az + b$. Also note that this new locally compact quantum group is the first analogue of a non-compact semi-simple Lie group. By atomic we mean that these examples (up till now) can not be constructed out of simpler quantum groups through different existing theoretical construction procedures of which the most important one is arguably the double crossed product construction. In turn, this new example $\widetilde{SU}_q(1,1)$ can now serve as one of the ingredients in these construction procedures. It is moreover very conceivable that the importance of $SU(1,1)$ in the classical group theory will be parallelled by that of $\widetilde{SU}_q(1,1)$ in the quantum group setting.

In the immediate future we hope to calculate the dual of $\widetilde{SU}_q(1,1)$, unravel its corepresentation theory and obtain a formula for the Plancherel measure. For one, we most definitely plan to fit the corepresentations of [21] rigorously in the framework of locally compact quantum groups. It is to be expected that new results for q-hypergeometric functions can emerge this way.

Since the ideas of Korogodsky are a chief motivation for this paper, we will briefly discuss the most important results of [9]. Let $(\mathcal{A}_q, \Delta)$ be the Hopf*-algebra associated to $\widetilde{SU}_q(1,1)$ (in [9], $\mathcal{A}_q$ is denoted by $\mathfrak{S}$). We give a precise definition of $(\mathcal{A}_q, \Delta)$ in Eqs. (1.1) and (1.2). The *-algebra $\mathcal{A}_q$ is generated by elements α, γ and a central self-adjoint involution e. Define two central orthogonal projections $p_\pm$ in $\mathcal{A}_q$ as $p_\pm = \frac{1}{2}(1 \pm e)$ and define the *-subalgebras $\mathcal{A}_q^\pm$ of $\mathcal{A}_q$ as $\mathcal{A}_q^\pm = p_\pm \mathcal{A}_q$. Thus, $\mathcal{A}_q = \mathcal{A}_q^+ \oplus \mathcal{A}_q^-$ (in [9], $\mathcal{A}_q^\pm$ are denoted by $\mathfrak{R}^\pm$). Let us also set $\gamma_\pm = p_\pm \gamma$.
Define a *-homomorphism $\Delta_+ : \mathcal{A}_q^+ \to \mathcal{A}_q^+ \otimes \mathcal{A}_q^+$ such that $\Delta_+(a) = (p_+ \otimes p_+)\Delta(a)$ for all $a \in \mathcal{A}_q^+$. Then $(\mathcal{A}_q^+, \Delta_+)$ is a Hopf*-algebra which should be thought of as the q-deformation of the coordinate Hopf*-algebra associated to the group $SU(1,1)$.
Most of the relevant representations of $\mathcal{A}_q$ are infinite dimensional so that a lot of care has to be taken to make the notion of representations of $\mathcal{A}_q$ more precise (see [23]). Consider a Hilbert space H, a dense subspace D of H and a unital algebra representation π of $\mathcal{A}_q$ on D such that $\langle \pi(a)v, w \rangle = \langle v, \pi(a^*)w \rangle$ for all $a \in \mathcal{A}_q$ and $v, w \in D$. Then we call π a *-representation of $\mathcal{A}_q$ *in* H. We set $\mathcal{D}(\pi) = D$. We call π well-behaved if $\pi(\gamma^*\gamma)$ is an essentially self-adjoint operator in H and if the spectrum $\sigma\big(\overline{\pi(\gamma^*\gamma)}\big) \subseteq q^{2\mathbb{Z}}$, where $\overline{\pi(\gamma^*\gamma)}$ denotes the closure of $\pi(\gamma^*\gamma)$ as an operator in H. Similar definitions are used for *-representations of $\mathcal{A}_q^\pm$ in H (where γ is replaced by $\gamma_\pm$). The spectral condition is not present in [9], a weaker spectral condition was introduced in [30].
It is clear that we can identify *-representations of $\mathcal{A}_q^\pm$ with *-representations π of $\mathcal{A}_q$ such that $\pi(p_\mp) = 0$, or equivalently $\pi(e) = \pm 1$.
Since e belongs to the center of $\mathcal{A}_q$, each irreducible *-representation of $\mathcal{A}_q^\pm$ is necessarily an irreducible *-representation of either $\mathcal{A}_q^+$ or $\mathcal{A}_q^-$. Korogodsky classified all well-behaved *-representations of $\mathcal{A}_q$ in [9, Prop. 2.4], most of them are infinite dimensional. K. Schmüdgen pointed us to the necessity of imposing some sort of spectral condition for this classification result to be true (and which is completed in [30]).
Given two well-behaved infinite dimensional irreducible *-representations π_1, π_2 of $\mathcal{A}_q$ in Hilbert spaces H_1, H_2 respectively. Korogodsky showed in [9, Thm. 6.1] that there does not exist a well-behaved *-representation π of $\mathcal{A}_q$ in $H_1 \otimes H_2$ such that $\mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2) \subseteq \mathcal{D}(\pi)$ and $\pi(a)\, v = (\pi_1 \odot \pi_2)(\Delta(a))\, v$ for all $a \in \mathcal{A}_q$ and $v \in \mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2)$, where $\odot$ denotes the algebraic tensor product. Thus, loosely speaking, the tensor product of two well-behaved infinite dimensional irreducible *-representations π_1 and π_2 of

$\mathcal{A}_q$ does not exist. Applied to two well-behaved infinite dimensional *-representations of $\mathcal{A}_q^+$, this result corresponds to the non-existence of quantum $SU(1,1)$ as a locally compact quantum group as proven in [27].
However, Korogodsky also showed that the situation is not as bad as the above result on first sight suggests. Therefore consider two well-behaved *-representations π_1, π_2 of $\mathcal{A}_q$ in Hilbert spaces H_1, H_2 such that for $i = 1,2$, π_i is the finite direct sum of well-behaved infinite dimensional irreducible *-representations of $\mathcal{A}_q$ for which the number of direct summands of π_i that are *-representations of $\mathcal{A}_q^+$ equals the number of direct summands of π_i that are *-representations of $\mathcal{A}_q^-$. Then [9, Thm. 6.2 and 6.3] imply the existence of a well-behaved *-representation π of $\mathcal{A}_q$ in $H_1 \otimes H_2$ such that $\mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2) \subseteq \mathcal{D}(\pi)$ and $\pi(a)\, v = (\pi_1 \odot \pi_2)(\Delta(a))\, v$ for all $a \in \mathcal{A}_q$ and $v \in \mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2)$. It should however be pointed out that π is not unique!

Around 1996, Woronowicz picked up on this observation to start his study of quantum $\widetilde{SU}_q(1,1)$ on the Hilbert space level (see [30]). Instead of focusing on all well-behaved *-representation of $\mathcal{A}_q$, he redirected his attention to *-representations that are direct sums (or more precisely, direct integrals) of the sort described above. For this purpose, Woronowicz introduces in [30, Def. 1.1] the notion of a $\widetilde{SU}_q(1,1)$-quadruples on a Hilbert space. Let us use some slightly different terminology in order to stay closer to Korogodsky's viewpoint.
Let π be a well-behaved *-representation of $\mathcal{A}_q$ in some Hilbert space H and Y a closed linear operator in H such that Phase $\overline{\pi(\gamma)}$ commutes with Y, the domain of Y and Y^* agree and $\pi\big(\sqrt{q}\, e\, \gamma^* - \alpha\big) \subseteq Y$. Then we call (π, Y) a $\widetilde{SU}_q(1,1)$-representation in H. It follows that for such a $\widetilde{SU}_q(1,1)$-representation, the quadruple $(\overline{\pi(\alpha)}, \overline{\pi(\gamma)}, \overline{\pi(e)}, Y)$ forms a $\widetilde{SU}_q(1,1)$ of unbounded type in the sense of [30].
If $(\tilde{\alpha}, \tilde{\gamma}, \tilde{e}, \tilde{Y})$ is a $\widetilde{SU}_q(1,1)$-quadruple of unbounded type, it follows from [30, Thm. 3.7] that it arises from such a $\widetilde{SU}_q(1,1)$-representation. Moreover, [30, Thm. 1.3] gives a precise meaning to the statement that π contains as much irreducible well-behaved *-representations of $\mathcal{A}_q^+$ as of $\mathcal{A}_q^-$. In [30, Thm. 1.4] it is then shown that, in accordance with the above formal discussion, the tensor product of two $\widetilde{SU}_q(1,1)$-quadruples can be defined as a new $\widetilde{SU}_q(1,1)$-quadruple. Recasted in the terminology of this introduction this means that given two $\widetilde{SU}_q(1,1)$-representations (H_1, π_1, Y_1), (H_2, π_2, Y_2) it is possible to construct a new $\widetilde{SU}_q(1,1)$-representation $(H_1 \otimes H_2, \pi, Y)$ such that $\mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2) \subseteq \mathcal{D}(\pi)$ and $\pi(a)\, v = (\pi_1 \odot \pi_2)(\Delta(a))\, v$ for all $a \in \mathcal{A}_q$ and $v \in \mathcal{D}(\pi_1) \odot \mathcal{D}(\pi_2)$. The extra information contained in Y moreover allows to get rid of the uniqueness problems mentioned before.
There is however one drawback to [30]. At the moment of writing this paper, there was still a gap in the proof of the associativity of the tensor product construction $\widetilde{SU}_q(1,1)$-quadruples. Although it is difficult to estimate the seriousness of this gap, it corresponds to the most difficult part of the proof of the coassociativity of the comultiplication in this paper.

In 1999, J. Stokman together with the first author studied the properties of a left invariant weight h on the C*-algebra C of quantum $SU(1,1)$ (see [14]). However, remember that it is impossible to define a comultiplication on C turning it into a locally compact quantum group. Instead, one looks at a suitable dense subspace B of C^* so that for ω_1, ω_2 it is possible to define the product $\omega_1 \star \omega_2$ so that on a formal level, $\omega_1 \star \omega_2 = (\omega_1 \otimes \omega_2)\Delta$. It turns out that the invariant weight h can be written as a sum of positive functionals in B and as a consequence $(\omega \star h)(x)$ can be defined as a pointwise convergent sum for suitable $\omega \in B$ and suitable $x \in C$. The left invariance statement then becomes $(\omega \star h)(x) = \omega(1)\, h(x)$. The benefit of [14] to this paper lies in the fact that the definition of $\omega_1 \star \omega_2$ depends on a class of special functions that arise formally as Clebsch-Gordan coefficients. These special functions can be extended in such a way that they serve as the principal set of data to construct the locally compact quantum group $\widetilde{SU}_q(1,1)$. Moreover, the formula for the left Haar weight of $\widetilde{SU}_q(1,1)$ is a straightforward generalization of the formula defining h.

A naive response to this all would be to presume that given all these existing results it should be not so difficult to construct $\widetilde{SU}_q(1,1)$ as a full blown locally compact quantum group. But this seems to be far from the truth. The family of special functions produced in [14] can be easily extended from the context of $SU(1,1)$ to the context of $\widetilde{SU}(1,1)$, but the orthogonality and completeness of the relevant family of functions needs some non-trivial special function theory that can be found in [2] and [3] (see Proposition

3.2). This is in fact the only place where we heavily rely on the theory of q-hypergeometric not covered in the appendix of this paper. Given these special functions it is not hard to define the comultiplication of $\widetilde{SU}_q(1,1)$, the hard part of the construction of $\widetilde{SU}_q(1,1)$ lies in the proof of the coassociativity of the comultiplication. Similarly, the left Haar weight is pretty easy to define, but checking that it is left invariant in the sense of [19] requires some non-trivial quantum group techniques.

Whereas [9] is mainly a motivational paper for this one, [30] is not only motivational (especially the use of the reflection operator u introduced in notation 2.2) but also gives an alternative approach to $\widetilde{SU}_q(1,1)$. The advantage of the approach of [30] lies mainly in the fact that the role of the generators and relations is more explicit and constructional. In order to do so, Woronowicz also presents a beautiful operator theoretic theory of balanced extensions, similar to the theory of self-adjoint extension of symmetric operators. As a consequence of this all, it should not be so difficult to prove that the C*-algebra introduced in proposition 4.15 is generated by the generators and relations, introduced in [30, Def. 1.1], and in the sense of [28]. We on the other hand take a more pragmatic stance and approach the subject by using special functions. As a disadvantage, it follows that the role of generators and relations is less clear (but still lurking in the background), but as a clear advantage we can prove the coassociativity of the comultiplication, prove the left invariance of the left Haar weight and also give an explicit formula for the multiplicative unitary. It also allows us to circumvent the theory of balanced extensions and stick to more standard operator theoretic techniques. As a consequence, the proofs of this paper are different from the ones in [30]. A notable exception is the use of an operator, introduced in [30], in the proof of the last (albeit simplest) step in the proof of Theorem 3.11.

The paper is organized as follows. In the first section we introduce the Hopf*-algebra associated to $\widetilde{SU}_q(1,1)$ and realize it as a *-algebra of unbounded operators on some Hilbert space. In the subsequent section the underlying von Neumann algebra is introduced. We define the comultiplication in section 3 and claim its coassociativity at the end of the section but the proof of this fact is given in section 5 in order to enhance the readability of the paper. In section 4 we construct the Haar weights and prove their invariance, thereby proving that $\widetilde{SU}_q(1,1)$ is indeed a locally compact quantum group. We end section 4 by calculating the underlying C*-algebraic quantum group. The appendix contains most of the basic results that we will need from the theory of special functions.

Acknowledgement. We would like to thank S. L. Woronowicz for insightful discussions on quantum $\widetilde{SU}(1,1)$, providing the preliminary version of [30] and giving a series of lectures on [30] in Trondheim (April 2000), which the second author attended.

Notations and conventions.

The set of all natural numbers, not including 0, is denoted by $\mathbb{N}$. Also, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.
Fix a number $q > 0$. Let $a \in \mathbb{C}$. If $n \in \mathbb{N}_0$, the q-shifted factorial $(a;q)_n \in \mathbb{C}$ is defined as $(a;q)_n = \prod_{i=0}^{n-1}(1-q^i a) \quad (\text{so } (a;q)_0 = 1)$. From now on, we assume that $q < 1$. Then $(a;q)_\infty \in \mathbb{C}$ is defined as $(a;q)_\infty = \lim_{n\to\infty}(a;q)_n$. Using some basic infinite product theory, one checks that this limit exists and that the function $a \in \mathbb{C} \mapsto (a;q)_\infty$ is analytic. Also, $(a;q)_\infty = 0$ if and only if $a \in q^{-\mathbb{N}_0}$. We also use the notation $(a_1,\ldots,a_m;q)_k = (a_1;q)_k \ldots (a_m;q)_k$ if $a_1,\ldots,a_m \in \mathbb{C}$ and $k \in \mathbb{N}_0 \cup \{\infty\}$.

If $a,b,z \in \mathbb{C}$, we define

$$\Psi\left(\begin{array}{c} a \\ b \end{array} ; q, z\right) = \Psi(a;b;q,z) := \sum_{n=0}^{\infty} \frac{(a;q)_n\,(b\,q^n;q)_\infty}{(q\,;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\,. \tag{1}$$

We collected some further basic information about these Ψ function in the appendix. See [6] for an extensive treatment on q-hypergeometric functions.

Let $B_1,\ldots,B_n$ be sets such that $B_i = \mathbb{T}$ or $B_i = -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$. Set $I = \{\, i \in \{1,\ldots,n\} \mid B_i = -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}\,\}$. Consider a set $K \subseteq (-q^{\mathbb{Z}} \cup q^{\mathbb{Z}})^I$ and a function $f : \{\,(x_1,\ldots,x_n) \in B_1 \times \ldots \times B_n \mid (x_i)_{i\in I} \in K\,\} \to \mathbb{C}$. Then we set $f(x) := 0$ for $(x_1,\ldots,x_n) \in B_1 \times \ldots \times B_n$ such that $(x_i)_{i\in I} \notin K$. (2)

If f is a function, the domain of f will be denoted by $D(f)$. If X is a set, the identity mapping on X will be denoted by ι_X and most of the time even by ι. If H is a Hilbert space, 1_H denotes ι_H. The set of all complex valued functions on X is denoted by $\mathcal{F}(X)$, the set of all elements in $\mathcal{F}(X)$ having finite support, is denoted by $\mathcal{K}(X)$. Let V be a vector space and S a subset of V. Then $\langle S \rangle$ denotes the linear span of S in V.

Consider a locally compact space Ω with a regular Borel measure μ on it. If $g \in \mathcal{L}^2(\Omega,\mu)$, we denote the class of g in $L^2(\Omega,\mu)$ by $[g]$. If f is a measurable function on Ω, we define the linear operator M_f in $L^2(\Omega,\mu)$ such that $D(M_f) = \{\, [g] \mid g \in \mathcal{L}^2(\Omega,\mu) \text{ such that } fg \in \mathcal{L}^2(\Omega,\mu) \,\}$ and $M_f([g]) = [fg]$ for all such classes $[g] \in D(M_f)$.

Let S, T be two linear operators acting in a Hilbert space H. We say that $S \subseteq T$ if $D(S) \subseteq D(T)$ and $S(v) = T(v)$ for all $v \in D(S)$. Following [30] we call S balanced if $D(S) = D(S^*)$.

The symbol $\odot$ will be used to denote the algebraic tensor product of vector spaces and linear mappings. The symbol $\otimes$ on the other hand will denote the tensor product of Hilbert spaces, von Neumann algebras and sufficiently continuous linear mappings.

If $n \in \mathbb{N}$ and X is locally compact space, X^n will denote the n-fold product set $X^n = X \times \ldots \times X$.

Consider von Neumann algebras M,N acting on Hilbert spaces H,K respectively and $\pi : M \to N$ a normal *-homomorphism. Let T be a densely defined, closed linear operator in H affiliated with M (in the von Neumann algebraic sense) and $T = U\,|T|$ the polar decomposition of T. Then there exists a unique positive operator P in K such that $f(P) = \pi(f(|T|))$ for all $f \in \mathcal{L}^\infty(\mathbb{R}^+)$. Now we set $\pi(T) = \pi(U)\,P$.

1. The Hopf*-algebra underlying quantum $\widetilde{SU}(1,1)$.

In order to resolve the problems surrounding quantum $SU(1,1)$, Korogodsky proposed in [9] to construct the quantum version of $\widetilde{SU}(1,1)$ instead of constructing the quantum version of $SU(1,1)$ itself. He suggested that the Hopf*-algebra of this quantum group should be the one that we describe now.

Throughout this paper, we fix a number $0 < q < 1$. Define $\mathcal{A}_q$ to be the unital *-algebra generated by elements α_0, γ_0 and e_0 and relations

$$\begin{array}{ll}
\alpha_0^\dagger \alpha_0 - \gamma_0^\dagger \gamma_0 = e_0 & \alpha_0 \alpha_0^\dagger - q^2\, \gamma_0^\dagger \gamma_0 = e_0 \\
\gamma_0^\dagger \gamma_0 = \gamma_0\, \gamma_0^\dagger & \\
\alpha_0\, \gamma_0 = q\, \gamma_0\, \alpha_0 & e_0^\dagger = e_0 \\
\alpha_0\, \gamma_0^\dagger = q\, \gamma_0^\dagger \alpha_0 & e_0^2 = 1 \\
\alpha_0\, e_0 = e_0\, \alpha_0 & \\
\gamma_0\, e_0 = e_0\, \gamma_0 &
\end{array} \tag{1.1}$$

where † denotes the *-operation on $\mathcal{A}_q$ (in order to distinguish this kind of adjoint with the adjoints of possibly unbounded operators in Hilbert spaces).

There exists a unique unital *-homomorphism $\Delta_0 : \mathcal{A}_q \to \mathcal{A}_q \odot \mathcal{A}_q$ such that

$$\begin{aligned}
\Delta_0(\alpha_0) &= \alpha_0 \otimes \alpha_0 + q\,(e_0\, \gamma_0^\dagger) \otimes \gamma_0 \\
\Delta_0(\gamma_0) &= \gamma_0 \otimes \alpha_0 + (e_0 \alpha_0^\dagger) \otimes \gamma_0 \\
\Delta_0(e_0) &= e_0 \otimes e_0 \ .
\end{aligned} \tag{1.2}$$

The pair $(\mathcal{A}_q, \Delta_0)$ turns out to be a Hopf*-algebra with counit ε_0 and antipode S_0 determined by

$$\begin{array}{ll}
S_0(\alpha_0) = e_0\, \alpha_0^\dagger & \varepsilon_0(\alpha_0) = 1 \\
S_0(\alpha_0^\dagger) = e_0\, \alpha_0 & \varepsilon_0(\gamma_0) = 0 \\
S_0(\gamma_0) = -q\, \gamma_0 & \varepsilon_0(e_0) = 1 \\
S_0(\gamma_0^\dagger) = -\frac{1}{q}\, \gamma_0^\dagger & \\
S_0(e_0) = e_0 \ . &
\end{array}$$

As always we want to represent this Hopf *algebra $\mathcal{A}_q$ by possibly unbounded operators in some Hilbert space in order to produce a locally compact quantum group in the sense of Definition 1 of the Introduction. In Proposition 2.4 of [9], Korogodsky produces a family of irreducible *-representations of the *-algebra

$\mathcal{A}_q$. We glue part of this family of irreducible *-representations together to a *-representation of $\mathcal{A}_q$ on one Hilbert space that will be the Hilbert space that our locally compact quantum group will act upon. For this purpose we define

$$I_q = \{\, -q^k \mid k \in \mathbb{N} \,\} \cup \{\, q^k \mid k \in \mathbb{Z} \,\}\ .$$

Set $I_q^- = \{\, x \in I_q \mid x < 0 \,\}$ and $I_q^+ = \{\, x \in I_q \mid x > 0 \,\}$. We will use the discrete topology on I_q.
Let $\mathbb{T}$ denote the group of complex numbers of modulus 1. We will consider the uniform measure on I_q and the normalized Haar measure on $\mathbb{T}$. Our *-representation of $\mathcal{A}_q$ will act in the Hilbert space H defined by $H = L^2(\mathbb{T}) \otimes L^2(I_q)$.

We let ξ denote the identity function on I_q, whereas ζ denotes the identity function on $\mathbb{T}$.
If $p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$, we define $\delta_p \in \mathcal{F}(I_q)$ such that $\delta_p(x) = \delta_{x,p}$ for all $x \in I_q$ (note that $\delta_p = 0$ if $p \not\in I_q$). The family $(\, \delta_p \mid p \in I_q \,)$ is the natural orthonormal basis of $L^2(I_q)$. Also recall the natural orthonormal basis $(\, \zeta^m \mid m \in \mathbb{Z} \,)$ for $L^2(\mathbb{T})$.

Instead of looking at the algebra $\mathcal{A}_q$ as the abstract algebra generated by generators and relations we will use an explicit realization of this algebra as linear operators on the dense subspace E of H defined by $E = \langle\, \zeta^m \otimes \delta_x \mid m \in \mathbb{Z}, x \in I_q, \rangle \subseteq H$. Of course, E inherits the inner product from H. Let $\mathcal{L}^+(E)$ denote the *-algebra of adjointable operators on E (see [23, Prop. 2.1.8]), i.e.

$$\mathcal{L}^+(E) = \{\, T \in \mathrm{End}(E) \mid \exists S \in \mathrm{End}(E), \forall v, w \in E : \langle Tv, w\rangle = \langle v, Sw\rangle \,\}\ .$$

The *-operation in $\mathcal{L}^+(E)$ (and $\mathcal{L}^+(E \odot E)$ for that matter) will be denoted by †. So if $T \in \mathcal{L}^+(E)$, the operator $T^\dagger \in \mathcal{L}^+(E)$ is defined to be the operator S in the above definition. It follows that $T^\dagger \subseteq T^*$ where T^* is the usual adjoint of T as an operator in the Hilbert space H. It also follows that T is a closable operator in H.

Define linear operators α_0, γ_0, e_0 on E such that

$$\begin{aligned} \alpha_0(\zeta^m \otimes \delta_p) &= \sqrt{\mathrm{sgn}(p) + p^{-2}}\ \zeta^m \otimes \delta_{qp} \\ \gamma_0(\zeta^m \otimes \delta_p) &= p^{-1}\ \zeta^{m+1} \otimes \delta_p \\ e_0(\zeta^m \otimes \delta_p) &= \mathrm{sgn}(p)\ \zeta^m \otimes \delta_p \end{aligned} \tag{1.3}$$

for all $p \in I_q$, $m \in \mathbb{Z}$.
Then α_0, γ_0 and e_0 belong to $\mathcal{L}^+(E)$, $e_0^\dagger = e_0$ and

$$\begin{aligned} \alpha_0^\dagger\,(\zeta^m \otimes \delta_p) &= \sqrt{\mathrm{sgn}(p) + q^2\, p^{-2}}\ \zeta^m \otimes \delta_{q^{-1}p} \\ \gamma_0^\dagger\,(\zeta^m \otimes \delta_p) &= p^{-1}\ \zeta^{m-1} \otimes \delta_p \end{aligned} \tag{1.4}$$

for all $p \in I_q$, $m \in \mathbb{Z}$. Note that $\alpha_0^\dagger\,(\zeta^m \otimes \delta_{-q}) = 0$.

Then $\mathcal{A}_q$ is the *-subalgebra of $\mathcal{L}^+(E)$ generated by α_0, γ_0 and e_0. Since $\mathcal{L}^+(E) \odot \mathcal{L}^+(E)$ is canonically embedded in $\mathcal{L}^+(E \odot E)$, we obtain $\mathcal{A}_q \odot \mathcal{A}_q$ as a *-subalgebra of $\mathcal{L}^+(E \odot E)$. As such, the comultiplication Δ is, according to Eqs. (1.2), given by

$$\begin{aligned} \Delta_0(\alpha_0) &= \alpha_0 \odot \alpha_0 + q\,(e_0\,\gamma_0^\dagger) \odot \gamma_0 \\ \Delta_0(\gamma_0) &= \gamma_0 \odot \alpha_0 + (e_0 \alpha_0^\dagger) \odot \gamma_0 \\ \Delta_0(e_0) &= e_0 \odot e_0\ . \end{aligned} \tag{1.5}$$

where $\odot$ denotes the algebraic tensor product of linear mappings.

2. The von Neumann algebra underlying quantum $\widetilde{SU}(1,1)$.

In this section we introduce the von Neumann algebra acting on H that underlies the von Neumann algebraic version of the quantum group $\widetilde{SU}_q(1,1)$. The reason for looking first at the von Neumann algebraic picture is twofold. The most important one is the lack of density conditions in the definition of von Neumann algebraic quantum groups (Definition 1 of the introduction), these are automatically satisfied. Moreover, in this case the von Neumann algebra turns out to be very simple. Towards the end of this paper we will use [19, Prop. 1.6] to produce the C*-algebraic version of $\widetilde{SU}_q(1,1)$.

In order to get into the framework of operator algebras, we need to introduce the topological versions of the algebraic objects α_0, γ_0 and e_0 as possibly unbounded operators in the Hilbert space H. So let α denote the closure of α_0, γ the closure of γ_0 and e the closure of e_0, all as linear operators in H. So e is a bounded linear operator on H, whereas α and γ are unbounded, closed, densely defined linear operators in H. Note also that α^* is the closure of $\alpha_0^\dagger$ and that γ^* is the closure of $\gamma_0^\dagger$.

Lemma 2.1. *Let $\mathcal{H}$ be a Hilbert space. Then the domains of $1_{\mathcal{H}}\otimes\alpha$, $1_{\mathcal{H}}\otimes\alpha^*$, $1_{\mathcal{H}}\otimes\gamma$ and $1_{\mathcal{H}}\otimes\gamma^*$ coincide.*

Proof. Since $1_{\mathcal{H}}\otimes\gamma$ is normal, $D(1_{\mathcal{H}}\otimes\gamma^*) = D(1_{\mathcal{H}}\otimes\gamma)$. We know that $\alpha_0^\dagger\alpha_0 = \gamma_0^\dagger\gamma_0 + e_0$, thus $\|(1_{\mathcal{H}}\odot\alpha_0)v\|^2 = \|(1_{\mathcal{H}}\odot\gamma_0)v\|^2 + \|(1_{\mathcal{H}}\otimes e)v\|^2$ for all $v \in \mathcal{H}\odot E$. Since $1_{\mathcal{H}}\otimes e$ is bounded, $1_{\mathcal{H}}\otimes\alpha$ is the closure of $1_{\mathcal{H}}\odot\alpha_0$ and $1_{\mathcal{H}}\otimes\gamma$ is the closure of $1_{\mathcal{H}}\odot\gamma_0$, it follows that $D(1_{\mathcal{H}}\otimes\alpha) = D(1_{\mathcal{H}}\otimes\gamma)$. Using the equality $\alpha_0\alpha_0^\dagger = q^2\,\gamma_0^\dagger\gamma_0 + e_0$, one proves in a similar way that $D(1_{\mathcal{H}}\otimes\alpha^*) = D(1_{\mathcal{H}}\otimes\gamma)$. □

All these operators can easily be realized as a combination of shift and multiplication operators.

Consider $p \in -q^{\mathbb{Z}}\cup q^{\mathbb{Z}}$. We define a translation operator T_p on $\mathcal{F}(\mathbb{T}\times I_q)$ such that for $f \in \mathcal{F}(\mathbb{T}\times I_q)$, $\lambda\in\mathbb{T}$ and $x \in I_q$, we have that $(T_p f)(\lambda,x) = f(\lambda,px)$. By discussion (2) in Notations and conventions, we get that $(T_p f)(\lambda,x) = 0$ if $px \notin I_q$. If $p,t\in I_q$ and $g\in\mathcal{F}(\mathbb{T})$, then $T_p(g\otimes\delta_t) = g\otimes\delta_{p^{-1}t}$, thus, $T_p(g\otimes\delta_t) = 0$ if $p^{-1}t\notin I_q$.

For instance, note that if $f\in\mathcal{F}(\mathbb{T}\times I_q)$, $g\in\mathcal{F}(\mathbb{T})$ and $\lambda\in\mathbb{T}$,

- $(T_{q^{-1}}f)(\lambda,-q) = 0$ and $T_q(g\otimes\delta_{-q}) = 0$,
- $(T_{-1}f)(\lambda,x) = 0$ for all $x\in I_q$ such that $x\geq 1$; $T_{-1}(g\otimes\delta_p) = 0$ for all $p\in I_q$ such that $p \geq 1$.

Notation 2.2. *Define the mapping $\rho : -q^{\mathbb{Z}}\cup q^{\mathbb{Z}} \to B(H)$ such that ρ_p equals the partial isometry on H induced by T_p for all $p\in -q^{\mathbb{Z}}\cup q^{\mathbb{Z}}$. Let us also single out the following special cases:*
1) *Define $w = \rho_{q^{-1}}$, which is an isometry on $B(H)$.*
2) *Define $u = \rho_{-1}$, which is a self-adjoint partial isometry on $B(H)$.*

In terms of multiplication and shift operators, our closed linear operators in H are easily recognized as

$$\alpha = w\,(1\otimes M_{\sqrt{\operatorname{sgn}(\xi)+\xi^{-2}}}) \qquad \gamma = M_\zeta\otimes M_{\xi^{-1}} \qquad e = 1\otimes M_{\operatorname{sgn}\xi}\ . \tag{2.1}$$

These tensor products are obtained by closing the algebraic tensor product mappings with respect to the norm topology on H.

Let us recall the following natural terminology. If $T_1,\ldots,T_n$ are closed, densely defined linear operators in H, the von Neumann algebra N on H generated by $T_1,\ldots,T_n$ is the one such that

$$N' = \{\, x\in B(H) \mid xT_i\subseteq T_i x \text{ and } xT_i^*\subseteq T_i^* x \text{ for } i=1,\ldots,n \,\}\ .$$

Almost by definition, N is the smallest von Neumann algebra acting on H so that $T_1,\ldots,T_n$ are affiliated with M in the von Neumann algebraic sense. If $w_1,\ldots,w_n$ are the partial isometries obtained from the polar decompositions of $T_1,\ldots,T_n$ respectively, then N is also the von Neumann algebra on H generated by

$$\bigcup_{i=1}^{n}\{w_i\}\cup\{\, f(T_i^*T_i) \mid f\in\mathcal{L}^\infty(\sigma(T_i^*T_i))\,\}\ . \tag{2.2}$$

It is now very tempting to define the von Neumann algebra underlying quantum $\widetilde{SU}_q(1,1)$ as the von Neumann algebra generated by α, γ and e. However, for reasons that will become clear later (see the discussion in the beginning of the next section and the remark after Proposition 3.8), the underlying von Neumann algebra will be the one generated by α, γ, e and u.

Definition 2.3. *We define M_q to be the von Neumann algebra on H generated by α, γ, e and u.*

So M_q will be the von Neumann algebra underlying $\widetilde{SU}_q(1,1)$. For convenience, we will also introduce $M_q^\diamond$ as the von Neumann algebra on H generated by α, γ and e.

Lemma 2.4. 1) *$M_q^\diamond$ is generated by $\{w\}\cup\{\, M_f \mid f\in\mathcal{L}^\infty(\mathbb{T}\times I_q)\,\}$,*
2) *M_q is generated by $\{w,u\}\cup\{\, M_f \mid f\in\mathcal{L}^\infty(\mathbb{T}\times I_q)\,\}$,*

3) $M_q = L^\infty(\mathbb{T}) \otimes B(L^2(I_q))$.

Proof. By Eq. (2.2) we know that $M_q^\diamond$ is generated by the set $\{w, M_{\zeta\otimes 1}, M_{1\otimes \mathrm{sgn}(\xi)}\} \cup \{\, f(\alpha^*\alpha) | f \in \mathcal{L}^\infty(\sigma(\alpha^*\alpha))\,\} \cup \{\, f(\gamma^*\gamma) | f \in \mathcal{L}^\infty(\sigma(\gamma^*\gamma))\,\}$. Since $\gamma^*\gamma = 1 \otimes M_{\xi^{-2}}$, it follows that the von Neumann algebra generated by $\{M_{1\otimes \mathrm{sgn}(\xi)}\} \cup \{\, f(\gamma^*\gamma) | f \in \mathcal{L}^\infty(\sigma(\gamma^*\gamma))\,\}$ equals the von Neumann algebra generated by $\{M_{1\otimes \mathrm{sgn}(\xi)}\} \cup \{\, M_{1\otimes f(|\xi|)} | f \in \mathcal{L}^\infty(I_q^+)\,\}$. Thus, the von Neumann algebra generated by $\{M_{1\otimes \mathrm{sgn}(\xi)}\} \cup \{\, f(\gamma^*\gamma) | f \in \mathcal{L}^\infty(\sigma(\gamma^*\gamma))\,\}$ equals $\{\, M_{1\otimes f} \mid f \in \mathcal{L}^\infty(I_q)\,\}$. Because $\alpha^*\alpha = 1 \otimes M_{\mathrm{sgn}(\xi)+\xi^{-2}}$, it now follows that $M_q^\diamond$ is generated by $\{w, M_{\zeta\otimes 1}\} \cup \{\, M_{1\otimes f} \mid f \in \mathcal{L}^\infty(I_q)\,\}$. Therefore the first statement holds, and as a consequence also the second one holds.

Define the partial isometries w_0, u_0 on $L^2(I_q)$ such that $(w_0 f)(x) = f(q^{-1}x)$ and $(u_0 f)(x) = f(-x)$ for all $f \in L^2(I_q)$ and $x \in I_q$. Letting M_0 denote the von Neumann algebra generated by $\{w_0, u_0\} \cup L^\infty(I_q)$, the second statement implies that $M_q = L^\infty(\mathbb{T}) \otimes M_0$. It is not so difficult to see that every rank one projector of the form $L^2(I_q) \to L^2(I_q) : v \mapsto \langle v, \delta_p\rangle\, \delta_t$, where $p, t \in I_q$, belongs to M_0, thus $M_0 = B(L^2(I_q))$. □

So we get in particular that $\rho_p \in M_q$ for all $p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$. Let us collect some further elementary results about the map ρ. If $p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$, we define $i_p = \rho_p^*\rho_p$, the initial projection of ρ_p. Consider $f \in \mathcal{L}^2(\mathbb{T}\times I_q)$ and define $g \in \mathcal{L}^2(\mathbb{T}\times I_q)$ such that for $\lambda \in \mathbb{T}$, $x \in I_q$, we have that $g(\lambda, x) = f(\lambda, x)$ if $p^{-1}x \in I_q$ and $g(\lambda, x) = 0$ if $p^{-1}x \notin I_q$. Then $i_p([f]) = [g]$.

Lemma 2.5. *Consider $p, t \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and $f \in \mathcal{L}^\infty(\mathbb{T}\times I_q)$, then*

$$\rho_p\, \rho_t = \rho_{p\,t}\, i_t \qquad \text{and} \qquad \rho_p^* = \rho_{p^{-1}}$$

and

$$\rho_p\, M_f = M_{T_p f}\, \rho_p \qquad \text{and} \qquad \rho_p\, M_{T_{p^{-1}} f} = M_f\, \rho_p\ .$$

Proof. We only prove the first equality, the other ones are even more straightforward to check. Take $f \in \mathcal{L}^2(\mathbb{T}\times I_q)$ such that $f(\lambda, x) = 0$ for all $\lambda \in \mathbb{T}$ and $x \in I_q$ such that $t^{-1}\,x \notin I_q$. Now $(\rho_p\,\rho_t)([f]) = [T_p(T_t(f))]$, whereas $\rho_{p\,t}([f]) = [T_{p\,t}(f)]$. Let $y \in I_q$ and $\nu \in \mathbb{T}$. We consider 4 different cases:

(1) If $py \in I_q$ and $p\,ty \in I_q$, then $(T_p(T_t(f))(\nu, y) = T_t(f)(\nu, py) = f(\nu, p\,ty)$. Thus $T_{p\,t}(f)(\nu, y) = f(\nu, p\,ty) = (T_p(T_t(f))(\nu, y)$.
(2) Now suppose that $py \notin I_q$ and $p\,ty \in I_q$. Then $(T_p(T_t(f)))(\nu, y) = 0$. On the other hand, $T_{p\,t}(f)(\nu, y) = f(\nu, p\,ty)$. Since $t^{-1}(p\,ty) = py \notin I_q$, we have that $f(\nu, p\,ty) = 0$. Thus $T_{p\,t}(f)(\nu, y) = 0 = (T_p(T_t(f))(\nu, y)$.
(3) Suppose that $py \in I_q$ and $p\,ty \notin I_q$. Then $\big(T_p(T_t(f))\big)(\nu, y) = T_t(f)(\nu, py) = 0$ since $t(py) \notin I_q$. Thus $T_{p\,t}(f)(\nu, y) = 0 = \big(T_p(T_t(f))\big)(\nu, y)$.
(4) If $py \notin I_q$ and $p\,ty \notin I_q$, then clearly $T_{p\,t}(f)(\nu, y) = 0 = \big(T_p(T_t(f))\big)(\nu, y)$.

So we see that $T_p(T_t(f)) = T_{p\,t}(f)$ for this kind of function f. Thus $\rho_p\,\rho_t\, i_t = \rho_{p\,t}\, i_t$. □

So we see that ρ is almost a group representation, but not quite. It is possible to use the results in the previous lemma to perform a partial cross product construction. Since we do not need this approach in the rest of the paper, we will not pursue this matter any further. Instead we focus on a more useful and simpler picture of M_q:

For every $p, t \in I_q$ and $m \in \mathbb{Z}$ we define $\Phi(m, p, t) = \rho_{p^{-1}t}\, M_{\zeta^m \otimes \delta_t} \in M_q$. So if $x \in I_q$ and $r \in \mathbb{Z}$, then

$$\Phi(m, p, t)\, (\zeta^r \otimes \delta_x) = \delta_{x,t}\, \zeta^{m+r} \otimes \delta_p\ . \tag{2.3}$$

Define $M_q^\circ = \langle\, \Phi(m, p, t) \mid m \in \mathbb{Z}, p, t \in I_q\,\rangle$. Using Eq. (2.3), it is easy to see that the family $(\, \Phi(m, p, t) \mid m \in \mathbb{Z}, p, t \in I_q\,)$ is a linear basis of M_q°.

The multiplication and *-operation are easily expressed in terms of these basis elements:

$$\begin{aligned} &\Phi(m_1, p_1, t_1)\, \Phi(m_2, p_2, t_2) = \delta_{p_2, t_1}\ \Phi(m_1 + m_2, p_1, t_2) \\ &\Phi(m, p, t)^* = \Phi(-m, t, p) \end{aligned} \tag{2.4}$$

for all $m, m_1, m_2 \in \mathbb{Z}$, $p, p_1, p_2, t, t_1, t_2 \in I_q$. So we see that M_q° is a σ-weakly dense sub*-algebra of M_q.

3. THE COMULTIPLICATION OF QUANTUM $\widetilde{SU}(1,1)$.

In this section we introduce the comultiplication of $\widetilde{SU}_q(1,1)$. In the first part we start with a motivation for the formulas appearing in Definition 3.1. Although the discussion is not really needed in the build up of $\widetilde{SU}_q(1,1)$, it is important and clarifying to know how we arrived at the formulas in Definition 3.1.

But first we introduce two auxiliary functions

(1) $\chi : -q^{\mathbb{Z}} \cup q^{\mathbb{Z}} \to \mathbb{Z}$ such that $\chi(x) = \log_q(|x|)$ for all $x \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$,

(2) $\kappa : \mathbb{R} \to \mathbb{R}$ such that $\kappa(x) = \operatorname{sgn}(x)\, x^2$ for all $x \in \mathbb{R}$.

Our purpose is to define a comultiplication $\Delta : M_q \to M_q \otimes M_q$. Assume for the moment that this has already been done. It is natural to require Δ to be closely related to the comultiplication Δ_0 on $\mathcal{A}_q$ as defined in Eqs. (1.5). The least that we expect is $\Delta_0(T_0) \subseteq \Delta(T)$ and $\Delta_0(T_0^\dagger) \subseteq \Delta(T)^*$ for $T = \alpha, \gamma$. In the rest of this discussion we will focus on the inclusion $\Delta_0(\gamma_0^\dagger \gamma_0) \subseteq \Delta(\gamma^*\gamma)$, where $\Delta_0(\gamma_0^\dagger\gamma_0) \in \mathcal{L}^+(E \odot E)$. Because $\gamma^*\gamma$ is self-adjoint, the element $\Delta(\gamma^*\gamma)$ would also be self-adjoint. So the hunt is on for self-adjoint extensions of the explicit operator $\Delta_0(\gamma_0^\dagger\gamma_0)$. Unlike in the case of quantum $E(2)$ (see [27]), the operator $\Delta_0(\gamma_0^\dagger\gamma_0)$ is not essentially self-adjoint. But it was already known in [9] that $\Delta_0(\gamma_0^\dagger\gamma_0)$ has self-adjoint extensions (this follows easily because the operator in (3.2) commutes with complex conjugation, implying that the deficiency spaces are isomorphic).

Let us make a small detour to quantum $SU(1,1)$. In this case, the operators α_0 and γ_0 are replaced by their restrictions to $L^2(\mathbb{T} \otimes I_q^+)$. Then $\Delta_0(\gamma_0^\dagger\gamma_0)$ still has self-adjoint extensions for each of which the domain is obtained by imposing a boundary condition on functions in its domain (this boundary condition is a simple relation between the function and its Jackson derivative in the limit towards 0, see [9, Eq. (6.2)]). However, in this case the closure of the operator $\Delta_0(\alpha_0)$ does not leave the domain of such a self-adjoint extension invariant because it distorts the boundary condition. From a practical point of view, it should also be said that an explicit manageable spectral decomposition in terms of special functions for these self-adjoint extensions is missing, cf. [13, Rem. 2.7].

Now we return to the case of quantum $\widetilde{SU}(1,1)$. Although $\Delta_0(\gamma_0^\dagger\gamma_0)$ has a self-adjoint extension, it is not unique. We have to make a choice for this self-adjoint extension, but we cannot extract the information necessary to make this choice from α and γ alone. This is why we do not work with $M_q^\diamond$ but with M_q which has the above extra extension information contained in the element u. These kind of considerations were already present in [31] and were also introduced in [30] for quantum $\widetilde{SU}_q(1,1)$. In this paper, this principle is only lurking in the background but it is treated in a fundamental and rigorous way in [30].

Now we get into slightly more detail in our discussion about the extension of $\Delta_0(\gamma_0^\dagger\gamma_0)$. Define a linear map $L : \mathcal{F}(\mathbb{T} \times I_q \times \mathbb{T} \times I_q) \to \mathcal{F}(\mathbb{T} \times I_q \times \mathbb{T} \times I_q)$ such that

$$\begin{aligned}(Lf)(\lambda, x, \mu, y) \quad = \quad & [\, x^{-2}(\operatorname{sgn}(y) + y^{-2}) + (\operatorname{sgn}(x) + q^2\, x^{-2})\, y^{-2}\,]\, f(\lambda, x, \mu, y) \\ & + \operatorname{sgn}(x)\, q^{-1}\, \bar{\lambda}\mu\, x^{-1} y^{-1} \sqrt{(\operatorname{sgn}(x) + x^{-2})(\operatorname{sgn}(y) + y^{-2})}\, f(\lambda, qx, \mu, qy) \\ & + \operatorname{sgn}(x)\, q\, \lambda\bar{\mu}\, x^{-1} y^{-1} \sqrt{(\operatorname{sgn}(x) + q^2 x^{-2})(\operatorname{sgn}(y) + q^2 y^{-2})}\, f(\lambda, q^{-1}x, \mu, q^{-1}y)\end{aligned}$$

for all $\lambda, \mu \in \mathbb{T}$ and $x, y \in I_q$. A straightforward calculation reveals that if $f \in E \odot E$, then $\Delta_0(\gamma_0^\dagger\gamma_0)\,[f] = [L(f)]$. From this, it is a standard exercise to check that $[f] \in D(\Delta_0(\gamma_0^\dagger\gamma_0)^*)$ and $\Delta_0(\gamma_0^\dagger\gamma_0)^*[f] = [L(f)]$ if $f \in \mathcal{L}^2(\mathbb{T} \times I_q \times \mathbb{T} \times I_q)$ and $L(f) \in \mathcal{L}^2(\mathbb{T} \times I_q \times \mathbb{T} \times I_q)$ (without any difficulty, one can even show that $D(\Delta_0(\gamma_0^\dagger\gamma_0)^*)$ consists precisely of such elements $[f]$).

If $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$, we define $\ell'_\theta = \{\, (\lambda, x, \mu, y) \in \mathbb{T} \times I_q \times \mathbb{T} \times I_q \mid y = \theta x \,\}$. We consider $L^2(\ell'_\theta)$ naturally embedded in $L^2(\mathbb{T} \times I_q \times \mathbb{T} \times I_q)$. It follows easily from the above discussion that $\Delta_0(\gamma_0^\dagger\gamma_0)^*$ leaves $L^2(\ell'_\theta)$ invariant. Thus, if T is a self-adjoint extension of $\Delta_0(\gamma_0^\dagger\gamma_0)$, the obvious inclusion $T \subseteq \Delta_0(\gamma_0^\dagger\gamma_0)^*$ implies that T also leaves $L^2(\ell'_\theta)$ invariant.

Therefore every self-adjoint extension T of $\Delta_0(\gamma_0^\dagger\gamma_0)$ is obtained by choosing a self-adjoint extension T_θ of the restriction of $\Delta_0(\gamma_0^\dagger\gamma_0)$ to $L^2(\ell'_\theta)$ for every $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and setting $T = \oplus_{\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}} T_\theta$.

Therefore fix $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$. Define $J_\theta = \{\, z \in I_{q^2} \mid \kappa(\theta)\, z \in I_{q^2} \,\}$ which is a q^2-interval around 0. On J_θ we define a measure ν_θ such that $\nu_\theta(\{x\}) = |x|$ for all $x \in J_\theta$.

Now define the unitary transformation $U_\theta : L^2(\mathbb{T} \times \mathbb{T} \times J_\theta) \to L^2(\ell'_\theta)$ such that $U_\theta([f]) = [g]$ where $f \in \mathcal{L}^2(\mathbb{T} \times \mathbb{T} \times J_\theta)$ and $g \in \mathcal{L}^2(\ell'_\theta)$ are such that

$$g(\lambda, z, \mu, \theta z) = (\lambda \bar{\mu})^{\chi(z)} \, (-\mathrm{sgn}(\theta z))^{\chi(z)} \, |z| \, f(\lambda, \mu, \kappa(z)) \tag{3.1}$$

for all $\lambda, \mu \in \mathbb{T}$ and $z \in I_q$ such that $\theta z \in I_q$.
Define the linear operator $L_\theta : \mathcal{F}(J_\theta) \to \mathcal{F}(J_\theta)$ such that

$$\begin{aligned}(L_\theta f)(x) \quad = \quad & \frac{1}{\theta^2 x^2} \big(-\sqrt{(1+x)(1+\kappa(\theta)\,x)}\, f(q^2 x) - q^2 \sqrt{(1+q^{-2}x)(1+q^{-2}\kappa(\theta)\,x)}\, f(q^{-2}x) \\ & \qquad [(1+\kappa(\theta)\,x) + q^2(1+q^{-2}x)]\, f(x) \big)\end{aligned} \tag{3.2}$$

for all $f \in \mathcal{F}(J_\theta)$ and $x \in J_\theta$.
Then an easy calculation shows that $U_\theta^* \, \Delta_0(\gamma_0^\dagger \gamma_0)\restriction_{E \odot E} \; U_\theta = 1 \odot (L_\theta \restriction_{\mathcal{K}(J_\theta)})$. So our problem is reduced to finding self-adjoint extensions of $L_\theta \restriction_{\mathcal{K}(I_q)}$. This operator $L_\theta \restriction_{\mathcal{K}(I_q)}$ is a second order q-difference operator for which eigenfunctions in terms of q-hypergeometric functions are known.

We can use a reasoning similar to the one in [13, Sec. 2] to get hold of the self-adjoint extensions of $L_\theta \restriction_{\mathcal{K}(I_q)}$:
Let $\beta \in \mathbb{T}$. Then we define a linear operator $L_\theta^\beta : D(L_\theta^\beta) \subseteq L^2(J_\theta, \nu_\theta) \to L^2(J_\theta, \nu_\theta)$ such that

$$D(L_\theta^\beta) = \{\, f \in L^2(J_\theta, \nu_\theta) \mid L_\theta(f) \in L^2(J_\theta, \nu_\theta), f(0+) = \beta\, f(0-) \text{ and } (D_q f)(0+) = \beta\, (D_q f)(0-) \,\}$$

and L_θ^β is the restriction of L_θ to $D(L_\theta^\beta)$. Here, D_q denotes the Jackson derivative, that is, $(D_q f)(x) = (f(qx) - f(x))/(q-1)x$ for $x \in J_\theta$. Also, $f(0+) = \beta\, f(0-)$ is an abbreviated form of saying that the limits $\lim_{x \uparrow 0} f(x)$ and $\lim_{x \downarrow 0} f(x)$ exist and $\lim_{x \downarrow 0} f(x) = \beta \lim_{x \uparrow 0} f(x)$.
Then L_θ^β is a self-adjoint extension of $L_\theta \restriction_{\mathcal{K}(I_q)}$. If $\beta, \beta' \in \mathbb{T}$ and $\beta \neq \beta'$, then $L_\theta^\beta \neq L_\theta^{\beta'}$.

It is tempting to use the extension L_θ^1 to construct our final self-adjoint extension for $\Delta_0(\gamma_0^\dagger \gamma_0)$ (although there is no apparent reason for this choice). However, in order to obtain a coassociative comultiplication, it turns out that we have to use the extension $L_\theta^{\mathrm{sgn}(\theta)}$ to construct our final self-adjoint extension. This is reflected in the fact that the expression $s(x, y)$ appears in the formula for a_p in Definition 3.1.

This all would be only a minor achievement if we could not go any further. But the results and techniques used in the theory of q-hypergeometric functions will even allow us to find an explicit orthonormal basis consisting of eigenvectors of $L_\theta^{\mathrm{sgn}(\theta)}$. These eigenvectors are, up to a unitary transformation, obtained by restricting the functions a_p in Definition 3.1 to ℓ_θ, which is introduced after this definition. The special case $\theta = 1$ was already known to Korogodsky (see [9, Prop. A.1]), although the proof in [9] seems to contain quite a few gaps. For instance, the presentation in [3] shows that a lot of more care has to be taken to solve these kind of problems.

In order to compress the formulas even further, we introduce two other auxiliary functions

(1) $\nu : -q^{\mathbb{Z}} \cup q^{\mathbb{Z}} \to \mathbb{R}^+$ such that $\nu(t) = q^{\frac{1}{2}(\chi(t)-1)(\chi(t)-2)}$ for all $t \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$.

(2) Another auxiliary function $s : \mathbb{R}_0 \times \mathbb{R}_0 \to \{-1, 1\}$ is defined such that

$$s(x, y) = \begin{cases} -1 & \text{if } x > 0 \text{ and } y < 0 \\ 1 & \text{if } x < 0 \text{ or } y > 0 \end{cases}$$

for all $x, y \in \mathbb{R}_0$.

Let us also collect some basic manipulation rules. Therefore take $x, y, z \in \mathbb{R}_0$.

(1) If $x > 0$ then $s(x, y) = \mathrm{sgn}(yz)\, s(x, z)$ and $s(y, x) = s(z, x)$. If $x < 0$, then $s(x, y) = s(x, z)$ and $s(y, x) = \mathrm{sgn}(yz)\, s(z, x)$. So if $\mathrm{sgn}(xyz) = 1$, then $s(x, y) = s(x, z)$.

(2) It is clear that $s(x, y) = -\mathrm{sgn}(x)\, s(x, -y)$ and $s(x, y) = \mathrm{sgn}(y)\, s(-x, y)$.

(3) One easily checks that $s(x, y) = \mathrm{sgn}(xy)\, s(y, x)$. As a consequence we get also that $s(-x, y) = s(-y, x)$ and $s(x, -y) = s(y, -x)$.

If $a, b, z \in \mathbb{C}$, we define $\Psi\left(\begin{matrix} a \\ b \end{matrix} ; z\right)$ and $\Psi(a;b;z)$ to be equal to $\Psi\left(\begin{matrix} a \\ b \end{matrix} ; q^2; z\right)$.

We will also use the normalization constant $c_q = (\sqrt{2}\, q\, (q^2, -q^2; q^2)_\infty)^{-1}$.

Definition 3.1. *If $p \in I_q$, we define a function $a_p : I_q \times I_q \to \mathbb{R}$ such that a_p is supported on the set $\{\, (x,y) \in I_q \times I_q \mid \mathrm{sgn}(xy) = \mathrm{sgn}(p) \,\}$ and $a_p(x,y)$ is given by*

$$c_q\, s(x,y)\, (-1)^{\chi(p)}\, (-\mathrm{sgn}(y))^{\chi(x)}\, |y|\, \nu(py/x)\, \sqrt{\frac{(-\kappa(p), -\kappa(y); q^2)_\infty}{(-\kappa(x); q^2)_\infty}}\; \Psi\left(\begin{matrix} -q^2/\kappa(y) \\ q^2\kappa(x/y) \end{matrix} ; q^2\kappa(x/p)\right)$$

for all $(x,y) \in I_q \times I_q$ satisfying $\mathrm{sgn}(xy) = \mathrm{sgn}(p)$.

The presence of the expression $(-\mathrm{sgn}(y))^{\chi(x)}$ in this same formula can be traced back to the defining formula (3.1) for U_θ.

The extra vital information that we need is contained in the following proposition. For $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ we define $\ell_\theta = \{\, (x,y) \in I_q \times I_q \mid y = \theta x \,\}$.

Proposition 3.2. *Consider $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$. Then the family $(\, a_p\restriction_{\ell_\theta} \mid p \in I_q$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta)\,)$ is an orthonormal basis for $l^2(\ell_\theta)$.*

This result implies also a dual result, stemming from the following simple duality principle (in special function theory, these are referred to as dual orthogonality relations). Consider a set I and suppose that $l^2(I)$ has an orthonormal basis $(e_j)_{j\in J}$. For every $i \in I$, we define a function f_i on J by $f_i(j) = e_j(i)$. Then $(f_i)_{i\in I}$ is an orthonormal basis for $l^2(J)$. If we apply this principle to the line ℓ_θ, the previous proposition implies the next one.

Proposition 3.3. *Consider $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and define $J = I_q^+$ if $\theta > 0$ and $J = I_q^-$ if $\theta < 0$. For every $(x,y) \in \ell_\theta$ we define the function $e_{(x,y)} : J \to \mathbb{R}$ such that $e_{(x,y)}(p) = a_p(x,y)$ for all $p \in J$. Then the family $(\, e_{(x,y)} \mid (x,y) \in \ell_\theta\,)$ forms an orthonormal basis for $l^2(J)$.*

The first of these two propositions will be shortly used to define the comultiplication on M_q, both of them pop up in the proof of the left invariance of the Haar weight.

For the proofs of Propositions 3.2 and 3.3 we refer to the literature. For $1 \geq \theta > 0$ Proposition 3.3 is [3, Thm. 4.1] and Proposition 3.2 follows from Corollary 4.2 and the following remark of [3] in base q^2 and with $(c, q^{2\alpha})$ replaced by $(1, \kappa(\theta))$. For $\theta \geq 1$ we derive Propositions 3.2 and 3.3 similarly from [3, Thm. 4.1 and Cor. 4.2] in base q^2 and with $(c, q^{2\alpha})$ replaced by $(1, \kappa(\theta)^{-1})$. The case $\theta \geq 1$ can also be reduced to the case $0 < \theta \leq 1$ using the second symmetry property of Proposition 3.5. The functions $a_p(x,y)\restriction_{\ell^\theta}$ are eigenfunctions of the operator L_θ as considered in (3.2). The general theory used in the first half of [3] is explained in [12]. For $\theta < 0$ the statements in Propositions 3.2 and 3.3 reduce to statements on specific classes of orthogonal polynomials. For $\theta < 0$ the orthogonality relations in Proposition 3.2 are directly obtainable from the orthogonality relations for the Al-Salam and Carlitz polynomials $U_n^{(a)}(x;q)$ in base q^2 with a replaced by $\kappa(\theta)$, see Al-Salam's and Carlitz's original paper [2] or references in [11]. This time we have that $a_p(x,y)\restriction_{\ell^\theta}$ are eigenfunctions of the operator L_θ, due to the second order q-difference equation for the Al-Salam and Carlitz polynomials, see [11]. The dual result in Proposition 3.3, which follows immediately from Proposition 3.2 for $\theta < 0$ since the corresponding moment problem is determinate, can also be matched to orthogonal polynomials after applying elementary transformation formulas for basic hypergeometric series. The dual orthogonality relations split up in three summation results. Two of these summations can be matched to the orthogonality relations for the q-Charlier polynomials given in [11, Eq. (3.23.3)], but for different parameters. The remaining summation is an easy consequence of [6, Eq. (1.3.16)].

The functions $a_p(x,y)$ can be thought of as Clebsch-Gordan coefficients for the tensor product decomposition of the representation described in (2.1) with itself. Note that the corresponding Clebsch-Gordan coefficients for the quantum $SU(2)$ group are given in terms of Wall polynomials, see Koornwinder [15, Rem. 4.2], and we can write the Wall polynomials in terms of the functions $a_p(x,y)$, precisely for $y \in -q^{-\mathbb{N}_0}$, using [6, Eq. (III.3)].

There is a nice symmetry in $a_p(x,y)$ with respect to interchanging x, y and p. One of this symmetries is easy to see while the proof of the other one requires an extra lemma.

Lemma 3.4. *Consider $x,y,p \in I_{q^2}$ and $m \in \mathbb{Z}$ such that $y/x = \operatorname{sgn}(p)\, q^{2m}$. Then*

$$\sqrt{\frac{(-y;q^2)_\infty}{(-x;q^2)_\infty}}\; \Psi\left(\begin{array}{c} -q^2/y \\ q^2x/y \end{array} ; q^2x/p\right) = \operatorname{sgn}(p)^{\log_{q^2}(|p|)+1}\,(-q^2/p)^m\, \sqrt{\frac{(-x;q^2)_\infty}{(-y;q^2)_\infty}}\; \Psi\left(\begin{array}{c} -q^2/x \\ q^2y/x \end{array} ; q^2y/p\right) .$$

Proof. If $p>0$, this follows easily from Proposition 6.6(1). Now assume that $p<0$. By Result 6.4, the above equation is equivalent to

$$\Psi\left(\begin{array}{c} -q^2/p \\ q^2x/p \end{array} ; q^2x/y\right) = \operatorname{sgn}(p)^{\log_{q^2}(|p|)+1}\,(-q^2/p)^m\, \frac{(-x;q^2)_\infty}{(-y;q^2)_\infty}\; \Psi\left(\begin{array}{c} -q^2/p \\ q^2y/p \end{array} ; q^2y/x\right) .$$

This equality follows easily from Proposition 6.6(2). □

With this lemma in hand, one verifies the second of the next symmetries. Use Result 6.4 to prove the first symmetry, the third one is then a consequence of the previous two symmetries. For the first two symmetries you will also need two of the manipulation rules for $s(x,y)$ discussed before Definition 3.1, namely the last one of (1) and the first one of (3).

Proposition 3.5. *If $x,y,p \in I_q$, then*

$$\begin{aligned} a_p(x,y) &= (-1)^{\chi(yp)}\operatorname{sgn}(x)^{\chi(x)}\;|y/p|\; a_y(x,p) \\ a_p(x,y) &= \operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(x)^{\chi(x)}\operatorname{sgn}(y)^{\chi(y)}\, a_p(y,x) \\ a_p(x,y) &= (-1)^{\chi(xp)}\operatorname{sgn}(y)^{\chi(y)}\,|x/p|\, a_x(p,y) \ . \end{aligned}$$

Now we produce the eigenvectors of our self-adjoint extension of $\Delta_0(\gamma_0^\dagger\gamma_0)$ (see the remarks after the proof of proposition 3.9). We will use these eigenvectors to define a unitary operator that will induce the comultiplication. The presence of $\lambda^{\chi(y)}$ and $\mu^{\chi(x)}$ in the formulas for $\mathcal{F}_{r,s,m,p}$ can be traced back to Eq. (3.1). The dependence of $\mathcal{F}_{r,s,m,p}$ on r,s and p is chosen in such a way that Proposition 3.9 is true.

Definition 3.6. *Consider $r,s \in \mathbb{Z}$, $m \in \mathbb{Z}$ and $p \in I_q$. We define the element $\mathcal{F}_{r,s,m,p} \in H \otimes H$ such that*

$$\mathcal{F}_{r,s,m,p}(\lambda,x,\mu,y) = \begin{cases} a_p(x,y)\,\lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)} & \text{if } y = \operatorname{sgn}(p)\,q^m\,x \\ 0 & \text{otherwise} \end{cases}$$

for all $x,y \in I_q$ and $\lambda,\mu \in \mathbb{T}$.

We know from Proposition 3.2 that the family $(\,\mathcal{F}_{r,s,m,p} \mid r,s \in \mathbb{Z}, m \in \mathbb{Z}, p \in I_q\,)$ forms an orthonormal basis for $H \otimes H$. As a consequence, we get the following result.

Proposition 3.7. *If $x,y \in I_q$ and $r,s \in \mathbb{Z}$, then, using L^2-convergence,*

$$\zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y = \sum_{p \in I_q} a_p(x,y)\; \mathcal{F}_{r-\chi(y/p),s+\chi(x/p),\chi(y/x),p} \ .$$

Now we are ready to introduce the comultiplication of quantum $\widetilde{SU}_q(1,1)$.

Proposition 3.8. *Define the unitary transformation $V : H \otimes H \to L^2(\mathbb{T}) \otimes L^2(\mathbb{T}) \otimes H$ such that $V(\mathcal{F}_{r,s,m,p}) = \zeta^r \otimes \zeta^s \otimes \zeta^m \otimes \delta_p$ for all $r,s \in \mathbb{Z}$, $m \in \mathbb{Z}$ and $p \in I_q$. Then there exists a unique injective normal *-homomorphism $\Delta : M_q \to M_q \otimes M_q$ such that $\Delta(a) = V^*(1_{L^2(\mathbb{T})} \otimes 1_{L^2(\mathbb{T})} \otimes a)V$ for all $a \in M_q$.*

Proof. Define the *-homomorphism $\Delta : M_q \to B(H \otimes H)$ such that $\Delta(a) = V^*(1_{L^2(\mathbb{T})} \otimes 1_{L^2(\mathbb{T})} \otimes a)V$ for all $a \in M_q$. Fix $a \in M_q$. If $r, s, m \in \mathbb{Z}$, $p \in I_q$, then Definition 3.6 implies that $(1_H \otimes M_\zeta \otimes 1_{L^2(I_q)})\mathcal{F}_{r,s,m,p} = \mathcal{F}_{r,s+1,m,p}$. It follows that $V(1_H \otimes M_\zeta \otimes 1_{L^2(I_q)}) = (1_{L^2(\mathbb{T})} \otimes M_\zeta \otimes 1_H)V$. As a consequence, $V(1_H \otimes b \otimes 1_{L^2(I_q)}) = (1_{L^2(\mathbb{T})} \otimes b \otimes 1_H)V$ for all $b \in L^\infty(\mathbb{T})$. It follows that $\Delta(a)(1_H \otimes b \otimes 1_{L^2(I_q)}) = (1_H \otimes b \otimes 1_{L^2(I_q)})\Delta(a)$ for all $b \in L^\infty(\mathbb{T})$. Thus, $\Delta(a) \in (1_H \otimes L^\infty(\mathbb{T}) \otimes 1_{L^2(I_q)})' = B(H) \otimes L^\infty(\mathbb{T}) \otimes B(L^2(I_q)) = B(H) \otimes M_q$ by Lemma 2.4. In a similar way, $\Delta(a) \in M_q \otimes B(H)$. Hence $\Delta(a) \in M_q \otimes M_q$ by the commutator theorem for tensor products. □

The requirement that $\Delta(M_q) \subseteq M_q \otimes M_q$ is the primary reason for introducing the extra generator u. We cannot work with the von Neumann algebra $M_q^\diamond$ that is generated only by α, γ and e, because $\Delta(M_q^\diamond) \nsubseteq M_q^\diamond \otimes M_q^\diamond$.

This definition of Δ and Eq. (2.1) imply easily that $\langle\, \mathcal{F}_{r,s,m,p} \mid r, s \in \mathbb{Z}, m \in \mathbb{Z}, p \in I_q \,\rangle$ is a core for $\Delta(\alpha)$, $\Delta(\gamma)$ and

$$\begin{aligned} \Delta(\alpha)\, \mathcal{F}_{r,s,m,p} &= \sqrt{\operatorname{sgn}(p) + p^{-2}}\; \mathcal{F}_{r,s,m,pq} \\ \Delta(\gamma)\, \mathcal{F}_{r,s,m,p} &= p^{-1}\; \mathcal{F}_{r,s,m+1,p}\ . \end{aligned} \tag{3.3}$$

for $r, s \in \mathbb{Z}$, $m \in \mathbb{Z}$ and $p \in I_q$.

Since $\mathcal{F}_{r,s,m,p}(\lambda, x, \mu, y) = 0$ if $\operatorname{sgn}(x)\operatorname{sgn}(y) \neq \operatorname{sgn}(p)$ it follows that $(e \otimes e)F_{r,s,m,p} = \operatorname{sgn}(p)\, F_{r,s,m,p}$. Hence $V(e \otimes e) = (1_{L^2(\mathbb{T}^2)} \otimes e)V$. As a consequence, $\Delta(e) = e \otimes e$.

Recall the linear operators $\Delta_0(\alpha_0)$, $\Delta_0(\gamma_0)$ acting on $E \odot E$ (Eqs. (1.5)). Also recall the distinction between * and $\dagger$. Since $\Delta(\alpha) = V^*(1_{L^2(\mathbb{T})} \otimes 1_{L^2(\mathbb{T})} \otimes \alpha)V$ and $\Delta(\gamma) = V^*(1_{L^2(\mathbb{T})} \otimes 1_{L^2(\mathbb{T})} \otimes \gamma)V$, Lemma 2.1 implies that $\Delta(\alpha)$ and $\Delta(\gamma)$ are balanced. For the proof of the next proposition q-contiguous relations for q-hypergeometric functions are essential (see Lemma 6.5).

Proposition 3.9. *The following inclusions hold:* $\Delta_0(\alpha_0) \subseteq \Delta(\alpha)$, $\Delta_0(\alpha_0)^\dagger \subseteq \Delta(\alpha)^*$, $\Delta_0(\gamma_0) \subseteq \Delta(\gamma)$ *and* $\Delta_0(\gamma_0)^\dagger \subseteq \Delta(\gamma)^*$.

Proof. (1) The first step is to prove for $i, j, m \in \mathbb{Z}$, $p \in I_q$ and $r, s \in \mathbb{Z}$, $x, y \in I_p$,

$$\langle \Delta(\alpha)\mathcal{F}_{i,j,m,p}, \zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y \rangle = \langle \mathcal{F}_{i,j,m,p}, \Delta_0(\alpha_0^\dagger)(\zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y) \rangle\ . \tag{3.4}$$

and this boils down to the q-contiguous relations in Lemma 6.5. First we can rewrite Eq. (6.2) in terms of the functions a_p of Definition 3.1 by a straightforward verification as

$$\begin{aligned} &\sqrt{\operatorname{sgn}(p) + p^{-2}}\, a_{qp}(x, y) \\ &\quad = \sqrt{(\operatorname{sgn}(x) + q^2/x^2)(\operatorname{sgn}(y) + q^2/y^2)}\, a_p(q^{-1}x, q^{-1}y) + \operatorname{sgn}(x)\,(q/xy)\, a_p(x, y)\ . \end{aligned} \tag{3.5}$$

Here we apply Eq. (6.2) in base q^2 with a,b,z replaced by $-q^2/\kappa(y)$, $q^2\kappa(x/y)$, $\kappa(x/p)$ respectively (and take some care if $x = -q$ or $y = -q$).

Now consider the left hand side of Eq. (3.4). By Eq. (3.3) and Definition 3.6, the inner product on the left hand side of Eq. (3.4) is zero unless $|y/x| = q^m$, $\operatorname{sgn}(p) = \operatorname{sgn}(xy)$, $i + \chi(y/qp) = r$ and $j - \chi(x/qp) = s$ in which case, this inner product equals $\sqrt{\operatorname{sgn}(p) + p^{-2}}\, a_{qp}(x, y)$. (3.6)
For the right hand side of Eq. (3.4) we recall that $\Delta_0(\alpha_0^\dagger) \in \mathcal{L}^+(E \odot E)$ is given by $\Delta_0(\alpha_0^\dagger) = q\, e_0\gamma_0 \odot \gamma_0^\dagger + \alpha_0^\dagger \odot \alpha_0^\dagger$. From Eqs. (1.3) and (1.4), it follows that the right hand side of Eq. (3.4) equals

$$\begin{aligned} &\operatorname{sgn}(x)\,(q/xy)\, \langle \mathcal{F}_{i,j,m,p}, \zeta^{r+1} \otimes \delta_x \otimes \zeta^{s-1} \otimes \delta_y \rangle \\ &\quad + \sqrt{(\operatorname{sgn}(x) + q^2/x^2)(\operatorname{sgn}(y) + q^2/y^2)}\, \langle \mathcal{F}_{i,j,m,p}, \zeta^r \otimes \delta_{q^{-1}x} \otimes \zeta^s \otimes \delta_{q^{-1}y} \rangle\ . \end{aligned} \tag{3.7}$$

By Definition 3.6, this implies that the right hand side of Eq. (3.4) is zero unless $|y/x| = q^m$, $\operatorname{sgn}(p) = \operatorname{sgn}(xy)$, $i + \chi(y/qp) = r$ and $j - \chi(x/qp) = s$, which proves Eq. (3.4) if these conditions are violated. If, on the other hand, these conditions are satisfied, Definition 3.6 guarantees that (3.7) equals

$$\operatorname{sgn}(x)\,(q/xy)\, a_p(x, y) + \sqrt{(\operatorname{sgn}(x) + q^2/x^2)(\operatorname{sgn}(y) + q^2/y^2)}\, a_p(q^{-1}x, q^{-1}y)\ .$$

Therefore (3.6) and Eq. (3.5) imply Eq. (3.4) also in this case.

So we have established Eq. (3.4) in all possible cases. Since the elements $\digamma_{i,j,m,p}$ form a core for $\Delta(\alpha)$, we conclude that $\zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y \in D(\Delta(\alpha)^*)$ and $\Delta(\alpha)^*(\zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y) = \Delta_0(\alpha_0^\dagger)(\zeta^r \otimes \delta_x \otimes \zeta^s \otimes \delta_y)$, thus proving that $\Delta_0(\alpha_0^\dagger) \subseteq \Delta(\alpha)^*$. Taking the adjoint of this equation, we see that $\Delta(\alpha) \subseteq \Delta_0(\alpha_0^\dagger)^*$. In general, $\Delta_0(\alpha_0) \subseteq \Delta_0(\alpha_0^\dagger)^*$. Since $D(\Delta(\alpha)) = D(\Delta(\alpha)^*) \supseteq E \odot E$, we conclude that the inclusion $\Delta_0(\alpha_0) \subseteq \Delta(\alpha)$ also holds.

(2) The inclusions regarding $\Delta(\gamma)$ and $\Delta(\gamma)^*$ are treated in the same way but this time one applies Eq. (6.3) in base q^2 with a,b,z replaced by $-q^2/\kappa(y)$, $q^2\kappa(x/y)$, $q^2\kappa(x/p)$ respectively, to obtain

$$p^{-1}\, a_p(x,y) = \operatorname{sgn}(x)\, y^{-1} \sqrt{\operatorname{sgn}(x) + 1/x^2}\; a_p(qx,y) + x^{-1} \sqrt{\operatorname{sgn}(y) + q^2/y^2}\; a_p(x, q^{-1}y)$$

for all $x, y, p \in I_q$. ☐

This proposition implies also that $\Delta(\gamma^*\gamma)$ is an extension of $\Delta_0(\gamma_0^\dagger\gamma_0)$. We also know that $\langle\, \digamma_{r,s,m,p} \mid r,s \in \mathbb{Z}, m \in \mathbb{Z}, p \in I_q \,\rangle$ is a core for $\Delta(\gamma^*\gamma)$ and $\Delta(\gamma^*\gamma)\, \digamma_{r,s,m,p} = p^{-2}\, \digamma_{r,s,m,p}$ for $r,s,m \in \mathbb{Z}$, $p \in I_q$. Using this information it is not so difficult to check that $U_\theta^* \Delta(\gamma^*\gamma)\restriction_{\ell'_\theta} U_\theta = 1 \otimes L_\theta^{\operatorname{sgn}(\theta)}$ for all $\theta \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$, but we will not make any use of this fact in this paper.

In the next step we investigate the behavior of V with respect to the flip map $\Sigma : H \otimes H \to H \otimes H$. Later on, this will guarantee that the unitary antipode (see the discussion after the proof of [19, Prop. 1.4]) commutes with the comultiplication up to the flip map. It will also reduce some of the calculations in the proof of the coassociativity. This behavior is directly related to the second symmetry property of Proposition 3.5.

Let us define anti-unitary permutation operators $\Sigma' : L^2(\mathbb{T}^2) \to L^2(\mathbb{T}^2)$ and $\tilde{\Sigma} : L^2(\mathbb{T}^4) \to L^2(\mathbb{T}^4)$ such that $\Sigma'(v_1 \otimes v_2) = \bar{v}_2 \otimes \bar{v}_1$ and $\tilde{\Sigma}(v_1 \otimes v_2 \otimes v_3 \otimes v_4) = \bar{v}_4 \otimes \bar{v}_3 \otimes \bar{v}_2 \otimes \bar{v}_1$ for all $v_1, v_2, v_3, v_4 \in L^2(\mathbb{T})$.

We also introduce the anti-unitary operator $\breve{J}$ on H such that $\breve{J}[f] = [g]$, where $f, g \in \mathcal{L}^2(\mathbb{T} \times I_q)$ are such that $g(\lambda, x) = \operatorname{sgn}(x)^{\chi(x)}\, \overline{f(\lambda,x)}$ for all $x \in I_q$ and $\lambda \in \mathbb{T}$.

Proposition 3.10. *We have that* $V\Sigma = (\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})$ *and*

$$V_{13}(1_H \otimes V)\Sigma_{13} = (\tilde{\Sigma} \otimes \breve{J})(1_{L^2(\mathbb{T}^2)} \otimes V)(V \otimes 1_H)(\breve{J} \otimes \breve{J} \otimes \breve{J}) \ .$$

Proof. Take $r,s,m \in \mathbb{Z}$ and $p \in I_q$. Proposition 3.5 implies that $a_p(x,y) = \operatorname{sgn}(p)^{\chi(p)} \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\, a_p(y,x)$ for all $x, y \in I_q$. If $x, y \in I_q$ and $\lambda, \mu \in \mathbb{T}$, then Definition 3.6 implies that

$$\begin{aligned}
(\Sigma(\breve{J} \otimes \breve{J})\digamma_{r,s,m,p})(\lambda,x,\mu,y) &= ((\breve{J} \otimes \breve{J})\digamma_{r,s,m,p})(\mu,y,\lambda,x) = \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\, \overline{\digamma_{r,s,m,p}(\mu,y,\lambda,x)} \\
&= \delta_{x,\operatorname{sgn}(p)yq^m} \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\, \overline{\mu^{r+\chi(x/p)}\, \lambda^{s-\chi(y/p)}\, a_p(y,x)} \\
&= \delta_{y,\operatorname{sgn}(p)xq^{-m}} \operatorname{sgn}(p)^{\chi(p)}\, \lambda^{-s+\chi(y/p)}\, \mu^{-r-\chi(x/p)}\, a_p(x,y) = \operatorname{sgn}(p)^{\chi(p)}\, \digamma_{-s,-r,-m,p}(\lambda,x,\mu,y) \ .
\end{aligned}$$

Thus,

$$\begin{aligned}
V\Sigma(\breve{J} \otimes \breve{J})\digamma_{r,s,m,p} &= \operatorname{sgn}(p)^{\chi(p)}\, V\digamma_{-s,-r,-m,p} = \operatorname{sgn}(p)^{\chi(p)}\, \zeta^{-s} \otimes \zeta^{-r} \otimes \zeta^{-m} \otimes \delta_p \\
&= (\Sigma' \otimes \breve{J})(\zeta^r \otimes \zeta^s \otimes \zeta^m \otimes \delta_p) = (\Sigma' \otimes \breve{J})V\digamma_{r,s,m,p}
\end{aligned}$$

From this all we conclude that $V\Sigma = (\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})$. Notice that $\Sigma_{13} = \Sigma_{23}\Sigma_{12}\Sigma_{23}$. Thus

$$V_{13}(1_H \otimes V)\Sigma_{13} = V_{13}(1_H \otimes V)\Sigma_{23}\Sigma_{12}\Sigma_{23} = V_{13}(1_H \otimes [(\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})])\Sigma_{12}\Sigma_{23} \ .$$

Let $\hat{\Sigma} : L^2(\mathbb{T}^2) \otimes L^2(\mathbb{T}^2) \to L^2(\mathbb{T}^2) \otimes L^2(\mathbb{T}^2)$ denote the flip map. If we let $\Xi : L^2(\mathbb{T}^2) \otimes H \otimes H \to H \otimes L^2(\mathbb{T}^2) \otimes H$ denote the permutation map defined by $\Xi(u \otimes v \otimes w) = w \otimes u \otimes v$ for all $u \in L^2(\mathbb{T}^2)$ and $v, w \in H$, we get that

$$\begin{aligned}
V_{13}(1_H \otimes V)\Sigma_{13} &= V_{13}\, \Xi\, ([(\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})] \otimes 1_H) \\
&= (\hat{\Sigma} \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V\Sigma)\, ([(\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})] \otimes 1_H) \\
&= (\hat{\Sigma} \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes [(\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})])\, ([(\Sigma' \otimes \breve{J})V(\breve{J} \otimes \breve{J})] \otimes 1_H) \\
&= (\hat{\Sigma}(\Sigma' \otimes \Sigma') \otimes \breve{J})(1_{L^2(\mathbb{T}^2)} \otimes V)\, (V \otimes 1_H)(\breve{J} \otimes \breve{J} \otimes \breve{J}) \\
&= (\tilde{\Sigma} \otimes \breve{J})(1_{L^2(\mathbb{T}^2)} \otimes V)(V \otimes 1_H)(\breve{J} \otimes \breve{J} \otimes \breve{J}) \ .
\end{aligned}$$

☐

We end this section with the statement that the comultiplication is coassociative, the result of this paper that is the most technical to prove.

Theorem 3.11. *The* **-homomorphism* $\Delta : M_q \rightarrow M_q \otimes M_q$ *is coassociative, i.e.* $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$.

In order to enhance the readability of this paper, the proof will be given in section 5 but we stress that the results of section 4 do not play any role therein.

4. THE HAAR WEIGHT ON (M_q, Δ).

In this section we construct the left Haar weight on (M_q, Δ) and prove its left invariance through the construction of a partial isometry that afterwards turns out to be the multiplicative unitary of quantum $\widetilde{SU}(1,1)$. We also establish the unimodularity of (M_q, Δ).

Since $M_q = L^\infty(\mathbb{T}) \otimes B(L^2(I_q))$ we can consider the trace Tr on M_q given by $\text{Tr} = \text{Tr}_{L^\infty(\mathbb{T})} \otimes \text{Tr}_{B(L^2(I_q))}$, where $\text{Tr}_{L^\infty(\mathbb{T})}$ and $\text{Tr}_{B(L^2(I_q))}$ are the canonical traces on $L^\infty(\mathbb{T})$ and $B(L^2(I_q))$ which we choose to be normalized in such a way that $\text{Tr}_{L^\infty(\mathbb{T})}(1) = 1$ and $\text{Tr}_{B(L^2(I_q))}(P) = 1$ for every rank one projection P in $B(L^2(I_q))$.

Given a weight η on M_q, we use the following standard concepts from weight theory:

$$\mathcal{M}_\eta^+ = \{\, x \in M_q^+ \mid \eta(x) < \infty \,\}, \qquad \mathcal{M}_\eta = \text{linear span of } \mathcal{M}_\eta^+, \qquad \mathcal{N}_\eta = \{\, x \in M_q \mid \eta(x^*x) < \infty \,\}\ .$$

Next we introduce a GNS-construction for the trace Tr. Define $K = H \otimes L^2(I_q) = L^2(\mathbb{T}) \otimes L^2(I_q) \otimes L^2(I_q)$. If $m \in \mathbb{Z}$ and $p, t \in I_q$, we set $f_{m,p,t} = \zeta^m \otimes \delta_p \otimes \delta_t \in K$. Thus $(\, f_{m,p,t} \mid m \in \mathbb{Z}, p, t \in I_q \,)$ is an orthonormal basis for K. Now define

(1) a linear map $\Lambda_{\text{Tr}} : \mathcal{N}_{\text{Tr}} \rightarrow K$ such that $\Lambda_{\text{Tr}}(a) = \sum_{p \in I_q} (a \otimes 1_{L^2(I_q)}) f_{0,p,p}$ for $a \in \mathcal{N}_{\text{Tr}}$.

(2) a unital *-homomorphism $\pi : M_q \rightarrow B(K)$ such that $\pi(a) = a \otimes 1_{L^2(I_q)}$ for all $a \in M_q$.

Then $(K, \pi, \Lambda_{\text{Tr}})$ is a GNS-construction for Tr.

Now we are ready to define the weight that will turn out to be left- and right invariant with respect to Δ. Use the remarks before [18, Prop. 1.15] to define a linear map $\Lambda = (\Lambda_{\text{Tr}})_{\gamma^*\gamma} : D(\Lambda) \subseteq M_q \rightarrow K$.

Definition 4.1. *We define the faithful normal semi-finite weight* φ *on* M_q *as* $\varphi = \text{Tr}_{\gamma^*\gamma}$. *By definition,* (K, π, Λ) *is a GNS-construction for* φ.

See [26] for more details about this definition. This definition of φ is of course compatible with the usual construction of absolutely continuous weights (see [22]). So we already know that the modular automorphism group σ^φ of φ is such that $\sigma_s^\varphi(x) = |\gamma|^{2is}\, x\, |\gamma|^{-2is}$ for all $x \in M_q$ and $s \in \mathbb{R}$.

The modular conjugation of φ with respect to (K, π, Λ) will be denoted by J, the modular operator of φ will be denoted by ∇. Since $\varphi = \text{Tr}_{\gamma^*\gamma}$ and $\Lambda = (\Lambda_{\text{Tr}})_{\gamma^*\gamma}$, J equals the modular conjugation of Tr with respect to $(K, \pi, \Lambda_{\text{Tr}})$.

Let us recall how these modular objects are related to the modular group. So fix $a \in \mathcal{N}_\varphi$. If $t \in \mathbb{R}$, then $\sigma_t^\varphi(a) \in \mathcal{N}_\varphi$ and $\nabla^{it}\Lambda(a) = \Lambda(\sigma_t^\varphi(a))$. Provided $a \in D(\sigma_{\frac{i}{2}}^\varphi)$, the element $\sigma_{\frac{i}{2}}^\varphi(a)^* \in \mathcal{N}_\varphi$ and $J\Lambda(a) = \Lambda\big(\sigma_{\frac{i}{2}}^\varphi(a)^*\big)$. For $x \in D(\sigma_{\frac{i}{2}}^\varphi)$, we have that $ax \in \mathcal{N}_\varphi$ and $\Lambda(ax) = J\pi(\sigma_{\frac{i}{2}}^\varphi(x)^*)J\Lambda(a)$.

Let us first establish some easy formulas that make working with the weight φ pretty easy.

Lemma 4.2. *Consider* $m \in \mathbb{Z}$ *and* $p, t \in I_q$. *Then*

(1) $\Phi(m,p,t) \in \mathcal{N}_\varphi$ *and* $\Lambda(\Phi(m,p,t)) = |t|^{-1}\, f_{m,p,t}$,

(2) $\Phi(m,p,t) \in \mathcal{M}_\varphi$ *and* $\varphi(\Phi(m,p,t)) = \delta_{m,0}\, \delta_{p,t}\, t^{-2}$,

(3) $\Phi(m,p,t)$ *is analytic with respect to* σ^φ *and* $\sigma_z^\varphi(\Phi(m,p,t)) = |p^{-1}t|^{2iz}\, \Phi(m,p,t)$ *for all* $z \in \mathbb{C}$.

Proof. (1) By Eq. (2.4), $\Phi(m,p,t)^*\Phi(m,p,t) = \Phi(0,t,t) = M_{1\otimes\delta_t}$ which clearly belongs to $\mathcal{M}_{\mathrm{Tr}}^+$. Thus $\Phi(m,p,t) \in \mathcal{N}_{\mathrm{Tr}}$ and

$$\Lambda_{\mathrm{Tr}}(\Phi(m,p,t)) = \sum_{p' \in I_q} (\Phi(m,p,t) \otimes 1) f_{0,p',p'} = f_{m,p,t} \ . \tag{4.1}$$

It is clear that $\Phi(m,p,t)\,|\gamma|$ is bounded and that the extension of this element to an an element of $B(H)$ is given by $|t|^{-1}\,\Phi(m,p,t) \in \mathcal{N}_{\mathrm{Tr}}$. Therefore the remarks before [18, Prop. 1.15] imply that $\Phi(m,p,t) \in \mathcal{N}_\varphi$ and $\Lambda(\Phi(p,m,t)) = |t|^{-1}\,\Lambda_{\mathrm{Tr}}(\Phi(m,p,t)) = |t|^{-1}\,f_{m,p,t}$.

(2) By Eq. (2.4), $\Phi(m,p,t) = \Phi(0,p,p)^*\Phi(m,p,t)$, thus $\varphi(\Phi(m,p,t)) = \langle \Lambda(\Phi(m,p,t)), \Lambda(\Phi(0,p,p))\rangle$. Hence (2) follows from (1).

(3) Choose $s \in \mathbb{R}$. By definition, $\Phi(m,p,t) = \rho_{p^{-1}t}\, M_{\zeta^m \otimes \delta_t}$. Thus Lemma 2.5 implies that

$$\begin{aligned} |\gamma|^{2is}\,\Phi(m,p,t) &= M_{1\otimes|\xi|^{-2is}}\,\rho_{p^{-1}t}\,M_{\zeta^m\otimes\delta_t} = \rho_{p^{-1}t}\,M_{T_{pt^{-1}}(1\otimes|\xi|^{-2is})}\,M_{\zeta^m\otimes\delta_t} \\ &= |p^{-1}t|^{2is}\,\rho_{p^{-1}t}\,M_{1\otimes|\xi|^{-2is}}\,M_{\zeta^m\otimes\delta_t} = |p^{-1}t|^{2is}\,\rho_{p^{-1}t}\,M_{\zeta^m\otimes\delta_t}\,M_{1\otimes|\xi|^{-2is}} = |p^{-1}t|^{2is}\,\Phi(m,p,t)\,|\gamma|^{2is} \ , \end{aligned}$$

from which it follows that $\sigma_s^\varphi(\Phi(m,p,t)) = |p^{-1}t|^{2is}\,\Phi(m,p,t)$ and the claim follows. ☐

These results imply easily the next lemma which is needed to extend results (involving Λ) that have been proven for the elements $\Phi(m,p,t)$ to the whole of $\mathcal{N}_\varphi$.

Corollary 4.3. *If $a \in \mathcal{N}_\varphi$, there exists a bounded net $(a_i)_{i\in I}$ in $\langle\, \Phi(m,p,t) \mid m \in \mathbb{Z}, p,t \in I_q \,\rangle$ such that $(a_i)_{i\in I}$ converges strongly* to a and $(\Lambda(a_i))_{i\in I}$ converges to $\Lambda(a)$.*

Proof. Define $C = \langle\, \Phi(m,p,t) \mid m \in \mathbb{Z}, p,t \in I_q \,\rangle$. Set $B = \{\, b \in M_q \mid \|b\| \leq \|a\| \,\}$ and equip B with the strong* topology. In $B \times K$, we consider the set G that is the closure of $\{\, (b, \Lambda(b)) \mid b \in B \cap C \,\}$.

Let $F(I_q)$ denote the set of all finite subsets of I_q and turn $F(I_q)$ into a directed set by inclusion. For $L \in F(I_q)$, we define the projection $P_L = \sum_{v \in L} \Phi(0,v,v)$. Thus, $(P_L)_{L \in F(I_q)}$ converges strongly to 1.

Fix $L \in F(I_q)$ for the moment. By Lemma 4.2(1) we see that $P_L \in \mathcal{N}_\varphi$. Since $\Lambda(bP_L) = \pi(b)\,\Lambda(P_L)$ for all $b \in M_q$, Kaplansky's density Theorem (see the proof of [7, Thm. 5.3.5]), applied to the *-algebra C, implies that $(aP_L, \Lambda(aP_L)) \in G$.

It is clear from Lemma 4.2(3) that $P_L \in D(\sigma^\varphi_{\frac{i}{2}})$ and $\sigma^\varphi_{\frac{i}{2}}(P_L) = P_L$. Thus, $\Lambda(aP_L) = J\pi(P_L)J\Lambda(a)$. It follows that $\big(\,(aP_L, \Lambda(aP_L))\,\big)_{L \in F(I_q)} \to (a, \Lambda(a))$, implying that $(a, \Lambda(a)) \in G$ and the lemma follows. ☐

Corollary 4.4. *Consider $m \in \mathbb{Z}$ and $p,t \in I_q$. Then*

1) $J f_{m,p,t} = f_{-m,t,p}$,
2) $f_{m,p,t} \in D(\nabla)$ *and* $\nabla f_{m,p,t} = |p^{-1}t|^2\, f_{m,p,t}$.

Proof. Recall that J is also the modular conjugation of the trace Tr. So Eq. (4.1) in the previous proof implies that $J f_{m,p,t} = J\Lambda_{\mathrm{Tr}}(\Phi(m,p,t)) = \Lambda_{\mathrm{Tr}}(\Phi(m,p,t)^*) = \Lambda_{\mathrm{Tr}}(\Phi(-m,t,p)) = f_{-m,t,p}$.

Let $s \in \mathbb{R}$. By Lemma 4.2, $\Phi(m,p,t) \in \mathcal{N}_\varphi$ and $\sigma_s^\varphi(\Phi(m,p,t)) = |p^{-1}t|^{2is}\,\Phi(m,p,t)$. So we get that $\nabla^{is}\Lambda(\Phi(m,p,t)) = \Lambda\big(\sigma_s^\varphi(\Phi(m,p,t))\big) = |p^{-1}t|^{2is}\,\Lambda(\Phi(m,p,t))$. Thus, since $f_{m,p,t} = |t|\,\Lambda(\Phi(m,p,t))$, $\nabla^{is} f_{m,p,t} = |p^{-1}t|^{2is}\, f_{m,p,t}$ and also the second claim follows. ☐

Now we know enough about the weight φ to proceed to the proof of the left invariance of φ with respect to Δ. For this purpose we introduce a partial isometry that will later turn out to be the multiplicative unitary of (M_q, Δ). The formula defining W^* in the next proposition is obtained by formally calculating $W^*\big(\Lambda(\Phi(m_1,p_1,t_1)) \otimes \Lambda(\Phi(m_2,p_2,t_2))\big) = (\Lambda \otimes \Lambda)(\Delta(\Phi(m_2,p_2,t_2))(\Phi(m_1,p_1,t_1) \otimes 1))$, but as this stage we do not know whether φ is left invariant so we cannot use this formula, as is done in the general theory, to define W^*.

Proposition 4.5. *There exists a unique surjective partial isometry W on $K \otimes K$ such that*

$$\begin{aligned} W^*(f_{m_1,p_1,t_1} \otimes f_{m_2,p_2,t_2}) = \sum_{\substack{y,\, z \,\in\, I_q \\ \mathrm{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q}} & |t_2/y|\, a_{t_2}(p_1,y)\, a_{p_2}(z, \mathrm{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \\ & \times \ f_{m_1+m_2-\chi(p_1p_2/t_2z),z,t_1} \otimes f_{\chi(p_1p_2/t_2z),\mathrm{sgn}(p_2t_2)(yz/p_1)q^{m_2},y} \end{aligned}$$

for all $m_1, m_2 \in \mathbb{Z}$ and $p_1, p_2, t_1, t_2 \in I_q$.

Proof. The proof of this proposition is based on the properties of the functions a_p, in particular on Propositions 3.2, 3.3 and 3.5. We now give details.
Fix $m_1, m_2 \in \mathbb{Z}$ and $p_1, p_2, t_1, t_2 \in I_q$. If $y, z, y', z' \in I_q$ such that $\operatorname{sgn}(p_2 t_2)(yz/p_1)q^{m_2}$, $\operatorname{sgn}(p_2 t_2)(y'z'/p_1)q^{m_2} \in I_q$ and $(y,z) \neq (y',z')$, then it is clear that the element $f_{m_1+m_2-\chi(p_1p_2/t_2z),z,t_1} \otimes f_{\chi(p_1p_2/t_2z),\operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2},y}$ is orthogonal to the element $f_{m_1+m_2-\chi(p_1p_2/t_2z'),z',t_1} \otimes f_{\chi(p_1p_2/t_2z'),\operatorname{sgn}(p_2t_2)(y'z'/p_1)q^{m_2},y'}$. By Propositions 3.2, 3.3 and 3.5 we get moreover

$$\begin{aligned}
&\sum_{\substack{y,\, z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q}} |t_2/y|^2\, |a_{t_2}(p_1,y)|^2\, |a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2})|^2 \\
&\quad = \sum_{y \in I_q} |t_2/y|^2\, |a_{t_2}(p_1,y)|^2 \sum_{\substack{z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q}} |a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2})|^2 \\
&\quad \leq \sum_{y \in I_q} |t_2/y|^2\, |a_{t_2}(p_1,y)|^2 = \sum_{y \in I_q} |a_y(p_2,t_1)|^2 = 1 \ . \qquad (4.2)
\end{aligned}$$

From this, it follows that we can define the element $v(m_1,p_1,t_1;m_2,p_2,t_2) \in K \otimes K$ as

$$\sum_{\substack{y,\, z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q}} |t_2/y|\, a_{t_2}(p_1,y)\, a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \\
\times\ f_{m_1+m_2-\chi(p_1p_2/t_2z),z,t_1} \otimes f_{\chi(p_1p_2/t_2z),\operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2},y} \ .$$

Note that $v(m_1,p_1,t_1;m_2,p_2,t_2)$ is just the right hand side of the formula we want to define W^* by. Now choose also $m_1', m_2' \in \mathbb{Z}$ and $p_1', p_2', t_1', t_2' \in I_q$. Notice that inequality (4.2) together with the Cauchy-Schwarz equality implies that

$$\sum_{\substack{y,\, z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q \\ \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'} \,\in\, I_q}} |\,(t_2/y)\,(t_2'/y)\, a_{t_2}(p_1,y)\, a_{t_2'}(p_1',y)\ \ a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \\
a_{p_2'}(z, \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'})\,|$$

is finite, which allows us to compute the next sum in any order we want:

$$\begin{aligned}
&\langle v(m_1,p_1,t_1;m_2,p_2,t_2), v(m_1',p_1',t_1';m_2',p_2',t_2') \rangle \\
&= \sum_{\substack{y,\, z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q \\ \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'} \,\in\, I_q}} |t_2/y|\, |t_2'/y|\, a_{t_2}(p_1,y)\, a_{t_2'}(p_1',y)\, a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \\
&\qquad \times\ a_{p_2'}(z, \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'})\ \langle f_{m_1+m_2-\chi(p_1p_2/t_2z),z,t_1}, f_{m_1'+m_2'-\chi(p_1'p_2'/t_2'z),z,t_1'} \rangle \\
&\qquad \times\ \langle f_{\chi(p_1p_2/t_2z),\operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2},y}, f_{\chi(p_1'p_2'/t_2'z),\operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'},y} \rangle \\
&= \sum_{\substack{y,\, z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q \\ \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'} \,\in\, I_q}} |t_2/y|\, |t_2'/y|\, a_{t_2}(p_1,y)\, a_{t_2'}(p_1',y)\, a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \\
&\qquad \times\ a_{p_2'}(z, \operatorname{sgn}(p_2't_2')(yz/p_1')q^{m_2'})\ \delta_{m_1+m_2,m_1'+m_2'}\, \delta_{t_1,t_1'}\, \delta_{|p_1p_2/t_2|,|p_1'p_2'/t_2'|} \\
&\qquad \times\ \delta_{\operatorname{sgn}(p_2t_2)q^{m_2}/p_1,\operatorname{sgn}(p_2't_2')q^{m_2'}/p_1'} \\
&= \delta_{m_1+m_2,m_1'+m_2'}\, \delta_{t_1,t_1'}\, \delta_{p_1p_2/t_2,p_1'p_2'/t_2'}\, \delta_{\operatorname{sgn}(p_2t_2)q^{m_2}/p_1,\operatorname{sgn}(p_2't_2')q^{m_2'}/p_1'} \sum_{y \in I_q} |t_2/y|\, |t_2'/y|\, a_{t_2}(p_1,y) \\
&\quad a_{t_2'}(p_1',y) \sum_{\substack{z \,\in\, I_q \\ \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2} \,\in\, I_q}} a_{p_2}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2})\, a_{p_2'}(z, \operatorname{sgn}(p_2t_2)(yz/p_1)q^{m_2}) \ .
\end{aligned}$$

Thus, using the orthogonality relation in Proposition 3.2 together with Proposition 3.5, we get that

$$\begin{aligned}
&\langle v(m_1,p_1,t_1;m_2,p_2,t_2), v(m_1',p_1',t_1';m_2',p_2',t_2')\rangle = \delta_{m_1+m_2,m_1'+m_2'}\,\delta_{t_1,t_1'}\,\delta_{p_1p_2/t_2,p_1'p_2'/t_2'} \\
&\qquad \delta_{\operatorname{sgn}(p_2t_2)q^{m_2}/p_1,\operatorname{sgn}(p_2't_2')q^{m_2'}/p_1'} \sum_{y\in I_q} |t_2/y|\,|t_2'/y|\,a_{t_2}(p_1,y)\,a_{t_2'}(p_1',y)\,\delta_{p_2,p_2'}\,\delta_{\operatorname{sgn}(y),\operatorname{sgn}(p_1t_2)} \\
&= \delta_{m_1+m_2,m_1'+m_2'}\,\delta_{t_1,t_1'}\,\delta_{p_1/t_2,p_1'/t_2'}\,\delta_{\operatorname{sgn}(t_2)q^{m_2}/p_1,\operatorname{sgn}(t_2')q^{m_2'}/p_1'}\,\delta_{p_2,p_2'} \\
&\quad \sum_{y\in I_q} (-1)^{\chi(yt_2)}\operatorname{sgn}(p_1)^{\chi(p_1)}\,(-1)^{\chi(yt_2')}\operatorname{sgn}(p_1')^{\chi(p_1')}\,a_y(p_1,t_2)\,a_y(p_1',t_2') \\
&= \delta_{m_1+m_2,m_1'+m_2'}\,\delta_{t_1,t_1'}\,\delta_{p_1/t_2,p_1'/t_2'}\,\delta_{\operatorname{sgn}(t_2)q^{m_2}/p_1,\operatorname{sgn}(t_2')q^{m_2'}/p_1'}\,\delta_{p_2,p_2'} \\
&\quad (-1)^{\chi(t_2t_2')}\operatorname{sgn}(p_1)^{\chi(p_1)}\operatorname{sgn}(p_1')^{\chi(p_1')}\sum_{y\in I_q} a_y(p_1,t_2)\,a_y(p_1',t_2')\ .
\end{aligned}$$

Because of the factor $\delta_{p_1/t_2,p_1'/t_2'}$ in the above expression, Proposition 3.3 implies that

$$\begin{aligned}
&\langle v(m_1,p_1,t_1;m_2,p_2,t_2), v(m_1',p_1',t_1';m_2',p_2',t_2')\rangle = \delta_{m_1+m_2,m_1'+m_2'}\,\delta_{t_1,t_1'} \\
&\quad \delta_{p_1/t_2,p_1'/t_2'}\,\delta_{\operatorname{sgn}(t_2)q^{m_2}/p_1,\operatorname{sgn}(t_2')q^{m_2'}/p_1'}\,\delta_{p_2,p_2'}\,(-1)^{\chi(t_2t_2')}\operatorname{sgn}(p_1)^{\chi(p_1)}\operatorname{sgn}(p_1')^{\chi(p_1')}\,\delta_{p_1,p_1'}\,\delta_{t_2,t_2'} \\
&= \delta_{m_1,m_1'}\,\delta_{m_2,m_2'}\,\delta_{p_1,p_1'}\,\delta_{p_2,p_2'}\,\delta_{t_1,t_1'}\,\delta_{t_2,t_2'} = \langle f_{m_1,p_1,t_1}\otimes f_{m_2,p_2,t_2}, f_{m_1',p_1',t_1'}\otimes f_{m_2',p_2',t_2'}\rangle\ .
\end{aligned}$$

Now the lemma follows easily. □

The next proposition deals with the essential step towards the left invariance of the weight φ and the realization that W is the multiplicative unitary associated to quantum $\widetilde{SU}(1,1)$.

Proposition 4.6. *Consider $\omega \in B(K)_*$ and $a \in \mathcal{N}_\varphi$. Then $(\omega\pi\otimes\iota)\Delta(a) \in \mathcal{N}_\varphi$ and $\Lambda((\omega\pi\otimes\iota)\Delta(a)) = (\omega\otimes\iota)(W^*)\,\Lambda(a)$.*

Proof. Choose $m, m_1, m_2 \in \mathbb{Z}$, $p, t, p_1, t_1, p_2, t_2 \in I_q$. Take also $v \in I_q$ and let us calculate the element $(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\Phi(m_1,p_1,t_1)\otimes\Phi(0,v,v))$. Choose $r, s \in \mathbb{Z}$ and $x, y \in I_q$. Then Proposition 3.7, Definition 3.6 and Eqs. (2.3), (2.4) tell us that

$$\begin{aligned}
&(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))\,(\zeta^r\otimes\delta_x\otimes\zeta^s\otimes\delta_y) \\
&\quad = \sum_{z\in I_q} a_z(x,y)\,(\Phi(-m_2,t_2,p_2)\otimes 1)V^*(1\otimes\Phi(m,p,t))V\,\digamma_{r-\chi(y/z),s+\chi(x/z),\chi(y/x),z} \\
&\quad = \sum_{z\in I_q} a_z(x,y)\,(\Phi(-m_2,t_2,p_2)\otimes 1)V^*(1\otimes\Phi(m,p,t))(\zeta^{r-\chi(y/z)}\otimes\zeta^{s+\chi(x/z)}\otimes\zeta^{\chi(y/x)}\otimes\delta_z) \\
&\quad = a_t(x,y)\,(\Phi(-m_2,t_2,p_2)\otimes 1)V^*(\zeta^{r-\chi(y/t)}\otimes\zeta^{s+\chi(x/t)}\otimes\zeta^{\chi(y/x)+m}\otimes\delta_p) \\
&\quad = a_t(x,y)\,(\Phi(-m_2,t_2,p_2)\otimes 1)\,\digamma_{r-\chi(y/t),s+\chi(x/t),\chi(y/x)+m,p} \\
&\quad = \sum_{\substack{z\,\in\, I_q \\ \operatorname{sgn}(p)|y/x|q^m z\,\in\, I_q}} a_t(x,y)\,a_p(z,\operatorname{sgn}(p)|y/x|q^m z)\,(\Phi(-m_2,t_2,p_2)\otimes 1) \\
&\qquad\qquad \zeta^{r-\chi(y/t)+\chi(q^m zy/px)}\otimes\delta_z\otimes\zeta^{s+\chi(x/t)-\chi(z/p)}\otimes\delta_{\operatorname{sgn}(p)|y/x|q^m z} \\
&\quad = \sum_{\substack{z\,\in\, I_q \\ \operatorname{sgn}(pt)(y/x)q^m z\,\in\, I_q}} a_t(x,y)\,a_p(z,\operatorname{sgn}(pt)(y/x)q^m z)\,(\Phi(-m_2,t_2,p_2)\otimes 1) \\
&\qquad\qquad \zeta^{r+m-\chi(px/zt)}\otimes\delta_z\otimes\zeta^{s+\chi(px/zt)}\otimes\delta_{\operatorname{sgn}(pt)(y/x)q^m z}
\end{aligned}$$

by Eq. (2.3). So we see that $(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))\,(\zeta^r\otimes\delta_x\otimes\zeta^s\otimes\delta_y) = 0$ if $\operatorname{sgn}(pt)(y/x)q^m p_2 \notin I_q$. If, on the other hand, $\operatorname{sgn}(pt)(y/x)q^m p_2 \in I_q$, then

$$\begin{aligned}
&(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))\,(\zeta^r\otimes\delta_x\otimes\zeta^s\otimes\delta_y) \\
&\quad = a_t(x,y)\,a_p(p_2,\operatorname{sgn}(pt)(y/x)q^m p_2)\,(\zeta^{r+m-m_2-\chi(px/p_2t)}\otimes\delta_{t_2}\otimes\zeta^{s+\chi(px/p_2t)}\otimes\delta_{\operatorname{sgn}(pt)(y/x)q^m p_2})\ .
\end{aligned}$$

Now we have for $r,s\in\mathbb{Z}$ and $x,y\in I_q$ that

$$\begin{aligned}&(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\Phi(m_1,p_1,t_1)\otimes\Phi(0,v,v))\,(\zeta^r\otimes\delta_x\otimes\zeta^s\otimes\delta_y)\\&\quad=\delta_{t_1,x}\,\delta_{v,y}\,(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\zeta^{m_1+r}\otimes\delta_{p_1}\otimes\zeta^s\otimes\delta_v)\ .\end{aligned}$$

So we see that $(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\Phi(m_1,p_1,t_1)\otimes\Phi(0,v,v))=0$ if $\operatorname{sgn}(pt)(v/p_1)q^m p_2\notin I_q$. If, on the other hand, $\operatorname{sgn}(pt)(v/p_1)q^m p_2\in I_q$, then

$$\begin{aligned}&(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\Phi(m_1,p_1,t_1)\otimes\Phi(0,v,v))\,(\zeta^r\otimes\delta_x\otimes\zeta^s\otimes\delta_y)\\&\quad=\delta_{t_1,x}\,\delta_{v,y}\,a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2)\\&\qquad\zeta^{r+m_1+m-m_2-\chi(pp_1/p_2t)}\otimes\delta_{t_2}\otimes\zeta^{s+\chi(pp_1/p_2t)}\otimes\delta_{\operatorname{sgn}(pt)(v/p_1)q^m p_2}\ .\end{aligned}$$

and thus

$$\begin{aligned}(\Phi(m_2,p_2,t_2)^*\otimes 1)\Delta(\Phi(m,p,t))(\Phi(m_1,p_1,t_1)\otimes\Phi(0,v,v))&=a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2)\\\times\,\Phi(m_1+m-m_2-\chi(pp_1/p_2t),t_2,t_1)&\otimes\Phi(\chi(pp_1/p_2t),\operatorname{sgn}(pt)(v/p_1)q^m p_2,v)\ .\end{aligned}$$

Applying $\varphi\otimes\iota$ to this equation and using Lemma 4.2, we see that

$$\begin{aligned}&[\,(\omega_{\Lambda(\Phi(m_1,p_1,t_1)),\Lambda(\Phi(m_2,p_2,t_2))}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,]\,\Phi(0,v,v)=\delta_{m_1+m-m_2,\chi(pp_1/p_2t)}\,\delta_{t_1,t_2}\\&\qquad\times(t_1)^{-2}\,a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2)\,\Phi(m+m_1-m_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2,v)\ .\end{aligned}$$

Or, again by Lemma 4.2,

$$\begin{aligned}&(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,\Phi(0,v,v)=\delta_{m_1+m-m_2,\chi(pp_1/p_2t)}\,\delta_{t_1,t_2}\\&\qquad\times|t_2/t_1|\,a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2)\,\Phi(m+m_1-m_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2,v)\ ,\end{aligned}$$

so we conclude that if $\operatorname{sgn}(pt)(v/p_1)q^m p_2\in I_q$, the element $(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))$ $\Phi(0,v,v)$ belongs to $\mathcal{N}_\varphi$ and

$$\begin{aligned}&\Lambda\big(\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,\Phi(0,v,v)\,\big)=\delta_{m_1+m-m_2,\chi(pp_1/p_2t)}\,\delta_{t_1,t_2}\\&\qquad\times|v|^{-1}\,a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2)f_{m+m_1-m_2,\operatorname{sgn}(pt)(v/p_1)q^m p_2,v}\ .\end{aligned}\tag{4.3}$$

Also recall that $(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,\Phi(0,v,v)=0$ if $\operatorname{sgn}(pt)(v/p_1)q^m p_2\notin I_q$.

Notice that Corollary 4.4 implies that $J\pi(\Phi(0,v,v))Jf_{n,x,y}=J\pi(\Phi(0,v,v))f_{-n,y,x}=\delta_{v,y}\,Jf_{-n,y,x}=\delta_{v,y}f_{n,x,y}$. So we get by Proposition 4.5 that

$$\begin{aligned}&J\pi(\Phi(0,v,v))J\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\otimes\iota)(W^*)\,f_{m,p,t}=\\&\quad=\delta_{t_1,t_2}\,\delta_{m_1+m-\chi(p_1p/tp_2),m_2}\sum_{\substack{y\,\in\,I_q\\ \operatorname{sgn}(pt)(yp_2/p_1)q^m\,\in\,I_q}}|t/y|\,a_t(p_1,y)\,a_p(p_2,\operatorname{sgn}(pt)(yp_2/p_1)q^m)\\&\qquad\qquad\times\ J\pi(\Phi(0,v,v))J\,f_{\chi(p_1p/tp_2),\operatorname{sgn}(pt)(yp_2/p_1)q^m,y}\\&\quad=\delta_{t_1,t_2}\,\delta_{m_1+m-\chi(p_1p/tp_2),m_2}\sum_{\substack{y\,\in\,I_q\\ \operatorname{sgn}(pt)(yp_2/p_1)q^m\,\in\,I_q}}|t/y|\,a_t(p_1,y)\,a_p(p_2,\operatorname{sgn}(pt)(yp_2/p_1)q^m)\\&\qquad\qquad\times\ \delta_{v,y}\,f_{m+m_1-m_2,\operatorname{sgn}(pt)(yp_2/p_1)q^m,y}\ .\end{aligned}$$

So we see that $J\pi(\Phi(0,v,v))J\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\otimes\iota)(W^*)\,f_{m,p,t}=0$ if $\operatorname{sgn}(pt)(vp_2/p_1)q^m\notin I_q$. If, on the other hand, $\operatorname{sgn}(pt)(vp_2/p_1)q^m\in I_q$,

$$\begin{aligned}&J\pi(\Phi(0,v,v))J\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\otimes\iota)(W^*)\,f_{m,p,t}\\&\quad=\delta_{t_1,t_2}\,\delta_{m_1+m-\chi(p_1p/tp_2),m_2}\,|t/v|\,a_t(p_1,v)\,a_p(p_2,\operatorname{sgn}(pt)(vp_2/p_1)q^m)\,f_{m+m_1-m_2,\operatorname{sgn}(pt)(vp_2/p_1)q^m,v}\ .\end{aligned}$$

From Eq. (4.3) and Lemma 4.2, we conclude that

$$\begin{aligned}&\Lambda\big(\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,\Phi(0,v,v)\,\big)\\&\qquad=J\pi(\Phi(0,v,v))J\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\otimes\iota)(W^*)\,\Lambda(\Phi(m,p,t))\ .\end{aligned}$$

We use the net of projections $(P_L)_{L\in F(I_q)}$ introduced in the proof of Corollary 4.3. If $L\in F(I_q)$, we know by the previous equation that $(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,P_L\in\mathcal{N}_\varphi$ and

$$\Lambda\big(\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi\otimes\iota)(\Delta(\Phi(m,p,t)))\,P_L\,\big)=J\pi(P_L)J\,(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\otimes\iota)(W^*)\,\Lambda(\Phi(m,p,t))\ .$$

Therefore the σ-strong* closedness of Λ implies that $(\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi \otimes \iota)(\Delta(\Phi(m,p,t))) \in \mathcal{N}_\varphi$ and

$$\Lambda\big((\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi \otimes \iota)(\Delta(\Phi(m,p,t)))\big) = (\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}} \otimes \iota)(W^*)\,\Lambda(\Phi(m,p,t))\ .$$

Since the linear span of such linear functionals $\omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}}\pi$ is norm dense in $(M_q)_*$, the proposition now follows easily from Corollary 4.3 and the σ-strong* closedness Λ. ☐

Now, the proof of the left invariance of φ can be dealt with as in the proof of [18, Prop. 8.15].

Proposition 4.7. *The weight φ is left invariant with respect to Δ.*

Proof. Take $a \in \mathcal{N}_\varphi$ and $\omega \in (M_q)^+_*$. Since $\pi(M_q)$ is in standard form with respect to K, there exists a vector $v \in K$ such that $\omega = \omega_{v,v}\pi$. Take also an orthonormal basis $(e_i)_{i\in I}$ for K. It follows from the proof of [18, Lem. A.5] that $\big(\sum_{i\in L}(\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))^*(\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))\big)_{L\in F(I)}$ is an increasing net in M_q^+ that converges strongly to $(\omega\otimes\iota)\Delta(a^*a)$ (as before, $F(I)$ denotes the set of finite subsets of I, directed by inclusion). Therefore the σ-weak lower semi-continuity of φ implies that

$$\varphi\big((\omega\otimes\iota)\Delta(a^*a)\big) = \sum_{i\in I}\varphi\big((\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))^*(\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))\big)\ .$$

By the previous proposition, this implies that

$$\begin{aligned}\varphi\big((\omega\otimes\iota)\Delta(a^*a)\big) &= \sum_{i\in I}\langle\Lambda\big((\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))\big),\Lambda\big((\omega_{v,e_i}\pi\otimes\iota)(\Delta(a))\big)\rangle\\ &= \sum_{i\in I}\langle(\omega_{v,e_i}\otimes\iota)(W^*)\Lambda(a),(\omega_{v,e_i}\otimes\iota)(W^*)\Lambda(a)\rangle = \sum_{i\in I}\langle(\omega_{v,e_i}\otimes\iota)(W^*)^*(\omega_{v,e_i}\otimes\iota)(W^*)\Lambda(a),\Lambda(a)\rangle\ .\end{aligned}$$

So if we use first [18, Lem. A.5] and afterwards the fact that W^* is an isometry (Proposition 4.5), we see that

$$\varphi\big((\omega\otimes\iota)\Delta(a^*a)\big) = \langle(\omega_{v,v}\otimes\iota)(WW^*)\Lambda(a),\Lambda(a)\rangle = \omega_{v,v}(1)\,\langle\Lambda(a),\Lambda(a)\rangle = \omega(1)\,\varphi(a^*a)$$

and we have proven the proposition. ☐

We have introduced the anti-unitary operator $\breve{J}$ on H before Proposition 3.10. Thus, $\breve{J}(\zeta^n\otimes\delta_x) = \mathrm{sgn}(x)^{\chi(x)}\,\zeta^{-n}\otimes\delta_x$ for all $n\in\mathbb{Z}$ and $x\in I_q$. This implies easily that $\breve{J}\Phi(m,p,t)\breve{J} = \mathrm{sgn}(p)^{\chi(p)}\,\mathrm{sgn}(t)^{\chi(t)}\,\Phi(-m,p,t)$ for $m\in\mathbb{Z}$ and $p,t\in I_q$. It shows first of all $\breve{J}M_q\breve{J} = M_q$. This allows us to define an anti-*-isomorphism $\breve{R}$ on M_q such that $\breve{R}(a) = \breve{J}a^*\breve{J}$ for all $a\in M_q$. Proposition 3.10 guarantees that $\chi(\breve{R}\otimes\breve{R})\Delta = \Delta\breve{R}$, where χ denotes the flip map $\chi : M_q\otimes M_q \to M_q\otimes M_q$. Later on, $\breve{R}$ turns out to be intimately connected with the unitary antipode.

Proposition 4.8. *We have that $\varphi = \varphi\breve{R}$. Thus φ is also right invariant.*

Proof. If $m\in\mathbb{Z}$ and $p,t\in I_q$, then $\breve{R}(\Phi(m,p,t))^* = \breve{J}\Phi(m,p,t)\breve{J} = \mathrm{sgn}(p)^{\chi(p)}\,\mathrm{sgn}(t)^{\chi(t)}\,\Phi(-m,p,t)$. Hence, Lemma 4.2 implies that $\breve{R}(\Phi(m,p,t))^*\in\mathcal{N}_\varphi$. This same lemma implies moreover for $m,m'\in\mathbb{Z}$ and $p,t,p',t'\in I_q$:

$$\begin{aligned}&\langle\Lambda(\breve{R}(\Phi(m,p,t))^*),\Lambda(\breve{R}(\Phi(m',p',t'))^*)\rangle\\ &\quad= \mathrm{sgn}(p)^{\chi(p)}\,\mathrm{sgn}(t)^{\chi(t)}\,\mathrm{sgn}(t')^{\chi(t')}\,\mathrm{sgn}(p')^{\chi(p')}\,\langle\Lambda(\Phi(-m,p,t)),\Lambda(\Phi(-m',p',t'))\rangle\\ &\quad= \mathrm{sgn}(p)^{\chi(p)}\,\mathrm{sgn}(t)^{\chi(t)}\,\mathrm{sgn}(t')^{\chi(t')}\,\mathrm{sgn}(p')^{\chi(p')}\,|tt'|^{-1}\delta_{m,m'}\,\delta_{p,p'}\,\delta_{t,t'}\\ &\quad= |tt'|^{-1}\,\delta_{m,m'}\,\delta_{p,p'}\,\delta_{t,t'} = \langle\Lambda(\Phi(m',p',t')),\Lambda(\Phi(m,p,t))\rangle\ .\end{aligned}$$

If $a\in\langle\,\Phi(m,p,t)\mid m\in\mathbb{Z},p,t\in I_q\,\rangle$, the above result implies that $\breve{R}(a)^*\in\mathcal{N}_\varphi$ and $\|\Lambda(\breve{R}(a)^*)\| = \|\Lambda(a)\|$. By Corollary 4.3 and the σ-strong* closedness of Λ this implies that for all $a\in\mathcal{N}_\varphi$, the element $\breve{R}(a)^*\in\mathcal{N}_\varphi$ and $\|\Lambda(\breve{R}(a)^*)\| = \|\Lambda(a)\|$, thus $\varphi(\breve{R}(a^*a)) = \varphi(\breve{R}(a)\breve{R}(a)^*) = \|\Lambda(\breve{R}(a)^*)\| = \|\Lambda(a)\| = \varphi(a^*a)$. Since $\breve{R}^2 = \iota$, it follows that $\varphi = \varphi\breve{R}$. Because $\chi(\breve{R}\otimes\breve{R})\Delta = \Delta\breve{R}$ and φ is left invariant, we get that φ is right invariant. ☐

Combined with Theorem 3.11, we have proven the main result of this paper (see Definition 1 of the Introduction for used terminology):

Theorem 4.9. *The pair (M_q, Δ) is a von Neumann algebraic quantum group which we denote by $\widetilde{SU}_q(1,1)$.*

By Proposition 4.6 and the remark after [19, Thm.1.2], we also conclude that

Proposition 4.10. *The element W is the multiplicative unitary of (M_q, Δ) with respect to the GNS-construction (K, π, Λ).*

The unitarity can also be proved directly from Propositions 4.5, 3.2, 3.3 and 3.5. We denote the antipode, unitary antipode and scaling group of (M_q, Δ) by S, R and τ respectively. All these objects are defined after the proof of [19, Prop. 1.4]. The modular element and scaling constant of a quantum group are introduced at the same place. Because (M_q, Δ) is unimodular, that is, possesses a weight that is left and right invariant, we get the following result.

Proposition 4.11. *We have that $\varphi R = \varphi$. Furthermore, the modular element of (M_q, Δ) is the identity operator and the scaling constant of (M_q, Δ) equals 1.*

Proof. We denote the modular element and scaling constant of (M_q, Δ) by δ and ν respectively, see the discussion after the proof of [19, Prop. 1.4]. We know that $\sigma_s^{\varphi}(\delta) = \nu^s\, \delta$ for all $s \in \mathbb{R}$ (*) and $\varphi R = \varphi_\delta$. Because φ is right invariant, the uniqueness of right Haar weights implies the existence of $\lambda > 0$ such that $\varphi R = \lambda\, \varphi$. Thus, the uniqueness of the Radon-Nykodim derivative guarantees that $\delta = \lambda 1$. But, since $\Delta(\delta) = \delta \otimes \delta$, it follows that $\delta = 1$. Therefore, (*) implies that $\nu = 1$. $\square$

In the next part of this section we calculate explicit expressions for S, R and τ acting on elements $\Phi(m,p,t)$. For this purpose we calculate some explicit slices $(\iota \otimes \omega)(W^*)$, where $\omega \in B(K)_*$.

Consider $m, m_1, m_2 \in \mathbb{Z}$, $p, p_1, p_2, t, t_1, t_2 \in I_q$. It follows easily from Proposition 4.5 that

$$\begin{aligned}(\iota \otimes \omega_{f_{m_1,p_1,t_1}, f_{m_2,p_2,t_2}})(W^*)\, f_{m,p,t} \\ = |t_1/t_2|\, a_{t_1}(p,t_2)\, a_{p_1}(\operatorname{sgn}(p_1 t_1)(p p_2/t_2) q^{-m_1}, p_2)\, f_{m_1-m_2+m, \operatorname{sgn}(p_1 t_1)(p p_2/t_2) q^{-m_1}, t}\end{aligned} \tag{4.4}$$

if $\operatorname{sgn}(p_1 t_1)(p p_2/t_2) q^{-m_1} \in I_q$ and $\chi(p_1 t_2/p_2 t_1) = m_2 - m_1$. If, on the other hand, $\operatorname{sgn}(p_1 t_1)(p p_2/t_2) q^{-m_1} \notin I_q$ or $\chi(p_1 t_2/p_2 t_1) \neq m_2 - m_1$, then $(\iota \otimes \omega_{f_{m_1,p_1,t_1}, f_{m_2,p_2,t_2}})(W^*)\, f_{m,p,t} = 0$.

Proposition 4.12. *Consider $m_1, m_2, m_3, m_4 \in \mathbb{Z}$ and $p, t \in I_q$. Then we define $\omega \in B(K)_*$ as the absolutely convergent sum*

$$\begin{aligned}\omega = \sum_{\substack{x,\, y \,\in\, I_q \\ \operatorname{sgn}(p) q^{m_3} x \,\in\, I_q \\ \operatorname{sgn}(t) q^{m_4} y \,\in\, I_q}} & \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\, (-1)^{\chi(xy)}\, |xy| a_p(\operatorname{sgn}(p) q^{m_3} x, x)\, a_t(\operatorname{sgn}(t) q^{m_4} y, y) \\ & \times\ \omega_{f_{m_1+\chi(x/y), \operatorname{sgn}(p) q^{m_3} x, \operatorname{sgn}(t) q^{m_4} y}, f_{m_2+\chi(x/y), x, y}}\ .\end{aligned} \tag{4.5}$$

Then $(\iota \otimes \omega)(W^*) = t^2\, q^{-m_1-m_3}\, (-1)^{m_2}\, \delta_{\chi(t/p), m_1}\, \delta_{m_3-m_4, m_2-m_1}\, \pi(\Phi(m_1 - m_2, p, t))$.

Proof. Let $s \in I_q$ and $m \in \mathbb{Z}$. By Definition 3.1, we have for $z \in I_q$ satisfying $\operatorname{sgn}(s)\, q^m z \in I_q$,

$$\begin{aligned}& |a_s(\operatorname{sgn}(s) q^m z, z)| \\ & \quad \leq c_q\, \nu(s q^{-m}) \sqrt{\frac{(-\kappa(s); q^2)_\infty}{(q^2; q^2)_\infty}}\, |z|\, \sqrt{(-\kappa(z); q^2)_\infty}\, |\Psi(-q^2/\kappa(s); (q^{2(1+m)}/s^2)\kappa(z); \operatorname{sgn}(s) q^{2(m+1)})|\ .\end{aligned}$$

which by Lemma 6.8 implies that $\sum_{z \in I_q, \operatorname{sgn}(s) q^m z \in I_q} |a_s(\operatorname{sgn}(s) q^m z, z)| < \infty$. It follows that the sum in the statement of this proposition is absolutely convergent. So, as a norm convergent limit of finite sums of vector functionals, we obtain $\omega \in B(K)_*$. Take $n \in \mathbb{Z}$ and $c, d \in I_q$. Now

$$\begin{aligned}(\iota \otimes \omega)(W^*)\, f_{n,c,d} = \sum_{\substack{x,\, y \,\in\, I_q \\ \operatorname{sgn}(p) q^{m_3} x \,\in\, I_q \\ \operatorname{sgn}(t) q^{m_4} y \,\in\, I_q}} & \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\, (-1)^{\chi(xy)}\, |xy| a_p(\operatorname{sgn}(p) q^{m_3} x, x) \\ \times\ a_t(\operatorname{sgn}(t) q^{m_4} y, y)\, & (\iota \otimes \omega_{f_{m_1+\chi(x/y), \operatorname{sgn}(p) q^{m_3} x, \operatorname{sgn}(t) q^{m_4} y}, f_{m_2+\chi(x/y), x, y}})(W^*)\, f_{n,c,d}\ .\end{aligned}$$

By Eq. (4.4), we get immediately that $(\iota\otimes\omega)(W^*)\, f_{n,c,d} = 0$ if $\mathrm{sgn}(pt)cq^{-m_1} \notin I_q$ or $m_3 - m_4 \neq m_2 - m_1$. Now suppose that $\mathrm{sgn}(pt)cq^{-m_1} \in I_q$ and $m_3 - m_4 = m_2 - m_1$. Then Eq. (4.4) and Proposition 3.5 imply that

$$\begin{aligned}
&(\iota\otimes\omega)(W^*)\, f_{n,c,d} \\
&= \sum_{\substack{x,y\in I_q \\ \mathrm{sgn}(p)q^{m_3}x\in I_q \\ \mathrm{sgn}(t)q^{m_4}y\in I_q}} \mathrm{sgn}(x)^{\chi(x)}\,\mathrm{sgn}(y)^{\chi(y)}\,(-1)^{\chi(xy)}\,|xy| a_p(\mathrm{sgn}(p)q^{m_3}x,x)\,a_t(\mathrm{sgn}(t)q^{m_4}y,y) \\
&\qquad\times\; q^{m_4}\,a_{\mathrm{sgn}(t)q^{m_4}y}(c,y)\,a_{\mathrm{sgn}(p)q^{m_3}x}(\mathrm{sgn}(pt)cq^{-m_1},x)\,f_{m_1-m_2+n,\mathrm{sgn}(pt)cq^{-m_1},d} \\
&= \sum_{\substack{x,y\in I_q \\ \mathrm{sgn}(p)q^{m_3}x\in I_q \\ \mathrm{sgn}(t)q^{m_4}y\in I_q}} \mathrm{sgn}(x)^{\chi(x)}\,\mathrm{sgn}(y)^{\chi(y)}\,(-1)^{\chi(xy)}\,|xy| a_p(\mathrm{sgn}(p)q^{m_3}x,x)\,a_t(\mathrm{sgn}(t)q^{m_4}y,y) \\
&\qquad\times\; q^{m_4}\,(-1)^{\chi(cy)+m_4}\,\mathrm{sgn}(y)^{\chi(y)}\,|c/q^{m_4}y|\,a_c(\mathrm{sgn}(t)q^{m_4}y,y)\,(-1)^{\chi(cx)+m_3+m_1} \\
&\qquad\times\; \mathrm{sgn}(x)^{\chi(x)}\,|cq^{-m_1}/q^{m_3}x|\,a_{\mathrm{sgn}(pt)cq^{-m_1}}(\mathrm{sgn}(p)q^{m_3}x,x)\,f_{m_1-m_2+n,\mathrm{sgn}(pt)cq^{-m_1},d} \\
&= \sum_{\substack{x,y\in I_q \\ \mathrm{sgn}(p)q^{m_3}x\in I_q \\ \mathrm{sgn}(t)q^{m_4}y\in I_q}} c^2\,q^{-m_1-m_3}\,(-1)^{m_1+m_3+m_4}\,a_p(\mathrm{sgn}(p)q^{m_3}x,x)\,a_{\mathrm{sgn}(pt)cq^{-m_1}}(\mathrm{sgn}(p)q^{m_3}x,x) \\
&\qquad\times\; a_t(\mathrm{sgn}(t)q^{m_4}y,y)\,a_c(\mathrm{sgn}(t)q^{m_4}y,y)\,f_{m_1-m_2+n,\mathrm{sgn}(pt)cq^{-m_1},d}\;.
\end{aligned}$$

This sum is absolutely convergent so we can compute it in any order we deem useful. Therefore the orthogonality relations in Proposition 3.2 imply that

$$\begin{aligned}
(\iota\otimes\omega)(W^*)\, f_{n,c,d} &= c^2\,q^{-m_1-m_3}\,(-1)^{m_1+m_3+m_4}\,\delta_{p,\mathrm{sgn}(pt)cq^{-m_1}}\,\delta_{t,c}\;f_{n+m_1-m_2,\mathrm{sgn}(pt)cq^{-m_1},d} \\
&= t^2\,q^{-m_1-m_3}\,(-1)^{m_1+m_3+m_4}\,\delta_{|p|,|t|q^{-m_1}}\,\delta_{t,c}\;f_{n+m_1-m_2,p,d} \\
&= t^2\,q^{-m_1-m_3}\,(-1)^{m_1+m_3+m_4}\,\delta_{|p|,|t|q^{-m_1}}\,\pi(\Phi(m_1-m_2,p,t))\,f_{n,c,d}\;, \qquad (4.6)
\end{aligned}$$

where π is the GNS-representation of φ. If $\mathrm{sgn}(pt)cq^{-m_1} \notin I_q$, then

$$\delta_{|p|,|t|q^{-m_1}}\,\pi(\Phi(m_1-m_2,p,t))\,f_{n,c,d} = \delta_{|p|,|t|q^{-m_1}}\,\delta_{t,c}\,f_{m_1-m_2+n,p,d} = 0\;,$$

implying that Eq. (4.6) also holds if $\mathrm{sgn}(pt)cq^{-m_1} \notin I_q$ and $m_3 - m_4 = m_2 - m_1$. Therefore the lemma follows. □

Now we can easily calculate the action of S on elements of the form $\Phi(m,p,t)$.

Proposition 4.13. *Consider $m \in \mathbb{Z}$ and $p,t \in I_q$. Then $\Phi(m,p,t) \in D(S)$ and $S(\Phi(m,p,t)) = \mathrm{sgn}(p)^{\chi(p)}\,\mathrm{sgn}(t)^{\chi(t)}\,(-1)^m\,q^m\,\Phi(m,t,p)$.*

Proof. Define $\omega \in B(K)_*$ as in Eq. (4.5) for $m_1 = \chi(t/p)$, $m_2 = \chi(t/p) - m$, $m_3 = -\chi(t/p)$ and $m_4 = -\chi(t/p) + m$. Then Proposition 4.12 implies that $\pi(\Phi(m,p,t)) = t^{-2}\,(-1)^{m+\chi(pt)}\,(\iota\otimes\omega)(W^*)$. Thus [18, Prop. 8.3] implies that $\Phi(m,p,t) \in D(S^{-1})$ and

$$\pi\big(S^{-1}(\Phi(m,p,t))\big) = t^{-2}\,(-1)^{m+\chi(pt)}\,(\iota\otimes\omega)(W) = t^{-2}\,(-1)^{m+\chi(pt)}\,(\iota\otimes\bar{\omega})(W^*)^*\;, \qquad (4.7)$$

where, as is customary, $\bar{\omega} \in B(K)_*$ is given by $\bar{\omega}(a) = \overline{\omega(a^*)}$ for all $a \in B(K)$. By definition,

$$\begin{aligned}
\omega = &\sum_{\substack{x,y\in I_q \\ (p/|t|)x\in I_q \\ (|p|/t)q^m y\in I_q}} \mathrm{sgn}(x)^{\chi(x)}\,\mathrm{sgn}(y)^{\chi(y)}\,(-1)^{\chi(xy)}\,|xy| a_p((p/|t|)x,x)\,a_t((|p|/t)q^m y,y) \\
&\qquad\times\; \omega_{f_{\chi(t/p)+\chi(x/y),(p/|t|)x,(|p|/t)q^m y},f_{-m+\chi(t/p)+\chi(x/y),x,y}}\;.
\end{aligned}$$

Thus,

$$\begin{aligned}\bar{\omega} \;&=\; \sum_{\substack{x,\,y\,\in\, I_q \\ (p/|t|)x\,\in\, I_q \\ (|p|/t)q^m y\,\in\, I_q}} \operatorname{sgn}(x)^{\chi(x)} \operatorname{sgn}(y)^{\chi(y)}\,(-1)^{\chi(xy)}\,|xy|\,a_p((p/|t|)x,x)\,a_t((|p|/t)q^m y,y) \\ &\qquad\qquad \times\;\; \omega_{f_{-m+\chi(t/p)+\chi(x/y),x,y},f_{\chi(t/p)+\chi(x/y),(p/|t|)x,(|p|/t)q^m y}} \\ &=\; \sum_{\substack{x,\,y\,\in\, I_q \\ (p/|t|)x\,\in\, I_q \\ (|p|/t)q^m y\,\in\, I_q}} (-1)^{\chi(xy)}\,|xy|\operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}((p/|t|)x)^{\chi((p/|t|)x)}\,a_p(x,(p/|t|)x) \\ &\qquad\qquad \times\;\; \operatorname{sgn}(t)^{\chi(t)}\operatorname{sgn}((|p|/t)q^m y)^{\chi((|p|/t)q^m y)}\,a_t(y,(|p|/t)q^m y) \\ &\qquad\qquad \times\;\; \omega_{f_{-m+\chi(t/p)+\chi(x/y),x,y},f_{\chi(t/p)+\chi(x/y),(p/|t|)x,(|p|/t)q^m y}}\ ,\end{aligned}$$

where we used Proposition 3.5 in the last equation. A simple change in summation variables $x' = (p/|t|)x$ and $y' = (|p|/t)yq^m$ then reveals that

$$\begin{aligned}\bar{\omega} = \sum_{\substack{x',\,y'\,\in\, I_q \\ (|t|/p)x'\,\in\, I_q \\ (t/|p|)q^{-m}y'\,\in\, I_q}} &\operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,q^{-m}(t^2/p^2)\operatorname{sgn}(x')^{\chi(x')}\operatorname{sgn}(y')^{\chi(y')}\,(-1)^{\chi(x'y')}\,|x'y'| \\ &\times\;\; a_p((|t|/p)x',x')\,a_t((t/|p|)q^{-m}y',y') \\ &\times\;\; \omega_{f_{\chi(t/p)+\chi(x'/y'),(|t|/p)x',(t/|p|)q^{-m}y'},f_{m+\chi(t/p)+\chi(x'/y'),x',y'}}\ ,\end{aligned}$$

which upon close inspection shows that $\bar{\omega}$ is $\operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,q^{-m}(t^2/p^2)$ times the functional in Eq. (4.5) where $m_1 = \chi(t/p)$, $m_2 = \chi(t/p) + m$, $m_3 = \chi(t/p)$ and $m_4 = \chi(t/p) - m$. By Proposition 4.12 this implies that

$$\begin{aligned}(\iota \otimes \bar{\omega})(W^*) &= \operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,q^{-m}(t^2/p^2)\,t^2\,(p^2/t^2)\,(-1)^{m+\chi(pt)}\,\pi(\Phi(-m,p,t)) \\ &= \operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,q^{-m}\,t^2\,(-1)^{m+\chi(pt)}\,\pi(\Phi(-m,p,t))\ .\end{aligned}$$

Therefore Eq. (4.7) implies that

$$S^{-1}(\Phi(m,p,t)) = \operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,q^{-m}\,\Phi(m,t,p)\ ,$$

proving the result. ☐

It is now possible to recognize the polar decomposition of the antipode. For this purpose, define the anti-unitary operator I on H and the strictly positive operator Q in H such that $\langle\, \zeta^n \otimes \delta_x \mid n \in \mathbb{Z}, x \in I_q \,\rangle$ is a core for Q and $I(\zeta^n \otimes \delta_x) = (-1)^n \operatorname{sgn}(x)^{\chi(x)}\,\zeta^{-n} \otimes \delta_x$ and $Q(\zeta^n \otimes \delta_x) = q^{2n}\,\zeta^n \otimes \delta_x$ for all $n \in \mathbb{Z}$ and $x \in I_q$. Recall that the unitary antipode R and scaling group τ are introduced after the proof of [19, Prop. 1.4].

Proposition 4.14. *The unitary antipode R and the scaling group τ for (M_q, Δ) are such that $R(a) = Ia^*I$ and $\tau_s(a) = Q^{is}aQ^{-is}$ for all $a \in M_q$ and $s \in \mathbb{R}$. Let $m \in \mathbb{Z}$ and $p, t \in I_q$. Then $R(\Phi(m,p,t)) = \operatorname{sgn}(p)^{\chi(p)}\operatorname{sgn}(t)^{\chi(t)}\,(-1)^m\,\Phi(m,t,p)$. If $z \in \mathbb{C}$, then the element $\Phi(m,p,t)$ belongs to $D(\tau_z)$ and $\tau_z(\Phi(m,p,t)) = q^{2miz}\,\Phi(m,p,t)$.*

Proof. By Proposition 4.11 and the discussion after the proof of [19, Prop. 1.4] we know that there exists a strictly positive operator P on K such that $P^{is}\Lambda(a) = \Lambda(\tau_s(a))$ for all $a \in \mathcal{N}_\varphi$ and $s \in \mathbb{R}$. Note that $\pi(\tau_s(a)) = P^{is}\pi(a)P^{-is}$ for all $s \in \mathbb{R}$.
Choose $m \in \mathbb{Z}$ and $p, t \in \mathbb{R}$. Since $S = R\tau_{-\frac{i}{2}}$ and R and τ commute, we see that $\tau_{-i} = S^2$. Thus, Proposition 4.13 implies that $\Phi(m,p,t) \in D(\tau_{-i})$ and $\tau_{-i}(\Phi(m,p,t)) = q^{2m}\,\Phi(m,p,t)$. Thus, Lemma 4.2 and the von Neumann algebraic version of [16, Prop. 4.4] guarantee that $\Lambda(\Phi(m,p,t)) \in D(P)$ and $P\,\Lambda(\Phi(m,p,t)) = q^{2m}\,\Lambda(\Phi(m,p,t))$. Therefore, Lemma 4.2 tells us that $f_{m,p,t} \in D(P)$ and $Pf_{m,p,t} = q^{2m}\,f_{m,p,t} = (Q\otimes 1)f_{m,p,t}$. Because such elements $f_{m,p,t}$ form a core for $Q\otimes 1$, we conclude that $Q\otimes 1 \subseteq P$. Taking the adjoint of this inclusion, we see that $P \subseteq Q\otimes 1$. As a consequence, $P = Q\otimes 1$. So if $a \in M_q$ and $s \in \mathbb{R}$, we see that $\tau_s(a)\otimes 1 = \pi(\tau_s(a)) = P^{is}\pi(a)P^{-is} = Q^{is}aQ^{-is}\otimes 1$, implying that $\tau_s(a) = Q^{is}aQ^{-is}$.

Note that $Q^{is}(\zeta^n \otimes \delta_x) = q^{2nis}\, \zeta^n \otimes \delta_x$ for all $n \in \mathbb{Z}$, $x \in I_q$ and $s \in \mathbb{R}$. Now a straightforward calculation reveals that $\tau_s(\Phi(m,p,t)) = Q^{is}\Phi(m,p,t)Q^{-is} = q^{2mis}\, \Phi(m,p,t)$ for all $m \in \mathbb{Z}$, $p,t \in I_q$ and $s \in \mathbb{R}$.
Fix $m \in \mathbb{Z}$ and $p,t \in I_q$ for the moment. By definition $S = R\tau_{-\frac{i}{2}}$. The previous paragraph implies that $\tau_{-\frac{i}{2}}(\Phi(m,p,t)) = q^m\, \Phi(m,p,t)$. Therefore Proposition 4.13 guarantees that $R(\Phi(m,p,t)) = q^{-m}\, S(\Phi(m,p,t)) = \operatorname{sgn}(p)^{\chi(p)} \operatorname{sgn}(t)^{\chi(t)}\, (-1)^m\, \Phi(m,t,p)$. Using the definition of I, one checks easily that also $I\Phi(m,p,t)^* I = I\Phi(-m,t,p)I = \operatorname{sgn}(p)^{\chi(p)} \operatorname{sgn}(t)^{\chi(t)}\, (-1)^m\, \Phi(m,t,p)$ and thus by the previous calculation, $R(\Phi(m,p,t)) = I\Phi(m,p,t)^* I$. Since the von Neumann algebra M_q is generated by such elements $\Phi(m,p,t)$, we conclude that $R(a) = Ia^*I$ for all $a \in M_q$. □

One easily checks that R and $\breve{R}$ are connected through the formula $\breve{R}(\Phi(m,p,t)) = (-1)^m\, R(\Phi(m,p,t))$ if $m \in \mathbb{Z}$ and $p,t \in I_q$.

Consider $z \in \mathbb{C}$. Since the *-algebra $\langle\, \Phi(m,p,t) \mid m \in \mathbb{Z}, p,t \in I_q \,\rangle$ is σ-strongly* dense in M_q and this same *-algebra is clearly invariant under each τ_s, where $s \in \mathbb{R}$, it follows from the von Neumann algebraic version of [17, Cor. 1.22] that $\langle\, \Phi(m,p,t) \mid m \in \mathbb{Z}, p,t \in I_q \,\rangle$ is a σ-strong* core for τ_z.
If $n \in \mathbb{Z}$, the fact that $S^n = R\,\tau_{-n\frac{i}{2}}$ implies that $\langle\, \Phi(m,p,t) \mid m \in \mathbb{Z}, p,t \in I_q \,\rangle$ is a σ-strong* core for S^n.

Following [19, Prop. 1.6] we associate to the von Neumann algebraic quantum group (M_q, Δ) a reduced C*-algebraic quantum group in the sense of [18]. Therefore we define the C*-subalgebra A_q of M_q such that $\pi(A_q)$ is the norm closure, in $B(K)$, of the set $\{\, (\iota \otimes \omega)(W^*) \mid \omega \in B(H)_* \,\}$. Then [19, Prop. 1.6] guarantees that $(A_q, \Delta\restriction_{A_q})$ is a reduced C*-algebraic quantum group in the sense of [18]. In the next proposition we give an explicit description for A_q.

We will use the following notation. For $f \in C(\mathbb{T} \times I_q)$ and $x \in I_q$ we define $f_x \in C(\mathbb{T})$ such that $f_x(\lambda) = f(\lambda, x)$ for all $\lambda \in \mathbb{T}$.

Proposition 4.15. *Denote by $\mathcal{C}$ the C*-algebra of all functions $f \in C(\mathbb{T} \times I_q)$ such that* (1) *f_x converges uniformly to* 0 *as $x \to 0$ and* (2) *f_x converges uniformly to a constant function as $x \to \infty$. Then A_q is the norm closed linear span, in $B(H)$, of the set $\{\, \rho_p\, M_f \mid f \in \mathcal{C}, p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}} \,\}$.*

Proof. Let $p,t \in I_q$ and define $F_{p,t} \in \mathcal{F}(I_q)$ such that $F_{p,t}(x) = a_p(x,t)$ for all $x \in I_q$. The reason for introducing the function $F_{p,t}$ stems from the fact that

$$\begin{aligned}(\iota \otimes \omega_{f_{m_1,p_1,t_1}, f_{m_2,p_2,t_2}})(W^*) \\ = \delta_{\chi(p_1t_2/p_2t_1), m_2-m_1}\ |t_1/t_2|\ \pi\Big(\ \rho_{\operatorname{sgn}(p_1t_1)(t_2/p_2)q^{m_1}}\ M_{\zeta^{m_1-m_2} \otimes T_{\operatorname{sgn}(p_1t_1)(p_2/t_2)q^{-m_1}}(F_{p_1,p_2})F_{t_1,t_2}}\ \Big)\end{aligned} \tag{4.8}$$

for $p_1, p_2, t_1, t_2 \in I_q$ and $m_1, m_2 \in \mathbb{Z}$ (and which follows from Eq. (4.4)). Since $B(K)_*$ is the closed linear span of vector functionals, A_q is the closed linear span of elements of the form (4.8).

Let $p,t \in I_q$. Take $m \in \mathbb{Z}$ such that $|p/t| = q^m$. We consider two different cases:

$\operatorname{sgn}(\boldsymbol{p}) = \operatorname{sgn}(\boldsymbol{t})$ **:** Note first of all that in this case, $F_{p,t}(x) = 0$ for all $x \in I_q^-$. Definition 3.1 implies the existence of a bounded function $h : I_q^+ \to \mathbb{R}$ such that

$$F_{p,t}(x) = h(x)\, \frac{\nu(pt/x)}{\sqrt{(-\kappa(x); q^2)_\infty}}\ \Psi(-q^2/\kappa(t); q^2\kappa(x/t); q^2\kappa(x/p))$$

for all $x \in I_q^+$. If $x \to 0$, we have that $\nu(pt/x) \to 0$, $(-\kappa(x); q^2)_\infty \to 1$ and $\Psi(-q^2/\kappa(t); q^2\kappa(x/t); q^2\kappa(x/p)) \to \Psi(-q^2/\kappa(t); 0; 0)$. It follows that $F_{p,t}(x) \to 0$ as $x \to 0$.

Proposition 3.5 and Definition 3.1 imply for $x \in I_q^+$,

$$\begin{aligned}F_{p,t}(x) = a_p(x,t) = \operatorname{sgn}(p)^{\chi(pt)}\, a_p(t,x) = (-1)^{\chi(pt)} \operatorname{sgn}(p)^{\chi(pt)}\, c_q\, (-\kappa(p); q^2)_\infty^{\frac{1}{2}} \\ (-\kappa(t); q^2)_\infty^{-\frac{1}{2}}\ x\, \nu(px/t)\, (-\kappa(x); q^2)_\infty^{\frac{1}{2}}\ \Psi(-q^2/\kappa(x); q^2\kappa(t/x); q^2\kappa(t/p))\end{aligned} \tag{4.9}$$

If $x \to \infty$, then $\Psi(-q^2/\kappa(x); q^2\kappa(t/x); q^2\kappa(t/p)) \to \Psi(0; 0; q^2\kappa(t/p)) = (q^2\kappa(t/p); q^2)_\infty$, where we used [6, Eq. (1.3.16)]. (4.10)

Let $x \in I_q^+$ and choose $s \in \mathbb{Z}$ such that $x = q^s$. Then Result 6.1 implies that

$$\begin{aligned} x\,\nu(px/t)\,(-\kappa(x);q^2)_\infty^{\frac{1}{2}} &= q^s\, q^{\frac{1}{2}(s+m-1)(s+m-2)}\,\sqrt{(-q^{2s};q^2)_\infty} \\ &= q^s\, q^{\frac{1}{2}s(s-1)}\, q^{\frac{1}{2}(m-1)(m-2)}\, q^{(m-1)s}\, q^{-\frac{1}{2}s(s-1)}\,(-1,-q^2;q^2)_\infty^{\frac{1}{2}}\,(-q^{2(1-s)};q^2)_\infty^{-\frac{1}{2}} \\ &= q^{\frac{1}{2}(m-1)(m-2)}\,\sqrt{2}\,(-q^2;q^2)_\infty\, x^m\,(-q^2/x^2;q^2)_\infty^{-\frac{1}{2}}\ . \end{aligned}$$

So if $p = t$ (and thus $m = 0$), $x\,\nu(px/t)\,(-\kappa(x);q^2)_\infty^{\frac{1}{2}} \rightarrow \sqrt{2}\,q\,(-q^2;q^2)_\infty$ as $x \rightarrow \infty$. If, on the other hand, $p > t$ (and thus $m < 0$), we see that $x\,\nu(px/t)\,(-\kappa(x);q^2)_\infty^{\frac{1}{2}} \rightarrow 0$ as $x \rightarrow \infty$. If we combine this with Eq. (4.9) and the convergence in (4.10), we conclude, since $c_q^{-1} = \sqrt{2}\,q\,(q^2,-q^2;q^2)_\infty$, that

(1) $F_{p,p}(x) \rightarrow 1$ as $x \rightarrow \infty$,
(2) if $p > t$, then $F_{p,t}(x) \rightarrow 0$ as $x \rightarrow \infty$.

But Proposition 3.5 implies that $|F_{p,t}| = |t/p|\,|F_{t,p}|$. By the convergence in (2), this implies that if $p < t$, also $F_{p,t}(x) \rightarrow 0$ as $x \rightarrow \infty$.

$\mathrm{sgn}(\boldsymbol{p}) \neq \mathrm{sgn}(\boldsymbol{t})$: Note that in this case, $F_{p,t}(x) = 0$ for all $x \in I_q^+$. Completely similar as in the beginning of the first part of this proof, one shows that $F_{p,t}(x) \rightarrow 0$ as $x \rightarrow 0$.

From the previous discussion we only have to remember that $F_{p,t}(x) \rightarrow 0$ as $x \rightarrow 0$. Provided $p \neq t$, we have that $F_{p,t}(x) \rightarrow 0$ as $x \rightarrow \infty$. And $F_{p,p}(x) \rightarrow 1$ as $x \rightarrow \infty$. (4.11)

Let B denote the norm closed linear span of $\{\,\rho_p\, M_f \mid f \in \mathcal{C}, p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}\,\}$ in $B(H)$.
Use the notation of Eq. (4.8). If $T_{\mathrm{sgn}(p_1 t_1)(p_2/t_2)q^{-m_1}}(F_{p_1,p_2})F_{t_1,t_2} \in C_0(I_q)$, then Eqs. (4.8) and (4.11) immediately guarantees that $(\iota \otimes \omega_{f_{m_1,p_1,t_1},f_{m_2,p_2,t_2}})(W^*) \in \pi(B)$. If, on the other hand, $T_{\mathrm{sgn}(p_1 t_1)(p_2/t_2)q^{-m_1}}(F_{p_1,p_2})F_{t_1,t_2} \notin C_0(I_q)$, we must by (4.11) necessarily have that $t_1 = t_2$ and $p_1 = p_2$. Thus, Eq. (4.8) implies in this case that

$$\begin{aligned} (\iota \otimes \omega_{f_{m_1,p_1,t_1},f_{m_2,p_1,t_1}})(W^*) &= \delta_{0,m_2-m_1}\,\pi\Big(\,\rho_{q^{m_1}|t_1/p_1|}\, M_{\zeta^{m_1-m_2}\otimes T_{q^{-m_1}|p_1/t_1|}(F_{p_1,p_1})F_{t_1,t_1}}\,\Big) \\ &= \delta_{0,m_2-m_1}\,\pi\Big(\,\rho_{q^{m_1}|t_1/p_1|}\, M_{1\otimes T_{q^{-m_1}|p_1/t_1|}(F_{p_1,p_1})F_{t_1,t_1}}\,\Big) \in \pi(B)\ , \end{aligned}$$

where we used (4.11) in the last relation. Hence $\pi(A_q) \subseteq \pi(B)$, thus $A_q \subseteq B$.

Proposition 4.12 implies that $\rho_{t^{-1}p}\, M_{\zeta^m \otimes \delta_p} = \Phi(m,t,p) \in A_q$ for all $p,t \in I_q$ and $m \in \mathbb{Z}$. It follows that $\rho_x\, M_{\zeta^m \otimes \delta_p} \in A_q$ for all $m \in \mathbb{Z}$, $x \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and $p \in I_q$. Thus, $\rho_x\, M_f \in A_q$ for all $x \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and $f \in C_0(\mathbb{T} \times I_q)$.
If $x \in -q^{\mathbb{Z}}$ and $f \in \mathcal{C}$, it is easy to see that there exists $g \in C_0(\mathbb{T} \times I_q)$ such that $\rho_x\, M_f = \rho_x\, M_g$. Hence, $\rho_x\, M_f \in A_q$ in this case.
Now fix $x \in q^{\mathbb{Z}}$ and $f \in \mathcal{C}$. Take $m \in \mathbb{Z}$ such that $x = q^m$. Because $f \in \mathcal{C}$, there exists $c \in \mathbb{C}$ such that $f_y \rightarrow c\,1_{\mathbb{T}}$ uniformly as $y \rightarrow \infty$. Consequently, $f - c\,(1_{\mathbb{T}} \otimes T_{x^{-1}}(F_{1,1})F_{1,1})$ belongs to $C_0(\mathbb{T} \times I_q)$ by (4.11). As such, $\rho_x\, M_{f-c\,(1_{\mathbb{T}}\otimes T_{x^{-1}}(F_{1,1})F_{1,1})} \in A_q$. Moreover, Eq. (4.8) guarantees that $\pi(\rho_x\, M_{1\otimes T_{x^{-1}}(F_{1,1})F_{1,1}}) = (\iota \otimes \omega_{f_{m,1,1},f_{m,1,1}})(W^*) \in \pi(A_q)$. Hence $\rho_x\, M_{1\otimes T_{x^{-1}}(F_{1,1})F_{1,1}} \in A_q$. It follows that $\rho_x\, M_f \in A_q$. Hence $B \subseteq A_q$, proving that $B = A_q$. $\square$

5. PROOF OF THE COASSOCIATIVITY OF THE COMULTIPLICATION

This section is devoted to the proof of Theorem 3.11. First one proves that $(\Delta \otimes \iota)\Delta(\gamma) = (\iota \otimes \Delta)\Delta(\gamma)$. It is easy to prove that both operators agree on the intersection of their domain, but it takes a lot of work to establish the equality of their domains. In order to enhance the continuity of the exposition, we give the proof in the second half of this section. Once we have proven that $(\Delta \otimes \iota)\Delta(\gamma) = (\iota \otimes \Delta)\Delta(\gamma)$, it is straightforward to show that $(\Delta \otimes \iota)\Delta(x) = (\iota \otimes \Delta)\Delta(x)$ for all $x \in M_q^{\diamond}$. In the final step of the proof of the coassociativity it is shown that $(\Delta \otimes \iota)\Delta(u) = (\iota \otimes \Delta)\Delta(u)$.

Since $\Delta(x) = V^*(1_{L^2(\mathbb{T}^2)} \otimes x)V$ for all $x \in M_q$, we get for any closed densely defined operator T in H, affiliated with M_q,

$$(\Delta \otimes \iota)\Delta(T) = (V^* \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes T)(1_{L^2(\mathbb{T}^2)} \otimes V)(V \otimes 1_H) \tag{5.1}$$

and

$$(\iota \otimes \Delta)\Delta(T) = (1_H \otimes V^*)V_{13}^*(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes T)V_{13}(1_H \otimes V) \tag{5.2}$$

where $V_{13} : H \otimes L^2(\mathbb{T}^2) \otimes H \to L^2(\mathbb{T}^2) \otimes L^2(\mathbb{T}^2) \otimes H$ is defined as usual.

Using the map $\Delta_0^{(2)} : \mathcal{A}_q \to \mathcal{A}_q \odot \mathcal{A}_q \odot \mathcal{A}_q$ defined by $\Delta_0^{(2)} = (\Delta_0 \odot \iota)\Delta_0 = (\iota \odot \Delta_0)\Delta_0$, we get adjointable operators $\Delta_0^{(2)}(\alpha_0)$, $\Delta_0^{(2)}(\alpha_0^\dagger)$, $\Delta_0^{(2)}(\gamma_0)$, $\Delta_0^{(2)}(\gamma_0^\dagger)$ in $\mathcal{L}^+(E \odot E \odot E)$. Proposition 3.9 will imply the next result.

Proposition 5.1. *If $T = \alpha$ or $T = \gamma$, then*

$$\begin{array}{lcllcl} \Delta_0^{(2)}(T_0) & \subseteq & (\Delta \otimes \iota)\Delta(T) & \qquad \Delta_0^{(2)}(T_0) & \subseteq & (\iota \otimes \Delta)\Delta(T) \\ \Delta_0^{(2)}(T_0^\dagger) & \subseteq & (\Delta \otimes \iota)\Delta(T^*) & \qquad \Delta_0^{(2)}(T_0^\dagger) & \subseteq & (\iota \otimes \Delta)\Delta(T^*)\ . \end{array}$$

Proof. We only proof the inclusion for $T = \alpha$. The other ones are dealt with in the same way. Proposition 3.9 tells us that $\alpha_0 \odot \alpha_0 + q\, e_0\gamma_0^\dagger \odot \gamma_0 \subseteq V^*(1_{L^2(\mathbb{T}^2)} \otimes \alpha)V$ and thus,

$$1_{L^2(\mathbb{T}^2)} \odot \alpha_0 \odot \alpha_0 + q\, 1_{L^2(\mathbb{T}^2)} \odot e_0\gamma_0^\dagger \odot \gamma_0 \subseteq (1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)(1_{L^2(\mathbb{T}^2)} \otimes V)\ . \tag{5.3}$$

By Lemma 2.1, $D(1_{L^2(\mathbb{T}^2)} \otimes \alpha) = D(1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*)$. Let $v \in D(1_{L^2(\mathbb{T}^2)} \otimes \alpha)$ and $w \in E$. There exists a sequence $(v_n)_{n=1}^\infty$ in $L^2(\mathbb{T}^2) \odot E$ such that $(v_n)_{n=1}^\infty \to v$ and $\big(\,(1_{L^2(\mathbb{T}^2)} \odot \alpha_0)(v_n)\,\big)_{n=1}^\infty \to (1_{L^2(\mathbb{T}^2)} \otimes \alpha)(v)$. Since $(1_{L^2(\mathbb{T}^2)} \odot \alpha_0^\dagger)(1_{L^2(\mathbb{T}^2)} \odot \alpha_0) = (1_{L^2(\mathbb{T}^2)} \odot e_0\gamma_0)(1_{L^2(\mathbb{T}^2)} \odot e_0\gamma_0^\dagger) + 1_{L^2(\mathbb{T}^2)} \odot e_0$, $1_{L^2(\mathbb{T}^2)} \odot e_0$ is bounded and $1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*$ is closed, it follows that $\big(\,(1_{L^2(\mathbb{T}^2)} \odot e_0\gamma_0^\dagger)(v_n)\,\big)_{n=1}^\infty \to (1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*)(v)$.
Therefore the net

$$\big(\,(1_{L^2(\mathbb{T}^2)} \odot \alpha_0 \odot \alpha_0 + q\, 1_{L^2(\mathbb{T}^2)} \odot e_0\gamma_0^\dagger \odot \gamma_0)(v_n \otimes w)\,\big)_{n=1}^\infty$$

converges to $(\,(1_{L^2(\mathbb{T}^2)} \otimes \alpha) \odot \alpha_0 + q\,(1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*) \odot \gamma_0\,)(v \otimes w)$. Hence, inclusion (5.3) and the fact that the operator on the right hand side of this inclusion is closed imply that $v \otimes w \in D\big((1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)(1_{L^2(\mathbb{T}^2)} \otimes V)\big)$ and

$$(\,(1_{L^2(\mathbb{T}^2)} \otimes \alpha) \odot \alpha_0 + q\,(1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*) \odot \gamma_0)(v \otimes w) = (1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)(1_{L^2(\mathbb{T}^2)} \otimes V)(v \otimes w)\ .$$

In other words,

$$(1_{L^2(\mathbb{T}^2)} \otimes \alpha) \odot \alpha_0 + q\,(1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*) \odot \gamma_0 \subseteq (1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)(1_{L^2(\mathbb{T}^2)} \otimes V)\ .$$

As a consequence, proposition 3.9 implies that

$$\begin{aligned} \Delta_0^{(2)}(\alpha_0) &= \Delta_0(\alpha_0) \odot \alpha_0 + q\,\Delta_0(e_0\gamma_0^\dagger) \odot \gamma_0 \subseteq V^*(1_{L^2(\mathbb{T}^2)} \otimes \alpha)V \odot \alpha_0 + q\,V^*(1_{L^2(\mathbb{T}^2)} \otimes e\gamma^*)V \odot \gamma_0 \\ &\subseteq (V^* \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V^*)(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)(1_{L^2(\mathbb{T}^2)} \otimes V)(V \otimes 1_H) = (\Delta \otimes \iota)\Delta(\alpha)\ . \end{aligned}$$

□

For later purposes we introduce some extra notation: if $z, w \in \mathbb{C}$, we set

$$\phi(w; z) = \sum_{r=0}^{\infty} \frac{(wq^{2r}; q^2)_\infty}{(q^2; q^2)_r}\, q^{2r(r-1)}\, z^r\ . \tag{5.4}$$

Note that this ϕ-function is closely related to the ${}_0\varphi_2$-function (see [6, Def. (1.2.22)]).

The next step of the proof consists of showing that

Proposition 5.2. $(\Delta \otimes \iota)\Delta(\gamma) = (\iota \otimes \Delta)\Delta(\gamma)$.

The proof of this result is a tedious but not so difficult task and is given in the second half of this section, after Proposition 5.4. It is however important to remember that this result is needed for the last two parts (Corollary 5.3 and Proposition 5.4) of the proof of the coassociativity of Δ.

Recall that we introduced $M_q^\diamond$ after Definition 2.3.

Corollary 5.3. *We have that $(\Delta \otimes \iota)\Delta(x) = (\iota \otimes \Delta)\Delta(x)$ for all $x \in M_q^\diamond$.*

Proof. The previous proposition implies that $D\big((\Delta \otimes \iota)\Delta(\gamma)\big) = D\big((\iota \otimes \Delta)\Delta(\gamma)\big)$. (5.5)
By Eq. (5.1), we know that

$$D\big((\Delta \otimes \iota)\Delta(T)\big) = (V^* \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V^*)\, D(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes T)$$

for $T = \gamma$ and $T = \alpha$. Lemma 2.1 guarantees that $D(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \gamma) = D(1_{L^2(\mathbb{T}^2)} \otimes 1_{L^2(\mathbb{T}^2)} \otimes \alpha)$. As a consequence, $D\big((\Delta \otimes \iota)\Delta(\gamma)\big) = D\big((\Delta \otimes \iota)\Delta(\alpha)\big)$. In a similar way one shows that $D\big((\iota \otimes \Delta)\Delta(\gamma)\big) = D\big((\iota \otimes \Delta)\Delta(\alpha)\big)$. By Eq. (5.5), this guarantees that $D\big((\Delta \otimes \iota)\Delta(\alpha)\big) = D\big((\iota \otimes \Delta)\Delta(\alpha)\big)$.
Proposition 5.1 tells us that $\Delta_0^{(2)}(\alpha_0^\dagger) \subseteq (\Delta \otimes \iota)\Delta(\alpha^*)$ and $\Delta_0^{(2)}(\alpha_0^\dagger) \subseteq (\iota \otimes \Delta)\Delta(\alpha^*)$. Taking the adjoints of this inclusions, we see that $(\Delta \otimes \iota)\Delta(\alpha)$ and $(\iota \otimes \Delta)\Delta(\alpha)$ are both restrictions of $\Delta_0^{(2)}(\alpha_0^\dagger)^*$. The previous discussion learned us that also the domains of $(\Delta \otimes \iota)\Delta(\alpha)$ and $(\iota \otimes \Delta)\Delta(\alpha)$ agree. Thus, $(\Delta \otimes \iota)\Delta(\alpha) = (\iota \otimes \Delta)\Delta(\alpha)$.

Since $M_\zeta \otimes 1$ is the unitary in the polar decomposition of γ and $(\Delta\otimes\iota)\Delta(\gamma) = (\iota\otimes\Delta)\Delta(\gamma)$, the uniqueness of the polar decomposition implies that $(\Delta\otimes\iota)\Delta(M_\zeta\otimes 1) = (\iota\otimes\Delta)\Delta(M_\zeta\otimes 1)$ (*) and $(\Delta\otimes\iota)\Delta(|\gamma|) = (\iota\otimes\Delta)\Delta(|\gamma|)$. Since $\Delta(e) = e\otimes e$, $(\Delta\otimes\iota)\Delta(e) = (\iota\otimes\Delta)\Delta(e)$, so we get that $(\Delta\otimes\iota)\Delta(e\,|\gamma|) = (\iota\otimes\Delta)\Delta(e\,|\gamma|)$. Applying functional calculus to this equation, we see that $(\Delta \otimes \iota)\Delta(M_{1\otimes g}) = (\iota \otimes \Delta)\Delta(M_{1\otimes g})$ for all $g \in \mathcal{L}^\infty(I_q)$. Combining this with (*), it follows that $(\Delta \otimes \iota)\Delta(\pi(f)) = (\iota \otimes \Delta)\Delta(\pi(f))$ for all $f \in \mathcal{L}^\infty(\mathbb{T} \times I_q)$ (cf. the proof of Lemma 2.4).
Because w is the isometry in the polar decomposition of α and $(\Delta\otimes\iota)\Delta(\alpha) = (\iota\otimes\Delta)\Delta(\alpha)$, the uniqueness of the polar decomposition implies that $(\Delta\otimes\iota)\Delta(w) = (\iota\otimes\Delta)\Delta(w)$. The lemma follows by Lemma 2.4. $\square$

In order to finalize the proof of the coassociativity we rely on formulas discovered by S.L. Woronowicz. In particular, we borrow the use of the extended Hilbert space, the definition of $\tilde{\Theta}$ as in (5.10) and Eq. (5.11) in the proof below. These formulas can be found in [30, Sec. 5] (Eq. (5.11) is a slight variation of the one in [30], where e is replaced by 1). S.L. Woronowicz told us that he has a proof for these identities, not included in the version of [30] yet. Our proof of the strong convergence in (5.10) and Eq. (5.11) are probably different from his ones, due to the difference of both approaches. The proof of the equality $(\Delta \otimes \iota)\Delta(u) = (\iota \otimes \Delta)\Delta(u)$ is not present in [30], but it is possible that S.L. Woronowicz also knows a proof for this. However, the property of tensor products of $\widetilde{SU}_q(1,1)$-quadruples that corresponds to the equality $D((\Delta \otimes \iota)\Delta(\gamma)) = D((\iota \otimes \Delta)\Delta(\gamma))$ is still not proven within the approach of [30].

Proposition 5.4. *The *-homomorphism $\Delta : M_q \to M_q \otimes M_q$ is coassociative.*

Proof. Since M_q is generated by $M_q^\diamond \cup \{u\}$, it only remains to show that $(\Delta \otimes \iota)\Delta(u) = (\iota \otimes \Delta)\Delta(u)$ by the previous corollary. Define first of all the isometry $v \in M_q^\diamond$ as $v = e\, w\, (M_\zeta^* \otimes 1)$.
Set $K_q = -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ and use the uniform measure on K_q. Define $\tilde{H} = L^2(\mathbb{T}) \otimes L^2(K_q)$. Then $\tilde{H}$ is obviously the orthogonal sum of H and $L^2(\mathbb{T}) \otimes L^2(K_q \setminus I_q)$.
Let L be a Hilbert space. Then we define a normal *-homomorphism $B(H \otimes L) \to B(\tilde{H} \otimes L) : a \to \hat{a}$ such that $\hat{a}\restriction_{H\otimes L} = a$ and $\hat{a}\restriction_{(H\otimes L)^\perp} = 0$. If L_1, L_2 are two Hilbert spaces, $(b \otimes c)\hat{\ } = \hat{b} \otimes c$ for all $b \in B(H \otimes L_1)$, $c \in B(L_2)$. (5.6)
On $\tilde{H}$ we define unitary operators $\tilde{e}$, $\tilde{w}$ and $\tilde{u}$ such that for $f \in L^2(\mathbb{T} \times K_q)$,

$$(\tilde{e} f)(\lambda, x) = \operatorname{sgn}(x)\, f(\lambda, x) \qquad (\tilde{w} f)(\lambda, x) = f(\lambda, q^{-1}x) \qquad (\tilde{u} f)(\lambda, x) = f(\lambda, -x)$$

for almost all $(\lambda, x) \in \mathbb{T} \times K_q$. Now define the unitary element $\tilde{v}$ in $B(\tilde{H})$ as $\tilde{v} = \tilde{e}\, \tilde{w}\, (M_\zeta^* \otimes 1)$.
Since $(L^2(\mathbb{T}) \otimes L^2(K_q \setminus I_q)) \otimes H$ is separable, we can extend the orthonormal basis for $H \otimes H$ given by the functions of Definition 3.6 to an orthonormal basis $(\, F_{r,s,m,p} \mid r, s, m \in \mathbb{Z}, p \in K_q\,)$ for $\tilde{H} \otimes H$. If $p \in K_q$, we define $\tilde{\delta}_p \in L^2(K_q)$ such that $\tilde{\delta}_p(x) = \delta_{x,p}$ for all $x \in K_q$. Now define the unitary operator $\tilde{V} : \tilde{H} \otimes H \to L^2(\mathbb{T}^2) \otimes \tilde{H}$ such that $\tilde{V} F_{r,s,m,p} = \zeta^r \otimes \zeta^s \otimes \zeta^m \otimes \tilde{\delta}_p$ for all $r, s, m \in \mathbb{Z}$ and $p \in K_q$. So the restriction of $\tilde{V}$ to $H \otimes H$ equals V and $\tilde{V}(H^\perp \otimes H) = L^2(\mathbb{T}^2) \otimes H^\perp$.
Define the normal injective *-homomorphism $\tilde{\Delta} : B(\tilde{H}) \to B(\tilde{H} \otimes H)$ such that $\tilde{\Delta}(a) = \tilde{V}^*(1_{L^2(\mathbb{T}^2)} \otimes a)\tilde{V}$ for all $a \in B(\tilde{H})$. Thus, if $a \in M_q$, then $\tilde{\Delta}(\hat{a}) = \Delta(a)\hat{\ }$. By Eq. (5.6), this implies that $(\tilde{\Delta} \otimes \iota)(\hat{b}) = (\Delta \otimes \iota)(b)\hat{\ }$ for all $b \in M_q \otimes B(H)$.

Take $r, s, m \in \mathbb{Z}$ and $p \in K_q$. Choose $n \in \mathbb{N}$. One easily checks that $\tilde{\Delta}(\tilde{v}) F_{r,s,m,p} = \operatorname{sgn}(p)\, F_{r,s,m-1,pq}$. Thus, $\tilde{\Delta}(\tilde{v})^n F_{r,s,m,p} = \operatorname{sgn}(p)^n\, F_{r,s,m-n,pq^n}$. Notice that $p\, q^n \in I_q$ if $n \geq 1 - \chi(p)$.

Assume that $n \geq 1-\chi(p)$. Choose $x \in K_q$, $y \in I_q$ and $\lambda, \mu \in \mathbb{T}$. If $q^n x \in I_q$ and $y/x = \operatorname{sgn}(p)\, q^m$, then $y/(q^n x) = \operatorname{sgn}(pq^n)\, q^{m-n}$ which by Definition 3.6 and Result 6.4 guarantees that

$$\begin{aligned}
&\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n \digamma_{r,s,m,p}\big)(\lambda,x,\mu,y) = \operatorname{sgn}(p)^n\,[(\tilde{v}^{-n}\otimes 1)\digamma_{r,s,m-n,pq^n}](\lambda,x,\mu,y) \\
&\quad = \operatorname{sgn}(p)^n \operatorname{sgn}(x)^n\,\lambda^n\,\digamma_{r,s,m-n,pq^n}(\lambda,q^n x,\mu,y) \\
&\quad = \operatorname{sgn}(y)^n\,\lambda^n\,a_{q^n p}(q^n x,y)\,\lambda^{r+\chi(y/pq^n)}\,\mu^{s-\chi(xq^n/pq^n)} \\
&\quad = \operatorname{sgn}(y)^n\,a_{q^n p}(q^n x,y)\,\lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)} \\
&\quad = \lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)}\operatorname{sgn}(y)^n\,c_q\,s(q^n x,y)\,(-1)^{\chi(q^n p)}\,(-\operatorname{sgn}(y))^{\chi(q^n x)}\,|y|\,\nu(q^n py/q^n x) \\
&\qquad \sqrt{\frac{(-\kappa(q^n p),-\kappa(y);q^2)_\infty}{(-\kappa(q^n x);q^2)_\infty}}\;\Psi\left(\begin{array}{c} -q^2/\kappa(q^n p) \\ q^2\kappa(q^n x/q^n p)\end{array};\,q^2\kappa(q^n x/y)\right) \\
&\quad = \lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)}\,c_q\,s(x,y)\,(-1)^{\chi(p)}\,(-\operatorname{sgn}(y))^{\chi(x)}\,|y|\,\nu(py/x) \\
&\qquad \sqrt{\frac{(-\kappa(q^n p),-\kappa(y);q^2)_\infty}{(-\kappa(q^n x);q^2)_\infty}}\;\Psi\left(\begin{array}{c} -q^2/\kappa(q^n p) \\ q^2\kappa(x/p)\end{array};\,q^2\kappa(q^n x/y)\right) \\
&\quad = \lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)}\,c_q\,s(x,y)\,(-1)^{\chi(p)}\,(-\operatorname{sgn}(y))^{\chi(x)}\,|x|\,q^m\,\nu(q^m p) \\
&\qquad \sqrt{\frac{(-\kappa(q^n p),-\operatorname{sgn}(p)\,q^{2m}\,\kappa(x);q^2)_\infty}{(-\kappa(q^n x);q^2)_\infty}}\;\Psi\left(\begin{array}{c} -q^2/\kappa(q^n p) \\ q^2\kappa(x/p)\end{array};\,\operatorname{sgn}(p)\,q^{2(1-m+n)}\right)\,.
\end{aligned} \tag{5.7}$$

If, on the other hand, $q^n x \notin I_q$ or $y/x \neq \operatorname{sgn}(p)\, q^m$, then

$$\begin{aligned}
\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n \digamma_{r,s,m,p}\big)(\lambda,x,\mu,y) &= \operatorname{sgn}(p)^n\,[(\tilde{v}^{-n}\otimes 1)\digamma_{r,s,m-n,pq^n}](\lambda,x,\mu,y) \\
&= \operatorname{sgn}(p)^n \operatorname{sgn}(x)^n\,\lambda^n\,\digamma_{r,s,m-n,pq^n}(\lambda,q^n x,\mu,y) = 0\,.
\end{aligned}$$

Using notation (5.4), we define the function $f_{r,s,m,p} : \mathbb{T} \times K_q \times \mathbb{T} \times I_q \rightarrow \mathbb{C}$ such that for $x \in K_q$, $y \in I_q$, $\lambda, \mu \in \mathbb{T}$, $f_{r,s,m,p}(\lambda,x,\mu,y)$ equals

$$\lambda^{r+\chi(y/p)}\,\mu^{s-\chi(x/p)}\,c_q\,s(x,y)\,(-1)^{\chi(p)}\,(-\operatorname{sgn}(y))^{\chi(x)}\,|y|\,\nu(q^m p)\,\sqrt{(-\kappa(y);q^2)_\infty}\;\phi(q^2\kappa(x/p);-q^{2(2-m)}/p^2) \tag{5.8}$$

if $y/x = \operatorname{sgn}(p)\, q^m$ and $f_{r,s,m,p}(\lambda,x,\mu,y) = 0$ if $y/x \neq \operatorname{sgn}(p)\, q^m$. Then it is clear from the above computations (and the proof of Lemma 6.2) that $\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n \digamma_{r,s,m,p}\big)_{n=1}^\infty$ converges pointwise to $f_{r,s,m,p}$. (5.9)

Define $J = \{\, x \in K_q \mid |x| \leq q^{1-m} \text{ or } \operatorname{sgn}(x) = \operatorname{sgn}(p)\,\}$. By Lemma 6.8 there exists a function $H \in l^2(J)^+$ such that

$$|\;|x|\,\sqrt{(-\operatorname{sgn}(p)\,q^{2m}\,\kappa(x);q^2)_\infty}\;\Psi(-q^2/\kappa(q^n p);q^2\kappa(x/p);\operatorname{sgn}(p)q^{2(1-m+n)})\;| \leq H(x)$$

for all $x \in J$ and $n \in \mathbb{N}$.

There exists clearly a number $C > 0$ such that $q^m\,\nu(q^m p)\,(-\kappa(q^n p);q^2)_\infty^{\frac{1}{2}}\,(q^2;q^2)_\infty^{-\frac{1}{2}} \leq C$ for all $n \in \mathbb{N}$. Let $(x,y) \in K_q \times I_q$ such that $y/x = \operatorname{sgn}(p)q^m$. If $|x| > q^{1-m}$ then $|q^m x| \geq 1$ and since $\operatorname{sgn}(p)q^m x = y \in I_q$, this implies that $\operatorname{sgn}(x) = \operatorname{sgn}(p)$. Hence, x belongs to J. So we can define an element $G \in \mathcal{L}^2(\mathbb{T} \times K_q \times \mathbb{T} \times I_q)$ such that for $\lambda, \mu \in \mathbb{T}$ and $x \in K_q$, $y \in I_q$, we have that $G(\lambda,x,\mu,y) = 0$ if $y/x \neq \operatorname{sgn}(p)q^m$ and $G(\lambda,x,\mu,y) = C\,H(x)$ if $y/x = \operatorname{sgn}(p)q^m$. Then Eq. (5.7) and the remarks thereafter imply that $|(\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n \digamma_{r,s,m,p}| \leq G$ for all $n \in \mathbb{N}$ such that $n \geq 1-\chi(p)$. Therefore the convergence in (5.9) and the dominated convergence theorem imply that $f_{r,s,m,p}$ belongs to $\tilde{H} \otimes H$ and that the sequence $\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n \digamma_{r,s,m,p}\big)_{n=1}^\infty$ converges to $f_{r,s,m,p}$ in $\tilde{H} \otimes H$.

Since the sequence $\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n\big)_{n=1}^\infty$ is a sequence of isometries and the linear span of such elements $\digamma_{r,s,m,p}$ is dense in $\tilde{H} \otimes H$, we conclude that there exists an isometry $\tilde{\Theta} : \tilde{H} \otimes H \rightarrow \tilde{H} \otimes H$ such that the sequence $\big((\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v})^n\big)_{n=1}^\infty$ converges strongly to $\tilde{\Theta}$. (5.10)
Thus, $\tilde{\Theta}\digamma_{r,s,m,p} = f_{r,s,m,p}$ for all $r,s,m \in \mathbb{Z}$ and $p \in K_q$. Using Eq. (5.8) and the definition of $\tilde{\Delta}$ and $\tilde{u}$, one easily checks that $(\tilde{u} \otimes e)f_{r,s,m,p} = f_{r,s,m,-p}$ and $\tilde{\Delta}(\tilde{u})\digamma_{r,s,m,p} = \digamma_{r,s,m,-p}$ for all $r,s,m \in \mathbb{Z}$ and

$p \in K_q$ (note that $s(-x,y) = \operatorname{sgn}(y)\, s(x,y)$ for all $x, y \in \mathbb{R} \setminus \{0\}$).
As a consequence, $\tilde{\Theta}\tilde{\Delta}(\tilde{u}) = (\tilde{u} \otimes e)\tilde{\Theta}$. (5.11)

Define $\hat{\Theta} \in B(\tilde{H} \otimes H)$ such that $\hat{\Theta}\restriction_{H\otimes H} = \tilde{\Theta}\restriction_{H\otimes H}$ and $\hat{\Theta}\restriction_{H^\perp \otimes H} = 0$. One sees that $\Delta(v)F_{r,s,m,p} = \operatorname{sgn}(p)F_{r,s,m-1,qp} = \tilde{\Delta}(\tilde{v})F_{r,s,m,p}$ for all $r, s, m \in \mathbb{Z}$, $p \in I_q$, implying that $\Delta(v)$ is the restriction of $\tilde{\Delta}(\tilde{v})$ to $H \otimes H$. So we get for every $n \in \mathbb{N}$ that $\tilde{\Delta}(\hat{v})^n\restriction_{H\otimes H} = \big(\Delta(v)\hat{\ }\big)^n\restriction_{H\otimes H} = \tilde{\Delta}(\tilde{v})^n\restriction_{H\otimes H}$, whereas $\tilde{\Delta}(\hat{v})^n\restriction_{H^\perp\otimes H} = 0$. It follows that $\hat{\Theta}$ is the strong limit of the sequence of isometries $\big(\,(\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\hat{v})^n\big)_{n=1}^{\infty}$. As such, $\hat{\Theta}$ belongs to $B(\tilde{H}) \otimes M_q$.
We set $P = u^*u$, which belongs to $M_q^\diamond$ because it is a multiplication operator. Multiplying Eq. (5.11) from the right by $\tilde{\Delta}(\hat{P})$ and using the fact that $\tilde{u}\,\hat{P} = \hat{u}$, we see that $\tilde{\Theta}\tilde{\Delta}(\hat{u}) = (\tilde{u} \otimes e)\tilde{\Theta}\tilde{\Delta}(\hat{P})$. Thus, $\hat{\Theta}\Delta(u)\hat{\ } = (\tilde{u} \otimes e)\hat{\Theta}\Delta(P)\hat{\ }$. (5.12)
Restricting the previous equation to $H \otimes H$, we get $\hat{\Theta}\Delta(u) = (\tilde{u} \otimes e)\hat{\Theta}\Delta(P)$. Applying $\iota \otimes \Delta$ to this equality, it follows that $(\iota \otimes \Delta)(\hat{\Theta})\,(\iota \otimes \Delta)(\Delta(u)) = (\tilde{u} \otimes e \otimes e)(\iota \otimes \Delta)(\hat{\Theta})(\iota \otimes \Delta)\Delta(P)$. (5.13)
If we apply $\tilde{\Delta}\otimes\iota$ to Eq. (5.12), we get $(\tilde{\Delta}\otimes\iota)(\hat{\Theta})(\Delta\otimes\iota)(\Delta(u))\hat{\ } = (\tilde{\Delta}(\tilde{u})\otimes e)(\tilde{\Delta}\otimes\iota)(\hat{\Theta})(\Delta\otimes\iota)(\Delta(P))\hat{\ }$. So the restriction of this equation to $H \otimes H \otimes H$ gives $(\tilde{\Delta}\otimes\iota)(\hat{\Theta})(\Delta\otimes\iota)\Delta(u) = (\tilde{\Delta}(\tilde{u})\otimes e)(\tilde{\Delta}\otimes\iota)(\hat{\Theta})(\Delta\otimes\iota)\Delta(P)$. Multiplying this equation from the left by $\tilde{\Theta} \otimes 1$, we see that

$$\begin{aligned}(\tilde{\Theta} \otimes 1)\,(\tilde{\Delta} \otimes \iota)(\hat{\Theta})(\Delta \otimes \iota)\Delta(u) &= (\tilde{\Theta}\tilde{\Delta}(\tilde{u}) \otimes e)(\tilde{\Delta} \otimes \iota)(\hat{\Theta})(\Delta \otimes \iota)\Delta(P) \\ &= (\tilde{u} \otimes e \otimes e)(\tilde{\Theta} \otimes 1)(\tilde{\Delta} \otimes \iota)(\hat{\Theta})(\iota \otimes \Delta)\Delta(P) \qquad (5.14)\end{aligned}$$

where in the last equality we used Eq. (5.11), the fact that $P \in M_q^\diamond$ and Corollary 5.3.
Now we calculate $(\tilde{\Theta} \otimes 1)\,(\tilde{\Delta} \otimes \iota)(\hat{\Theta})$. Let $n \in \mathbb{N}$. Since v belongs to $M_q^\diamond$, Corollary 5.3 guarantees that $(\Delta \otimes \iota)\Delta(v) = (\iota \otimes \Delta)\Delta(v)$ which implies that $(\tilde{\Delta} \otimes \iota)\tilde{\Delta}(\hat{v}) = (\iota \otimes \Delta)\tilde{\Delta}(\hat{v})$. Hence,

$$\begin{aligned}(\,[(\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\tilde{v}^n)] \otimes 1)(\tilde{\Delta} \otimes \iota)\big(\,(\tilde{v}^{-n} \otimes 1)\tilde{\Delta}(\hat{v}^n)\,\big) &= (\tilde{v}^{-n} \otimes 1 \otimes 1)\,(\tilde{\Delta} \otimes \iota)\tilde{\Delta}(\hat{v}^n) \\ &= (\tilde{v}^{-n} \otimes 1 \otimes 1)\,(\iota \otimes \Delta)\tilde{\Delta}(\hat{v}^n)\ .\end{aligned}$$

If we let n tend to infinity in this equality, the expression we started with converges strongly to $(\tilde{\Theta} \otimes 1)(\tilde{\Delta} \otimes \iota)(\hat{\Theta})$ whereas the expression we end up converges strongly to $(\iota \otimes \Delta)(\hat{\Theta})$. Thus, $(\tilde{\Theta} \otimes 1)(\tilde{\Delta} \otimes \iota)(\hat{\Theta}) = (\iota \otimes \Delta)(\hat{\Theta})$. Inserting this in Eq. (5.14), we conclude that $(\iota \otimes \Delta)(\hat{\Theta})\,(\Delta \otimes \iota)\Delta(u) = (\tilde{u} \otimes e \otimes e)(\iota \otimes \Delta)(\hat{\Theta})\,(\iota \otimes \Delta)\Delta(P)$. Comparing this with Eq. (5.13), we conclude that

$$(\iota \otimes \Delta)(\hat{\Theta})\,(\iota \otimes \Delta)\Delta(u) = (\iota \otimes \Delta)(\hat{\Theta})\,(\Delta \otimes \iota)\Delta(u)\ . \qquad (5.15)$$

The initial projection of $\hat{\Theta}$ is given by $X\otimes 1$ where $X \in B(\tilde{H})$ is the orthogonal projection onto H. In other words, $\hat{\Theta}^*\hat{\Theta} = X\otimes 1$. Thus, $(\iota\otimes\Delta)(\hat{\Theta})^*(\iota\otimes\Delta)(\hat{\Theta}) = X\otimes 1\otimes 1$, implying that $(\iota\otimes\Delta)(\hat{\Theta})$ is a partial isometry with initial space $H \otimes H \otimes H$. Therefore Eq. (5.15) guarantees that $(\iota \otimes \Delta)\Delta(u) = (\Delta \otimes \iota)\Delta(u)$. ☐

It is clear from the end of the proof above that we need $\hat{\Theta}$ to be an isometry on $H \otimes H \otimes H$. This is precisely the reason for using the extended Hilbert space $\tilde{H}$, a definition of $\hat{\Theta}$ as the strong limit of $\big(\,((v^*)^n \otimes 1)\Delta(v)^n\,\big)_{n=1}^{\infty}$ would not result in an isometry. The need for using $\tilde{\Theta}$ should be clear from the fact that the range of the operator $(\tilde{\Delta} \otimes \iota)(\hat{\Theta})(\Delta \otimes \iota)\Delta(u)$ in Eq. (5.14) is not contained in $H \otimes H \otimes H$.

In the last part of this section we provide the proof of Proposition 5.2. The most intricate part of this proof is contained in Proposition 5.8. The results before this proposition reduce the problem to a simpler form. We stress that this exposition does not depend on Corollary 5.3 or Proposition 5.4.

In order to filter out some distracting features we will change our Hilbert space $H^{\otimes 3}$. For this purpose, define a measure Ω on $I_{q^2}^3$ such that

$$\Omega(\{(x,z,y)\}) = |x|\,|y|\,\frac{(-x,-y;q^2)_\infty}{(-z;q^2)_\infty}$$

for all $x, y, z \in I_{q^2}$. In this section, all L^2-spaces that involve $I_{q^2}^3$ are defined by this measure.

Define the unitary transformation $U : H^{\otimes 3} \rightarrow L^2(\mathbb{T}^3) \otimes L^2(I_{q^2}^3, \Omega)$ such that $U\,[f] = [g]$, where the functions $f \in \mathcal{L}^2((\mathbb{T} \times I_q)^3)$ and $g \in \mathcal{L}^2(\mathbb{T}^3 \times I_{q^2}^3)$ are such that

$$\begin{aligned} g(\lambda, \mu, \eta, x, z, y) \quad = \quad & s(xz, y)\, s(z, x)\, \lambda^{-\log_{q^2}|x|}\, \mu^{\log_{q^2}|x/y|}\, \eta^{\log_{q^2}|y|}\, (-1)^{\log_{q^2}|z|}\, \operatorname{sgn}(z)^{\log_{q^2}|x|}\, \operatorname{sgn}(y)^{\log_{q^2}|xz|} \\ & |x|^{-1/2}\, |y|^{-1/2} \sqrt{\frac{(-z; q^2)_\infty}{(-x, -y; q^2)_\infty}}\; f(\lambda, \kappa^{-1}(x), \mu, \kappa^{-1}(z), \eta, \kappa^{-1}(y)) \end{aligned}$$

for all $\lambda, \mu, \eta \in \mathbb{T}$ and $x, z, y \in I_{q^2}$.

If $\theta \in -q^{2\mathbb{Z}} \cup q^{2\mathbb{Z}}$, we define the sets (see Fig. 1 and Fig. 2 in the proof of proposition 5.8)

$$K(\theta) = \{\, (x, y) \mid x, y \in I_{q^2} \text{ such that } \theta x y \in I_{q^2} \,\} \quad \text{ and } \quad L(\theta) = \{\, (x, \theta x y, y) \mid (x, y) \in K(\theta) \,\} \; .$$

Consider $m, k \in \mathbb{Z}$ and $p \in I_{q^2}$. Define the function $F^k_{m,p} \in \mathcal{F}(K(\operatorname{sgn}(p) q^{2k}))$ such that for $(x, y) \in K(\operatorname{sgn}(p) q^{2k})$,

$$\begin{aligned} F^k_{m,p}(x, y) \quad = \quad & c_q^2\, q^{\frac{1}{2}(m+k+1)(m+k+2)}\, \nu(\kappa^{-1}(p) q^m)\, \sqrt{(-p; q^2)_\infty} \\ & \Psi\left(\begin{matrix} -q^2/p \\ q^{2(1-m)}\, y/|p| \end{matrix} \; ; \operatorname{sgn}(p)\, q^{2(1-m)} \right) \Psi\left(\begin{matrix} -q^2/x \\ \operatorname{sgn}(p)\, q^{2(1+k)} y \end{matrix} \; ; q^{2(1+m+k)} x \right) \end{aligned} \tag{5.16}$$

if $\operatorname{sgn}(p) q^{-2m} y \in I_{q^2}$ and $F^k_{m,p}(x, y) = 0$ if $\operatorname{sgn}(p) q^{-2m} y \notin I_{q^2}$. We also define the function $G^k_{m,p} \in \mathcal{F}(K(\operatorname{sgn}(p) q^{2k}))$ by $G^k_{m,p}(x, y) = F^k_{-m,p}(y, x)$ for all $(x, y) \in K(\operatorname{sgn}(p) q^{2k})$.

Next we define functions $\tilde{F}^k_{m,p}, \tilde{G}^k_{m,p} \in \mathcal{F}(I_{q^2}^3)$ both of which have support in $L(\operatorname{sgn}(p) q^{2k})$ and satisfy

$$\tilde{F}^k_{m,p}(x, \operatorname{sgn}(p) q^{2k} x y, y) = F^k_{m,p}(x, y) \qquad \text{and} \qquad \tilde{G}^k_{m,p}(x, \operatorname{sgn}(p) q^{2k} x y, y) = G^k_{m,p}(x, y)$$

for all $(x, y) \in K(\operatorname{sgn}(p) q^{2k})$.

Define the unitary element $\varsigma \in C(\mathbb{T}^3)$ such that $\varsigma(\lambda, \eta, \mu) = \lambda \mu \bar{\eta}$ for all $\lambda, \eta, \mu \in \mathbb{T}$.

Lemma 5.5. *The families* $\{\, \tilde{F}^k_{m,p} \mid p \in I_{q^2}, m, k \in \mathbb{Z} \,\}$ *and* $\{\, \tilde{G}^k_{m,p} \mid p \in I_{q^2}, m, k \in \mathbb{Z} \,\}$ *are both orthonormal bases of* $L^2(I_{q^2}^3, \Omega)$. *Moreover,*

1) *the space* $\langle\, v \otimes \tilde{F}^k_{m,p} \mid v \in L^2(\mathbb{T}^3), p \in I_{q^2}, m, k \in \mathbb{Z} \,\rangle$ *is a core for* $U\,(\Delta \otimes \iota)(\Delta(\gamma))\, U^*$,

2) *the space* $\langle\, v \otimes \tilde{G}^k_{m,p} \mid v \in L^2(\mathbb{T}^3), p \in I_{q^2}, m, k \in \mathbb{Z} \,\rangle$ *is a core for* $U\,(\iota \otimes \Delta)(\Delta(\gamma))\, U^*$,

3) *if* $v \in L^2(\mathbb{T}^3)$, $p \in I_{q^2}$ *and* $m, k \in \mathbb{Z}$, *then*

$$\begin{aligned} (U(\Delta \otimes \iota)(\Delta(\gamma)) U^*)\, (v \otimes \tilde{F}^k_{m,p}) \quad &= \quad |p|^{-\frac{1}{2}}\, M_\varsigma v \otimes \tilde{F}^k_{m+1,p} \\ (U(\iota \otimes \Delta)(\Delta(\gamma)) U^*)\, (v \otimes \tilde{G}^k_{m,p}) \quad &= \quad |p|^{-\frac{1}{2}}\, M_\varsigma v \otimes \tilde{G}^k_{m+1,p} \; . \end{aligned}$$

Proof. The proof of this fact is pretty straightforward. For $r, s, t, n, m \in \mathbb{Z}$ and $p \in I_q$, we define $F^{r,s,t,n}_{m,p}, G^{r,s,t,n}_{m,p} \in H^{\otimes 3}$ as

$$\begin{aligned} F^{r,s,t,n}_{m,p} \quad &= \quad (V^* \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V^*)(\zeta^r \otimes \zeta^s \otimes \zeta^t \otimes \zeta^n \otimes \zeta^m \otimes \delta_p) \\ G^{r,s,t,n}_{m,p} \quad &= \quad (1_H \otimes V^*) V_{13}^* (\zeta^r \otimes \zeta^s \otimes \zeta^t \otimes \zeta^n \otimes \zeta^m \otimes \delta_p) \; . \end{aligned}$$

From Proposoition 3.8, we get immediately that the families $(\, F^{r,s,t,n}_{m,p} \mid r, s, t, n, m \in \mathbb{Z}, p \in I_q \,)$ and $(\, G^{r,s,t,n}_{m,p} \mid r, s, t, n, m \in \mathbb{Z}, p \in I_q \,)$ are both orthonormal bases of $H^{\otimes 3}$. Moreover, Eqs. (2.1), (5.1) and (5.2) imply that

1) the space $\langle\, F^{r,s,t,n}_{m,p} \mid r, s, t, n, m \in \mathbb{Z}, p \in I_q \rangle$ is a core for the operator $(\Delta \otimes \iota)(\Delta(\gamma))$,

2) the space $\langle\, G^{r,s,t,n}_{m,p} \mid r, s, t, n, m \in \mathbb{Z}, p \in I_q \rangle$ is a core for the operator $(\iota \otimes \Delta)(\Delta(\gamma))$,

3) if $r, s, t, m, n \in \mathbb{Z}$ and $p \in I_q$, then

$$(\Delta \otimes \iota)(\Delta(\gamma))\, F^{r,s,t,n}_{m,p} = p^{-1}\, F^{r,s,t,n}_{m+1,p} \qquad \text{and} \qquad (\iota \otimes \Delta)(\Delta(\gamma))\, G^{r,s,t,n}_{m,p} = p^{-1}\, G^{r,s,t,n}_{m+1,p} \; . \tag{5.17}$$

By Definition 3.6, we get the following equalities (in which the infinite sums are L^2-convergent),

$$\begin{aligned}
F^{r,s,t,n}_{m,p} &= (V^* \otimes 1_H)(1_{L^2(\mathbb{T}^2)} \otimes V^*)(\zeta^r \otimes \zeta^s \otimes \zeta^t \otimes \zeta^n \otimes \zeta^m \otimes \delta_p) = (V^* \otimes 1_H)(\zeta^r \otimes \zeta^s \otimes \digamma_{t,n,m,p}) \\
&= \sum_{\substack{y \in I_q \\ \mathrm{sgn}(p)q^{-m}y \in I_q}} a_p(\mathrm{sgn}(p)q^{-m}y, y)\; (V^* \otimes 1_H)(\zeta^r \otimes \zeta^s \otimes \; \zeta^{t+\chi(y/p)} \otimes \\
&\qquad\qquad\qquad\qquad \delta_{\mathrm{sgn}(p)q^{-m}y} \otimes \zeta^{n-\chi(q^{-m}y/p)} \otimes \delta_y) \\
&= \sum_{\substack{y \in I_q \\ \mathrm{sgn}(p)q^{-m}y \in I_q}} a_p(\mathrm{sgn}(p)q^{-m}y, y)\; \digamma_{r,s,t+\chi(y/p),\mathrm{sgn}(p)q^{-m}y} \otimes \zeta^{n+m+\chi(p/y)} \otimes \delta_y \\
&= \sum_{\substack{y \in I_q \\ \mathrm{sgn}(p)q^{-m}y \in I_q}} \; \sum_{\substack{x \in I_q \\ \mathrm{sgn}(p)\mathrm{sgn}(y)q^{t+\chi(y/p)}x \in I_q}} a_{\mathrm{sgn}(p)q^{-m}y}(x, \mathrm{sgn}(p)\mathrm{sgn}(y)q^{t+\chi(y/p)}x) \\
&\qquad a_p(\mathrm{sgn}(p)q^{-m}y, y)\, \zeta^{r+t+\chi(y/p)+\chi(x)+m-\chi(y)} \otimes \delta_x \otimes \zeta^{s-\chi(x)-m+\chi(y)} \otimes \\
&\qquad \delta_{\mathrm{sgn}(p)\mathrm{sgn}(y)q^{t+\chi(y/p)}x} \otimes \zeta^{n+m+\chi(p/y)} \otimes \delta_y \\
&= \sum_{\substack{y \in I_q \\ \mathrm{sgn}(p)q^{-m}y \in I_q}} \; \sum_{\substack{x \in I_q \\ (q^t/p)\,xy \in I_q}} a_p(\mathrm{sgn}(p)q^{-m}y, y)\, a_{\mathrm{sgn}(p)q^{-m}y}(x, (q^t/p)\,xy) \\
&\qquad \zeta^{r+t+m+\chi(x/p)} \otimes \delta_x \otimes \zeta^{s-m+\chi(y/x)} \otimes \delta_{(q^t/p)\,xy} \otimes \zeta^{n+m+\chi(p/y)} \otimes \delta_y \;. \qquad (5.18)
\end{aligned}$$

Choose $\lambda, \mu, \eta \in \mathbb{T}$ and $x, z, y \in I_{q^2}$. Look first at the case where $\mathrm{sgn}(p)\, q^{-2m}\, y \in I_{q^2}$ and $z = \mathrm{sgn}(p)\, q^{2(t-\chi(p))}\, xy$. Set $x' = \kappa^{-1}(x)$, $y' = \kappa^{-1}(y)$ and $z' = \kappa^{-1}(z)$. Thus, $\mathrm{sgn}(p)q^{-m}\, y' \in I_q$ and $(q^t/p)x'y' = z' \in I_q$. Set $d = \lambda^{r+t+m+\chi(x'/p)}\, \mu^{s-m+\chi(y'/x')}\, \eta^{n+m+\chi(p/y')}$. The above equation, Proposition 3.5 and Definition 3.1 imply that

$$\begin{aligned}
&F^{r,s,t,n}_{m,p}(\lambda, \kappa^{-1}(x), \mu, \kappa^{-1}(z), \eta, \kappa^{-1}(y)) \;= d\; a_p(\mathrm{sgn}(p)q^{-m}y', y')\, a_{\mathrm{sgn}(p)q^{-m}y'}(x', z') \\
&\quad = d\; (\mathrm{sgn}(p)\mathrm{sgn}(y'))^{m+\chi(y')}\, \mathrm{sgn}(z')^{\chi(z')}\, \mathrm{sgn}(x')^{\chi(x')}\, a_p(\mathrm{sgn}(p)q^{-m}y', y')\, a_{\mathrm{sgn}(p)q^{-m}y'}(z', x') \\
&\quad = d\; (\mathrm{sgn}(p)\,\mathrm{sgn}(y'))^{m+\chi(y')}\, \mathrm{sgn}(z')^{\chi(x')}\, (\mathrm{sgn}(p)\,\mathrm{sgn}(x')\,\mathrm{sgn}(y'))^{\chi(p)+\chi(y')+t}\; \mathrm{sgn}(x')^{\chi(x')} \\
&\qquad c_q\, s(\mathrm{sgn}(p)q^{-m}y', y')\, (-1)^{\chi(p)+m+\chi(y')}\, \mathrm{sgn}(y')^{m+\chi(y')}\, |y'| \\
&\qquad \nu(pq^m) \sqrt{\frac{(-\kappa(p), -y; q^2)_\infty}{(-\mathrm{sgn}(p)q^{-2m}y; q^2)_\infty}}\; \Psi\left(\begin{matrix} -q^2/\kappa(p) \\ q^{2(1-m)}\, y/|\kappa(p)| \end{matrix} \;;\, \mathrm{sgn}(p)\, q^{2(1-m)} \right) \\
&\qquad c_q\, s(z', x')\, (-1)^{m+\chi(y')+\chi(z')}\, \mathrm{sgn}(x')^{\chi(z')}\, |x'| \\
&\qquad \nu(q^{-m}y'x'/z') \sqrt{\frac{(-\mathrm{sgn}(p)q^{-2m}y, -x; q^2)_\infty}{(-z; q^2)_\infty}}\; \Psi\left(\begin{matrix} -q^2/x \\ q^2 z/x \end{matrix} \;;\, \mathrm{sgn}(p)\, q^{2(1+m)} z/y \right) \\
&\quad = d\, c_q^2\, \nu(pq^m)\, \nu(q^{-m-t+\chi(p)})\, s(xz, y)\, s(z, x)\, (-1)^{\chi(p)+\chi(z')} \\
&\qquad \mathrm{sgn}(p)^{m+t+\chi(p)}\, \mathrm{sgn}(z')^{\chi(x')}\, \mathrm{sgn}(y')^{\chi(x'z')}\; |x|^{\frac{1}{2}}\, |y|^{\frac{1}{2}} \\
&\qquad \sqrt{\frac{(-\kappa(p), -x, -y; q^2)_\infty}{(-z; q^2)_\infty}}\;\; \Psi\left(\begin{matrix} -q^2/\kappa(p) \\ q^{2(1-m)}\, y/|\kappa(p)| \end{matrix} \;;\, \mathrm{sgn}(p)\, q^{2(1-m)} \right) \\
&\qquad \Psi\left(\begin{matrix} -q^2/x \\ \mathrm{sgn}(p)\, q^{2(1+t-\chi(p))} y \end{matrix} \;;\, q^{2(1+m+t-\chi(p))} x \right) \\
&\quad = \lambda^{\log_{q^2}|x|}\, \mu^{\log_{q^2}|y/x|}\, \eta^{-\log_{q^2}|y|}\;\; s(xz, y)\, s(z, x)\, (-1)^{\log_{q^2}|z|}\, \mathrm{sgn}(z)^{\log_{q^2}|x|}\, \mathrm{sgn}(y)^{\log_{q^2}|xz|}\, |x|^{\frac{1}{2}}\, |y|^{\frac{1}{2}} \\
&\qquad \sqrt{\frac{(-x, -y; q^2)_\infty}{(-z; q^2)_\infty}}\, (-1)^{\chi(p)}\, \mathrm{sgn}(p)^{m+t+\chi(p)}\;\; \lambda^{r+t+m-\chi(p)}\, \mu^{s-m}\, \eta^{n+m+\chi(p)}\;\; \tilde{F}^{t-\chi(p)}_{m,\kappa(p)}(x, z, y)\;.
\end{aligned}$$

If $\mathrm{sgn}(p)\, q^{-2m}\, y \notin I_{q^2}$ or $z \neq \mathrm{sgn}(p)\, q^{2(t-\chi(p))}\, xy$, then Eq. (5.18) implies immediately that $F^{r,s,t,n}_{m,p}(\lambda, \kappa^{-1}(x), \mu, \kappa^{-1}(z), \eta, \kappa^{-1}(y)) = 0$. So we see that

$$U F^{r,s,t,n}_{m,p} = (-1)^{\chi(p)}\, \mathrm{sgn}(p)^{m+t+\chi(p)}\;\; \zeta^{r+t+m-\chi(p)} \otimes \zeta^{s-m} \otimes \zeta^{n+m+\chi(p)}\;\; \otimes \tilde{F}^{t-\chi(p)}_{m,\kappa(p)}\;.$$

Proposition 3.10 implies that

$$\begin{aligned} UG^{r,s,t,n}_{m,p} &= U\Sigma_{13}(\breve{J}\otimes\breve{J}\otimes\breve{J})(V^*\otimes 1_H)(1_{L^2(\mathbb{T}^2)}\otimes V^*)(\tilde{\Sigma}\otimes\breve{J})(\zeta^r\otimes\zeta^s\otimes\zeta^t\otimes\zeta^n\otimes\zeta^m\otimes\delta_p) \\ &= \operatorname{sgn}(p)^{\chi(p)}\, U\Sigma_{13}(\breve{J}\otimes\breve{J}\otimes\breve{J})(V^*\otimes 1_H)(1_{L^2(\mathbb{T}^2)}\otimes V^*)(\zeta^{-n}\otimes\zeta^{-t}\otimes\zeta^{-s}\otimes\zeta^{-r}\otimes\zeta^{-m}\otimes\delta_p) \\ &= \operatorname{sgn}(p)^{\chi(p)}\, U\Sigma_{13}(\breve{J}\otimes\breve{J}\otimes\breve{J})F^{-n,-t,-s,-r}_{-m,p}\ . \end{aligned}$$

Define the anti-unitary operator $\breve{I}$ on $L^2(\mathbb{T}^3)\otimes L^2(I^3_{q^2},\Omega)$ such that $\breve{I}\,[f]=[g]$ where $f,g\in\mathcal{L}^2(\mathbb{T}^3\times I^3_{q^2})$ are such that $g(\lambda,\mu,\eta,x,z,y)=\operatorname{sgn}(xyz)^{\log_{q^2}|xyz|}\ \overline{f(\eta,\mu,\lambda,y,z,x)}$ for all $x,y,z\in I_{q^2}$ and $\lambda,\mu,\eta\in\mathbb{T}$.

Let $x,y,z\in I_{q^2}$. If $z>0$, it follows from the considerations before Definition 3.6 that $s(xz,y)=\operatorname{sgn}(xy)\,s(yz,x)$ and $s(z,x)=\operatorname{sgn}(xy)\,s(z,y)$. If on the other hand $z<0$, we have $s(xz,y)=s(yz,x)$ and $s(z,x)=s(z,y)$. Thus, in both cases $s(xz,y)\,s(z,x)=s(yz,x)\,s(z,y)$. Now it is straightforward to check that $U\Sigma_{13}(\breve{J}\otimes\breve{J}\otimes\breve{J})=\breve{I}U$. Thus,

$$\begin{aligned} UG^{r,s,t,n}_{m,p} &= \operatorname{sgn}(p)^{\chi(p)}\,\breve{I}UF^{-n,-t,-s,-r}_{-m,p} \\ &= (-1)^{\chi(p)}\operatorname{sgn}(p)^{m+s}\,\breve{I}\,(\zeta^{-n-s-m-\chi(p)}\otimes\zeta^{-t+m}\otimes\zeta^{-r-m+\chi(p)}\otimes\tilde{F}^{-s-\chi(p)}_{-m,\kappa(p)})\ . \end{aligned}$$

Since $\tilde{F}^{-s-\chi(p)}_{-m,\kappa(p)}$ is supported on the set $L(q^{-2s}/\kappa(p))$ and $\operatorname{sgn}(xyz)^{\log_{q^2}|xyz|}=\operatorname{sgn}(p)^{s+\chi(p)}$ for all $(x,y,z)\in L(q^{-2s}/\kappa(p))$, we get that

$$UG^{r,s,t,n}_{m,p} = (-1)^{\chi(p)}\operatorname{sgn}(p)^{m+\chi(p)}\,\zeta^{r+m-\chi(p)}\otimes\zeta^{t-m}\otimes\zeta^{n+s+m+\chi(p)}\otimes\tilde{G}^{-s-\chi(p)}_{m,\kappa(p)}\ .$$

This implies for $r',s',t'\in\mathbb{Z}$, $k,m\in\mathbb{Z}$ and $p'\in I_{q^2}$,

$$\begin{aligned} \zeta^{r'}\otimes\zeta^{s'}\otimes\zeta^{t'}\otimes\tilde{F}^k_{m,p'} &= (-1)^{\log_{q^2}|p'|}\operatorname{sgn}(p')^{m+k}\,UF^{r'-m-k,s'+m,k+\log_{q^2}|p'|,t'-m-\log_{q^2}|p'|}_{m,\kappa^{-1}(p')} \\ \zeta^{r'}\otimes\zeta^{s'}\otimes\zeta^{t'}\otimes\tilde{G}^k_{m,p'} &= (-1)^{\log_{q^2}|p'|}\operatorname{sgn}(p')^{m+\log_{q^2}|p'|}\,UF^{r'-m+\log_{q^2}|p'|,-k-\log_{q^2}|p'|,s'+m,t'-m+k}_{m,\kappa^{-1}(p')}\ . \end{aligned}$$

From these two equations and the properties listed in (5.17) in the beginning of the proof, all the claims in the statement of this lemma easily follow. ☐

Set $D=\langle\,\zeta^r\mid r\in\mathbb{Z}\,\rangle\subseteq L^2(\mathbb{T})$.

Lemma 5.6. *The operator* $U\Delta^{(2)}_0(\gamma_0)U^*$ *is an adjointable operator on* $D^{\odot 3}\odot\mathcal{K}(I_{q^2})^{\odot 3}$ *such that* $(U\Delta^{(2)}_0(\gamma_0)U^*)^\dagger=U\Delta^{(2)}_0(\gamma_0^\dagger)U^*$. *Moreover*

$$\begin{aligned} (U\Delta^{(2)}_0(\gamma_0)U^*\,g)(\lambda,\mu,\eta,x,z,y) = q\,|xy/z|^{\frac{1}{2}}\,\frac{\lambda\bar{\mu}\eta}{xy}\,\big[\,&(1+q^{-2}y)\ f(\lambda,\mu,\eta,q^2x,z,q^{-2}y) \\ &-(1+q^{-2}y)\ g(\lambda,\mu,\eta,x,q^{-2}z,\eta,q^{-2}y)-(1+z)\ g(\lambda,\mu,\eta,q^2x,q^2z,y)+g(\lambda,\mu,\eta,x,z,y)\,\big] \end{aligned}$$

for $g\in D^{\odot 3}\odot\mathcal{K}(I_{q^2})^{\odot 3}$, $\lambda,\mu,\eta\in\mathbb{T}$ *and* $x,z,y\in I_{q^2}$.

Proof. Since the domain of $\Delta^{(2)}_0(\gamma_0)$ and $\Delta^{(2)}_0(\gamma_0^\dagger)$ is $E^{\odot 3}$ and clearly $U(E^{\odot 3})=D^{\odot 3}\odot\mathcal{K}(I_{q^2})^{\odot 3}$, the statements about the adjointability of $\Delta^{(2)}_0(\gamma_0)$ follow easily. Using Eqs. (1.5), one checks that

$$\Delta^{(2)}_0(\gamma_0)=\gamma_0\odot\alpha_0\odot\alpha_0+e_0\alpha_0^\dagger\odot\gamma_0\odot\alpha_0+e_0\alpha_0^\dagger\odot e_0\alpha_0^\dagger\odot\gamma_0+q\,\gamma_0\odot e_0\gamma_0^\dagger\odot\gamma_0\ .$$

So we get for $f\in E^{\odot 3}$, $x,y,z\in I_q$ and $\lambda,\mu,\eta\in\mathbb{T}$ that

$$\begin{aligned} &(\Delta^{(2)}_0(\gamma_0)\,f)(\lambda,x,\mu,z,\eta,y) \\ &\quad=\frac{\lambda}{x}\,\sqrt{\operatorname{sgn}(z)+q^2/z^2}\,\sqrt{\operatorname{sgn}(y)+q^2/y^2}\ f(\lambda,x,\mu,q^{-1}z,\eta,q^{-1}y) \\ &\qquad+\operatorname{sgn}(x)\,\frac{\mu}{z}\,\sqrt{\operatorname{sgn}(x)+1/x^2}\,\sqrt{\operatorname{sgn}(y)+q^2/y^2}\ f(\lambda,qx,\mu,z,\eta,q^{-1}y) \\ &\qquad+\operatorname{sgn}(xz)\frac{\eta}{y}\,\sqrt{\operatorname{sgn}(x)+1/x^2}\,\sqrt{\operatorname{sgn}(z)+1/z^2}\ f(\lambda,qx,\mu,qz,\eta,y)+\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{xyz}\,f(\lambda,x,\mu,z,\eta,y) \\ &\quad=\operatorname{sgn}(yz)\,\frac{q^2\lambda}{xyz}\,\sqrt{1+q^{-2}\,\kappa(z)}\,\sqrt{1+q^{-2}\kappa(y)}\ f(\lambda,x,\mu,q^{-1}z,\eta,q^{-1}y) \\ &\qquad+\operatorname{sgn}(y)\,\frac{q\mu}{xyz}\,\sqrt{1+\kappa(x)}\,\sqrt{1+q^{-2}\kappa(y)}\ f(\lambda,qx,\mu,z,\eta,q^{-1}y) \\ &\qquad+\frac{\eta}{xyz}\,\sqrt{1+\kappa(x)}\,\sqrt{1+\kappa(z)}\ f(\lambda,qx,\mu,qz,\eta,y)+\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{xyz}\,f(\lambda,x,\mu,z,\eta,y)\ . \end{aligned}$$

The inverse U^* of U is such that

$$\begin{aligned}(U^*f)(\lambda,x,\mu,z,\eta,y) \;&=\; s(xz,y)\, s(z,x)\, \lambda^{\log_q|x|}\, \mu^{\log_q|y/x|}\, \eta^{-\log_q|y|}\, (-1)^{\log_q|z|}\, \operatorname{sgn}(z)^{\log_q|x|}\, \operatorname{sgn}(y)^{\log_q|xz|} \\ &\qquad |x|\,|y| \sqrt{\frac{(-\kappa(x),-\kappa(y);q^2)_\infty}{(-\kappa(z);q^2)_\infty}}\; f(\lambda,\mu,\eta,\kappa(x),\kappa(z),\kappa(y))\end{aligned}$$

for $f \in D^{\odot 3} \odot \mathcal{K}(I_q)^{\odot 3}$, $\lambda,\mu,\eta \in \mathbb{T}$ and $x,z,y \in I_q$.
Then the following rules are easily established for $g \in D^{\odot 3} \odot \mathcal{K}(I_{q^2})^{\odot 3}$, $x,y,z \in I_{q^2}$ and $\omega = (\lambda,\eta,\mu) \in \mathbb{T}^3$.

(1) If $f \in D^{\odot 3} \odot \mathcal{K}(I_q)^{\odot 3}$, then $(UM_fU^*g)(\omega,x,z,y) = f(\lambda,\kappa^{-1}(x),\mu,\kappa^{-1}(z),\eta,\kappa^{-1}(y))\; g(\omega,x,z,y)$.

(2) $(U(1 \odot T_{q^{-1}} \odot T_{q^{-1}})U^*g)(\omega,x,z,y) = -q^{-1}\operatorname{sgn}(y)\,\bar{\mu}\eta\,\sqrt{(1+q^{-2}y)(1+q^{-2}z)^{-1}}\, g(\omega,x,q^{-2}z,q^{-2}y)$.

(3) $(U(T_q \odot 1 \odot T_{q^{-1}})U^*g)(\omega,x,z,y) = \operatorname{sgn}(yz)\,\lambda\bar{\mu}^2\eta\,\sqrt{(1+q^{-2}y)(1+x)^{-1}}\, g(\omega,q^2x,z,q^{-2}y)$.

(4) $(U(T_q \odot T_q \odot 1)U^*g)(\omega,x,z,y) = -q\operatorname{sgn}(z)\,\lambda\bar{\mu}\,\sqrt{(1+z)(1+x)^{-1}}\, g(\omega,q^2x,q^2z,y)$.

This implies that for $g \in D^{\odot 3} \odot \mathcal{K}(I_{q^2})^{\odot 3}$, $x,y,z \in I_{q^2}$ and $\omega = (\lambda,\eta,\mu) \in \mathbb{T}^3$,

$$\begin{aligned}&(U\Delta_0^{(2)}(\gamma_0)U^*\, g)(\omega,x,z,y) \\ &\quad = -\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{\kappa^{-1}(xyz)}\,(1+q^{-2}y)\; g(\omega,x,q^{-2}z,q^{-2}y) \\ &\qquad +\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{\kappa^{-1}(xyz)}\,(1+q^{-2}y)\; g(\omega,q^2x,z,q^{-2}y) \\ &\qquad -\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{\kappa^{-1}(xyz)}\,(1+z)\; g(\omega,q^2x,q^2z,y) \\ &\qquad +\operatorname{sgn}(z)\,\frac{q\lambda\bar{\mu}\eta}{\kappa^{-1}(xyz)}\, g(\omega,x,z,y)\;. \\ &\quad = q\,|xy/z|^{\frac{1}{2}}\,\frac{\lambda\bar{\mu}\eta}{xy}\,\big(\,(1+q^{-2}y)\; g(\omega,q^2x,z,q^{-2}y) - (1+q^{-2}y)\; g(\omega,x,q^{-2}z,q^{-2}y) \\ &\qquad -(1+z)\; g(\omega,q^2x,q^2z,y) + g(\omega,x,z,y)\,\big)\;.\end{aligned}$$

☐

Let us also get rid of part of the operators acting on $L^2(\mathbb{T}^3)$. For this purpose we introduce normal operators γ_l on γ_r in $L^2(I_q^3,\Omega)$ such that

(1) the space $\langle\, \tilde{F}^k_{m,p} \mid p \in I_{q^2}, m,k \in \mathbb{Z}\rangle$ is a core for γ_l,

(2) the space $\langle\, \tilde{G}^k_{m,p} \mid p \in I_{q^2}, m,k \in \mathbb{Z}\rangle$ is a core for γ_r,

(3) if $p \in I_{q^2}$ and $m,k \in \mathbb{Z}$, then $\gamma_l\,\tilde{F}^k_{m,p} = |p|^{-\frac{1}{2}}\,\tilde{F}^k_{m+1,p}$ and $\gamma_r\,\tilde{G}^k_{m,p} = |p|^{-\frac{1}{2}}\,\tilde{G}^k_{m+1,p}$.

By Lemma 5.5 we know that $U((\Delta\otimes\iota)\Delta(\gamma))U^* = M_\varsigma \otimes \gamma_l$ and $U((\iota\otimes\Delta)\Delta(\gamma))U^* = M_\varsigma \otimes \gamma_r$. So in order to prove that $(\Delta\otimes\iota)\Delta(\gamma) = (\iota\otimes\Delta)\Delta(\gamma)$, it is enough to show that $\gamma_l = \gamma_r$.

The description of γ_l and γ_r in terms of eigenvectors will not be sufficient to solve this problem. We will also need to know the explicit action of γ_l and γ_r on functions in its domain. For this purpose we introduce the linear operators $\tilde{\gamma}$, $\tilde{\gamma}^\dagger$ belonging to $\operatorname{End}(\mathcal{F}(I^3_{q^2}))$ such that

$$\begin{aligned}(\tilde{\gamma}\, f)(x,z,y) \;&=\; q\,|xy/z|^{\frac{1}{2}}\,\frac{1}{xy}\,\big(\,(1+q^{-2}y)\; f(q^2x,z,q^{-2}y) - (1+q^{-2}y)\; f(x,q^{-2}z,q^{-2}y) \\ &\qquad\qquad -(1+z)\; f(q^2x,q^2z,y) + f(x,z,y)\,\big)\end{aligned}$$

and

$$\begin{aligned}(\tilde{\gamma}^\dagger\, f)(x,z,y) \;&=\; q\,|xy/z|^{\frac{1}{2}}\,\frac{1}{xy}\,\big(\,(1+q^{-2}x)\; f(q^{-2}x,z,q^2y) - (1+z)\; f(x,q^2z,q^2y) \\ &\qquad\qquad -(1+q^{-2}x)\; f(q^{-2}x,q^{-2}z,y) + f(x,z,y)\,\big)\end{aligned}$$

for all $f \in \mathcal{F}(I^3_{q^2})$ and $x,y,z \in I_{q^2}$. Note that $\tilde{\gamma}^\dagger$ is obtained from $\tilde{\gamma}$ by interchanging x and y.

Lemma 5.7. *If $f \in D(\gamma_l)$ and $g \in D(\gamma_r^*)$, then $\gamma_l(f) = \tilde{\gamma}(f)$ and $\gamma_r^*(g) = \tilde{\gamma}^\dagger(g)$.*

Proof. Define the sesquilinear form $\langle\, .\, ,\, .\,\rangle : \mathcal{K}(I^3_{q^2}) \times \mathcal{F}(I^3_{q^2}) \to \mathbb{C}$ such that

$$\langle f, g\rangle = \sum_{(x,z,y)\in I^3_{q^2}} f(x,z,y)\, \bar{g}(x,z,y)\, |x|\, |y|\, \frac{(-x,-y;q^2)_\infty}{(-z;q^2)_\infty}$$

for all $f \in \mathcal{K}(I^3_{q^2})$ and $g \in \mathcal{F}(I^3_{q^2})$.

Under this pairing, $\mathcal{F}(I^3_{q^2})$ is anti-linearly isomorphic to the algebraic dual of $\mathcal{K}(I^3_{q^2})$. So there exist an anti-linear anti-homomorphism $^\circ : \mathrm{End}(\mathcal{K}(I^3_{q^2})) \to \mathrm{End}(\mathcal{F}(I^3_{q^2}))$ such that $\langle T(f), g\rangle = \langle f, T^\circ(g)\rangle$ for all $f \in \mathcal{K}(I^3_{q^2})$ and $g \in \mathcal{F}(I^3_{q^2})$. If $f \in \mathcal{K}(I^3_{q^2})$ and $g \in L^2(I^3_{q^2}, \Omega)$ then $\langle f, g\rangle$ equals the inner product in $L^2(I^3_{q^2}, \Omega)$. So if $T \in \mathrm{End}(\mathcal{K}(I^3_{q^2}))$, then $T^*(f) = T^\circ(f)$ for all $f \in D(T^*)$ (and the domain of T^* consists of all $f \in L^2(I^3_{q^2}, \Omega)$ such that $T^\circ(f) \in L^2(I^3_{q^2}, \Omega)$). Moreover, $T \in \mathrm{End}(\mathcal{K}(I^3_{q^2}))$ is adjointable if and only if $T^\circ(\mathcal{K}(I^3_{q^2})) \subseteq \mathcal{K}(I^3_{q^2})$ in which case $T^\dagger$ is the restriction of T° to $\mathcal{K}(I^3_{q^2})$.

If $p = (p_1, p_2, p_3) \in (-q^{2\mathbb{Z}} \cup q^{2\mathbb{Z}})^3$ and $g \in \mathcal{F}(I^3_{q^2})$, we define $\tilde{T}_p, \tilde{M}_g \in \mathrm{End}(\mathcal{F}(I^3_{q^2}))$ such that

$$\tilde{T}_p(f)(x,z,y) = f(p_1 x, p_2 z, p_3 y) \qquad\qquad \text{and} \qquad\qquad \tilde{M}_g(f)(x,z,y) = g(x,z,y)\, f(x,z,y)$$

for all $f \in \mathcal{F}(I^3_{q^2})$ and $x, z, y \in I_{q^2}$. We denote the restrictions of $\tilde{M}_g, \tilde{T}_p$ to $\mathcal{K}(I^3_{q^2})$ by $\hat{M}_g, \hat{T}_p \in \mathrm{End}(\mathcal{K}(I^3_{q^2}))$ respectively.

Let ξ_1, ξ_2, ξ_3 denote the 3 coordinate functions on $I^3_{q^2}$. Define also an auxiliary function $\omega : I^3_{q^2} \to \mathbb{R}$ such that for $x, y, z \in I_{q^2}$, $\omega(x,z,y) = (1 + q^{-2} z)^{-1}$ if $z \neq -q^2$ and $\omega(x,z,y) = 0$ if $z = -q^2$. It is not so difficult to check that

$$\begin{aligned}
&\hat{T}^\circ_{q^2,1,1} = q^{-2}\, \tilde{M}_{1+q^{-2}\xi_1}\, \tilde{T}_{q^{-2},1,1} &\qquad& \hat{T}^\circ_{1,1,q^{-2}} = q^2\, \tilde{M}_{(1+\xi_3)^{-1}}\, \tilde{T}_{1,1,q^2}\\
&\hat{T}^\circ_{1,q^2,1} = \tilde{M}_\omega\, \tilde{T}_{1,q^{-2},1} && \hat{T}^\circ_{1,q^{-2},1} = \tilde{M}_{1+\xi_2}\, \tilde{T}_{1,q^2,1}\\
&\hat{M}^\circ_g = \tilde{M}_{\bar{g}} \ \text{ if } g \in \mathcal{F}(I^3_{q^2})\ .
\end{aligned}$$

Thus

$$\begin{aligned}
&\hat{T}^\circ_{q^2,q^2,1} = q^{-2}\, \tilde{M}_{\omega(1+q^{-2}\xi_1)}\, \tilde{T}_{q^{-2},q^{-2},1} &\qquad& \hat{T}^\circ_{1,q^{-2},q^{-2}} = q^2\, \tilde{M}_{(1+\xi_2)(1+\xi_3)^{-1}}\, \tilde{T}_{1,q^2,q^2}\\
&\hat{T}^\circ_{q^2,1,q^{-2}} = \tilde{M}_{(1+q^{-2}\xi_1)(1+\xi_3)^{-1}}\, \tilde{T}_{q^{-2},1,q^2}\ .
\end{aligned}$$

Let $\hat{\gamma}$ denote the restriction of $\tilde{\gamma}$ to $\mathcal{K}(I^3_{q^2})$. Clearly, $\hat{\gamma} = \sum_p \hat{T}_p\, \hat{M}_{g_p}$ where p ranges over $(q^2, q^2, 1)$, $(1, q^{-2}, q^{-2})$, $(q^2, 1, q^{-2})$, $(1,1,1)$ and g_p belongs to $\mathcal{F}(I^3_q)$. Therefore $\hat{\gamma}$ is adjointable.
Moreover, $\hat{\gamma}^\circ = \sum_p \hat{M}_{g_p}\, \hat{T}^\circ_p$. Now an easy calculation reveals that $\hat{\gamma}^\circ = \tilde{\gamma}^\dagger$. Similarly, $(\hat{\gamma}^\dagger)^\circ = \tilde{\gamma}$.

Lemma 5.6 implies that $U\Delta^{(2)}_0(\gamma_0)U^* = M_\varsigma\restriction_{D^{\odot 3}} \odot \hat{\gamma}$ and $U\Delta^{(2)}_0(\gamma^\dagger)U^* = M^*_\varsigma\restriction_{D^{\odot 3}} \odot \hat{\gamma}^\dagger$.
By Proposition 5.1 and the remarks before this lemma we get that

$$M^*_\varsigma\restriction_{D^{\odot 3}} \odot \hat{\gamma}^\dagger = U\Delta^{(2)}_0(\gamma_0^\dagger)U^* \subseteq U((\Delta \otimes \iota)\Delta(\gamma^*))U^* = M^*_\varsigma \otimes \gamma^*_l\ .$$

Multiplying this inclusion from the right with $M_\varsigma \otimes 1_{L^2(I^3_{q^2},\Omega)}$, we get that $1_{D^{\odot 3}} \odot \hat{\gamma}^\dagger \subseteq 1_{L^2(\mathbb{T}^3)} \otimes \gamma^*_l$.
Fix a unit vector $w \in D^{\odot 3}$. Define the bounded linear operator $T : L^2(\mathbb{T}^3) \to \mathbb{C}$ such that $T(x) = \langle x, w\rangle$ for all $x \in L^2(\mathbb{T}^3)$. Then $(T \otimes 1_{L^2(I^3_{q^2},\Omega)})\,(1_{L^2(\mathbb{T}^3)} \otimes \gamma^*_l) \subseteq \gamma^*_l\,(T \otimes 1_{L^2(I^3_{q^2},\Omega)})$. So if $v \in \mathcal{K}(I^3_{q^2})$, then $w \otimes v \in D(1_{L^2(\mathbb{T}^3)} \otimes \gamma^*_l)$ and $(1_{L^2(\mathbb{T}^3)} \otimes \gamma^*_l)(w \otimes v) = w \otimes \hat{\gamma}^\dagger\, v$. Thus, $v = (T \otimes 1_{L^2(I^3_{q^2},\Omega)})(w \otimes v) \in D(\gamma^*_l)$ and

$$\gamma^*_l v = \gamma^*_l (T \otimes 1_{L^2(I^3_{q^2},\Omega)})(w \otimes v) = (T \otimes 1_{L^2(I^3_{q^2},\Omega)})(1_{L^2(\mathbb{T}^3)} \otimes \gamma^*_l)(w \otimes v) = (T \otimes 1_{L^2(I^3_{q^2},\Omega)})(w \otimes \hat{\gamma}^\dagger v) = \hat{\gamma}^\dagger v\ .$$

So we have shown that $\hat{\gamma}^\dagger \subseteq \gamma^*_l$. Taking the adjoint of this inclusion we see that $\gamma_l \subseteq (\hat{\gamma}^\dagger)^*$. By the remarks in the beginning of the proof, this implies for $f \in D(\gamma_l)$, $\gamma_l(f) = (\hat{\gamma}^\dagger)^*(f) = (\hat{\gamma}^\dagger)^\circ(f) = \tilde{\gamma}(f)$. Starting from the inclusion $\Delta^{(2)}_0(\gamma_0) \subseteq (\iota \otimes \Delta)\Delta(\gamma)$, the statement concerning γ^*_r is proven in the same way. ☐

Let us further reduce the 3-dimensional problem to a 2-dimensional one. If $\theta \in q^{2\mathbb{Z}} \cup -q^{-2\mathbb{Z}}$, we define $L^2_\theta(I^3_{q^2}, \Omega)$ to be the closed subspace of $L^2_\theta(I^3_{q^2}, \Omega)$ consisting of all functions that have support in $L(\theta)$. Then $L^2(I^3_{q^2}, \Omega)$ is the orthogonal sum $\oplus_{\theta \in q^{2\mathbb{Z}} \cup -q^{-2\mathbb{Z}}} L^2_\theta(I^3_{q^2}, \Omega)$. We know that $\tilde{F}^k_{m,p}, \tilde{G}^k_{m,p} \in L^2_{\mathrm{sgn}(p)\, q^{2k}}(I^3_{q^2}, \Omega)$ for all $p \in I_{q^2}$, $m, k \in \mathbb{Z}$.

Thus, lemma 5.5 implies that both families $(\,\tilde{F}_{m,p}^{\log_{q^2}|\theta|} \mid p \in I_{q^2}$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta), m \in \mathbb{Z}\,)$ and $(\,\tilde{G}_{m,p}^{\log_{q^2}|\theta|} \mid p \in I_{q^2}$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta), m \in \mathbb{Z}\,)$ are orthonormal bases of $L^2_\theta(I^3_{q^2}, \Omega)$.

It follows from the definition of γ_l and γ_r that $L^2_\theta(I^3_{q^2}, \Omega)$ is invariant under γ_l and γ_r.

Before starting the proof of the next result, let us first introduce some notation. Therefore suppose that $C \subseteq (-q^{2\mathbb{Z}} \cup q^{2\mathbb{Z}}) \times (-q^{2\mathbb{Z}} \cup q^{2\mathbb{Z}})$ and consider a function $f : C \to \mathbb{C}$. Then we define new functions $\partial_1 f, \partial_2 f : C \to \mathbb{C}$ such that $(\partial_1 f)(x,y) = \frac{f(q^2x,y)-f(x,y)}{x}$ and $(\partial_2 f)(x,y) = \frac{f(x,q^2y)-f(x,y)}{y}$ for all $(x,y) \in C$ (recall discussion (2) in Notation and conventions).

Proposition 5.8. *We have that* $\gamma_l = \gamma_r$.

Proof. Fix $\theta \in -q^{2\mathbb{Z}} \cup q^{2\mathbb{Z}}$. Set $k = \log_{q^2}|\theta|$. On $K(\theta)$ we define the measure Ω_θ such that

$$\Omega_\theta(\{(x,y)\}) = |x|\,|y|\,\frac{(-x,-y;q^2)_\infty}{(-\theta xy;q^2)_\infty}$$

for all $(x,y) \in K(\theta)$. Define a unitary transformation $\Xi : L_\theta(I^3_{q^2}, \Omega) \to L^2(K(\theta), \Omega_\theta)$ such that $(\Xi f)(x,y) = f(x, \theta xy, y)$ for all $f \in L_\theta(I^3_{q^2}, \Omega)$ and $(x,y) \in K(\theta)$. Notice that $\Xi \tilde{F}^k_{m,p} = F^k_{m,p}$ and $\Xi \tilde{G}^k_{m,p} = G^k_{m,p}$ for all $m \in \mathbb{Z}$ and $p \in I_{q^2}$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta)$. Set $\gamma^\theta_r = \Xi\, \gamma_r \restriction_{L_\theta(I^3_{q^2},\Omega)} \Xi^*$ and $\gamma^\theta_l = \Xi\, \gamma_l \restriction_{L_\theta(I^3_{q^2},\Omega)} \Xi^*$. Then,

(1) the space $\langle\, F^k_{m,p} \mid p \in I_{q^2}$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta), m \in \mathbb{Z}\,\rangle$ is a core of γ^θ_l,

(2) the space $\langle\, G^k_{m,p} \mid p \in I_{q^2}$ such that $\mathrm{sgn}(p) = \mathrm{sgn}(\theta), m \in \mathbb{Z}\,\rangle$ is a core of $(\gamma^\theta_r)^*$. (5.19)

Choose $f \in D(\gamma^\theta_l)$ and $g \in D((\gamma^\theta_r)^*)$. By Lemma 5.7, we know that for $(x,y) \in K(\theta)$,

$$(\gamma^\theta_l\, f)(x,y) = q\,|\theta|^{-\frac{1}{2}}\,\frac{1}{xy}\,\big(\,(1+q^{-2}y)\;f(q^2x, q^{-2}y) - (1+q^{-2}y)\;f(x, q^{-2}y) - (1+\theta xy)\;f(q^2x, y) + f(x,y)\,\big) \tag{5.20}$$

and

$$((\gamma^\theta_r)^*\, g)(x,y) = q\,|\theta|^{-\frac{1}{2}}\,\frac{1}{xy}\,\big(\,(1+q^{-2}x)\;g(q^{-2}x, q^2y) - (1+\theta xy)\;g(x, q^2y) - (1+q^{-2}x)\;g(q^{-2}x, y) + g(x,y)\,\big)\,. \tag{5.21}$$

Note the symmetry between the above formulas for γ^θ_l and $(\gamma^\theta_r)^*$ with respect to interchanging x and y.

Define the auxiliary function $h : K(\theta) \to \mathbb{C}$ such that $h(x,y) = \frac{(-x,-y;q^2)_\infty}{(-\theta xy;q^2)_\infty}$ for all $(x,y) \in K(\theta)$.

We want to show that $\langle \gamma^\theta_l f, g\rangle = \langle f, (\gamma^\theta_r)^* g\rangle$. The difference of these inner products are sums over the whole area $K(\theta)$ but we will start by approximating these sums by sums over a finite subset of $K(\theta)$ obtained by cutting off $K(\theta)$ by 4 rectangles (see Fig. 1 and Fig. 2 later in the proof). Then we let these rectangles get bigger and bigger. In this way, these finite sums converge certainly to the sum over the whole of $K(\theta)$. In the second part of the proof, we will then prove that these finite sums converge to 0, implying that $\langle \gamma^\theta_l f, g\rangle$ must be equal to $\langle f, (\gamma^\theta_r)^* g\rangle$.

So let us first calculate the contribution of one rectangle, lying in one of the quadrants, to the difference of the above inner product. Therefore let $a,b,c,d \in I_{q^2}$ such that $\mathrm{sgn}(a) = \mathrm{sgn}(b)$, $\mathrm{sgn}(c) = \mathrm{sgn}(d)$, $|a| < |b|$ and $|c| < |d|$ and set

$$C(a,b;c,d) = \sum_{\substack{(x,y)\,\in\, K(\theta) \\ x \text{ between } a \text{ and } b \\ y \text{ between } c \text{ and } d}} \big(\,(\gamma^\theta_l f)(x,y)\,\bar{g}(x,y)\; - f(x,y)\,\overline{((\gamma^\theta_r)^* g)(x,y)}\,\big)\,|x|\,|y|\,h(x,y)\;. \tag{5.22}$$

where the statement ‘x between a and b’ allows for x to be equal to a or b as well (similarly for y).

Using Eqs. (5.20) and (5.21) and recalling discussion (2) in Notations and conventions, we get that

$$
\begin{aligned}
&q^{-1}\,|\theta|^{\frac{1}{2}}\,\mathrm{sgn}(ac)\,C(a,b;c,d)\\
&\quad=\sum_{x=a}^{b}\sum_{y=c}^{d}(1+q^{-2}y)\ f(q^2x,q^{-2}y)\,\bar{g}(x,y)\,h(x,y)-\sum_{x=a}^{b}\sum_{y=c}^{d}(1+q^{-2}x)\ f(x,y)\,\bar{g}(q^{-2}x,q^2y)\,h(x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=c}^{d}(1+q^{-2}y)\ f(x,q^{-2}y)\,\bar{g}(x,y)\,h(x,y)+\sum_{x=a}^{b}\sum_{y=c}^{d}(1+\theta xy)\ f(x,y)\bar{g}(x,q^2y)\,h(x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=c}^{d}(1+\theta xy)\ f(q^2x,y)\,\bar{g}(x,y)\,h(x,y)+\sum_{x=a}^{b}\sum_{y=c}^{d}(1+q^{-2}x)\ f(x,y)\,\bar{g}(q^{-2}x,y)\,h(x,y)\\
&\quad=\sum_{x=a}^{b}\sum_{y=q^{-2}c}^{q^{-2}d}(1+y)\ f(q^2x,y)\,\bar{g}(x,q^2y)\,h(x,q^2y)-\sum_{x=q^{-2}a}^{q^{-2}b}\sum_{y=c}^{d}(1+x)\ f(q^2x,y)\,\bar{g}(x,q^2y)\,h(q^2x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=q^{-2}c}^{q^{-2}d}(1+y)\ f(x,y)\,\bar{g}(x,q^2y)\,h(x,q^2y)+\sum_{x=a}^{b}\sum_{y=c}^{d}(1+\theta xy)\ f(x,y)\bar{g}(x,q^2y)\,h(x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=c}^{d}(1+\theta xy)\ f(q^2x,y)\,\bar{g}(x,y)\,h(x,y)+\sum_{x=q^{-2}a}^{q^{-2}b}\sum_{y=c}^{d}(1+x)\ f(q^2x,y)\,\bar{g}(x,y)\,h(q^2x,y)\ .
\end{aligned}
\tag{5.23}
$$

Define the set $S=\{\,(x,y)\in(-q^{2\mathbb{Z}}\cup q^{2\mathbb{Z}})\times(-q^{2\mathbb{Z}}\cup q^{2\mathbb{Z}})\mid q^2\theta xy\in I_{q^2}\,\}$. Define a new auxiliary function $h':S\to\mathbb{C}$ such that $h'(x,y)=\frac{(-x,-y;q)_\infty}{(-q^2\theta xy;q^2)_\infty}$ for all $(x,y)\in S$. Now observe the following basic facts:

$$(1+y)h(x,q^2y)=h'(x,y)\quad\text{if } (x,q^2y)\in K(\theta)\qquad\qquad (1+x)h(q^2x,y)=h'(x,y)\quad\text{if } (q^2x,y)\in K(\theta)$$

$$(1+\theta xy)h(x,y)=h'(x,y)\quad\text{if } (x,y)\in K(\theta)\ .$$

Remembering that f and g are defined on $K(\theta)$, these facts combined with Eq. (5.23) imply that

$$
\begin{aligned}
&q^{-1}\,|\theta|^{\frac{1}{2}}\,\mathrm{sgn}(ac)\,C(a,b;c,d)\\
&\quad=\sum_{x=a}^{b}\sum_{y=q^{-2}c}^{q^{-2}d}f(q^2x,y)\,\bar{g}(x,q^2y)\,h'(x,y)-\sum_{x=q^{-2}a}^{q^{-2}b}\sum_{y=c}^{d}f(q^2x,y)\,\bar{g}(x,q^2y)\,h'(x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=q^{-2}c}^{q^{-2}d}f(x,y)\,\bar{g}(x,q^2y)\,h'(x,y)+\sum_{x=a}^{b}\sum_{y=c}^{d}f(x,y)\bar{g}(x,q^2y)\,h'(x,y)\\
&\qquad-\sum_{x=a}^{b}\sum_{y=c}^{d}f(q^2x,y)\,\bar{g}(x,y)\,h'(x,y)+\sum_{x=q^{-2}a}^{q^{-2}b}\sum_{y=c}^{d}f(q^2x,y)\,\bar{g}(x,y)\,h'(x,y)\ .
\end{aligned}
$$

In the above sum, most of the terms of the sums in the left column are cancelled by the terms of the sums in the right column. What remains is

$$
\begin{aligned}
&q^{-1}\,|\theta|^{\frac{1}{2}}\,\mathrm{sgn}(ac)\,C(a,b;c,d)\\
&\quad=\sum_{x=a}^{b}f(q^2x,q^{-2}d)\,\bar{g}(x,d)\,h'(x,q^{-2}d)+\sum_{y=q^{-2}c}^{d}f(q^2a,y)\,\bar{g}(a,q^2y)\,h'(a,y)\\
&\qquad-\sum_{x=q^{-2}a}^{b}f(q^2x,c)\,\bar{g}(x,q^2c)\,h'(x,c)-\sum_{y=c}^{d}f(b,y)\,\bar{g}(q^{-2}b,q^2y)\,h'(q^{-2}b,y)\\
&\qquad-\sum_{x=a}^{b}f(x,q^{-2}d)\,\bar{g}(x,d)\,h'(x,q^{-2}d)+\sum_{x=a}^{b}f(x,c)\bar{g}(x,q^2c)\,h'(x,c)\\
&\qquad-\sum_{y=c}^{d}f(q^2a,y)\,\bar{g}(a,y)\,h'(a,y)+\sum_{y=c}^{d}f(b,y)\,\bar{g}(q^{-2}b,y)\,h'(q^{-2}b,y)
\end{aligned}
$$

$$\begin{aligned}
&= \sum_{x=a}^{b} (f(q^2x, q^{-2}d) - f(x, q^{-2}d)\,)\,\bar{g}(x,d)\,h'(x,q^{-2}d) \\
&\quad - \sum_{y=c}^{d} f(b,y)\,(\bar{g}(q^{-2}b, q^2y) - \bar{g}(q^{-2}b,y)\,)\,h'(q^{-2}b,y) \\
&\quad + f(a,c)\,\bar{g}(a,q^2c)h'(a,c) - \sum_{x=q^{-2}a}^{b} (f(q^2x,c) - f(x,c)\,)\,\bar{g}(x,q^2c)\,h'(x,c) \\
&\quad - f(q^2a,c)\,\bar{g}(a,c)h'(a,c) + \sum_{y=q^{-2}c}^{d} f(q^2a,y)\,(\bar{g}(a,q^2y) - \bar{g}(a,y)\,)\,h'(a,y)\ . \qquad (5.24)
\end{aligned}$$

Set $I = \{\, v \in I_{q^2}^{+} \mid v < q^2, v < |\theta|^{-1} \text{ and } v < |\theta|^{-\frac{1}{2}} q \,\}$. If $v \in I$ we define $v' = q^2/v|\theta|$. Let us calculate the contribution of the 4 rectangles depicted in Fig. 1 and Fig. 2 to $\langle \gamma_l^\theta f, g\rangle - \langle f, (\gamma_r^\theta)^* g\rangle$. Therefore set $C(v) = C(v,v';v,v') + C(-v,-q^2;v,v') + C(-v,-q^2;-v,-q^2) + C(v,v';-v,-q^2)$.

It is clear from Eq. (5.22) and the dominated convergence theorem that $C(v)$ converges to $\langle \gamma_l^\theta f, g\rangle - \langle f, (\gamma_r^\theta)^* g\rangle$ as $v \to 0$. (5.25)



Fig. 1 : $\theta > 0$



Fig. 2 : $\theta < 0$

If $\theta < 0$, we set

$$C_{\infty,1}(v) = \sum_{x=-v}^{-q^2} (\partial_1 f)(x, q^{-2}v')\,\bar{g}(x,v')\,h'(x,q^{-2}v')\,|x| - \sum_{y=-v}^{-q^2} f(v',y)\,(\partial_2 \bar{g})(q^{-2}v',y)\,h'(q^{-2}v',y)\,|y|$$

and if $\theta > 0$, we set

$$C_{\infty,1}(v) = \sum_{x=v}^{v'} (\partial_1 f)(x, q^{-2}v')\,\bar{g}(x,v')\,h'(x,q^{-2}v')\,|x| - \sum_{y=v}^{v'} f(v',y)\,(\partial_2 \bar{g})(q^{-2}v',y)\,h'(q^{-2}v',y)\,|y|\ .$$

Regardless of the sign of θ, we set

$$\begin{aligned}
C_{\infty,0}(v) \quad = \quad & \operatorname{sgn}(\theta)\,f(v', -\operatorname{sgn}(\theta)\,v)\,\bar{g}(q^{-2}v', -\operatorname{sgn}(\theta)\,q^2 v)\,h'(q^{-2}v', -\operatorname{sgn}(\theta)\,v) \\
& -\operatorname{sgn}(\theta)\,f(-\operatorname{sgn}(\theta)\,q^2 v, q^{-2}v')\bar{g}(-\operatorname{sgn}(\theta)\,v, v')\,h'(-\operatorname{sgn}(\theta)\,v, q^{-2}v')
\end{aligned}$$

and

$$\begin{aligned} C_{0,0}(v) &= f(v,v)\bar{g}(v,q^2v)\,h'(v,v) - f(q^2v,v)\,\bar{g}(v,v)\,h'(v,v) \\ &\quad -f(-v,v)\,\bar{g}(-v,q^2v)\,h'(-v,v) + f(-q^2v,v)\,\bar{g}(-v,v)\,h'(-v,v) \\ &\quad +f(-v,-v)\,\bar{g}(-v,-q^2v)\,h'(-v,-v) - f(-q^2v,-v)\,\bar{g}(-v,-v)\,h'(-v,-v) \\ &\quad -f(v,-v)\,\bar{g}(v,-q^2v)\,h'(v,-v) + f(q^2v,-v)\,\bar{g}(v,-v)\,h'(v,-v)\ , \end{aligned}$$

$$\begin{aligned} C_{0,1}(v) &= \sum_{y=q^{-2}v}^{v'} f(q^2v,y)\,(\partial_2\bar{g})(v,y)\,h'(v,y)\,|y| - \sum_{y=q^{-2}v}^{v'} f(-q^2v,y)\,(\partial_2\bar{g})(-v,y)\,h'(-v,y)\,|y|\ , \qquad (5.26) \\ C_{0,2}(v) &= \sum_{y=-q^{-2}v}^{-q^2} f(q^2v,y)\,(\partial_2\bar{g})(v,y)\,h'(v,y)\,|y| - \sum_{y=-q^{-2}v}^{-q^2} f(-q^2v,y)\,(\partial_2\bar{g})(-v,y)\,h'(-v,y)\,|y|\ , \\ C_{0,3}(v) &= -\sum_{x=q^{-2}v}^{v'} (\partial_1 f)(x,v)\,\bar{g}(x,q^2v)\,h'(x,v)\,|x| + \sum_{x=q^{-2}v}^{v'} (\partial_1 f)(x,-v)\,\bar{g}(x,-q^2v)\,h'(x,-v)\,|x|\ , \\ C_{0,4}(v) &= -\sum_{x=-q^{-2}v}^{-q^2} (\partial_1 f)(x,v)\,\bar{g}(x,q^2v)\,h'(x,v)\,|x| + \sum_{x=-q^{-2}v}^{-q^2} (\partial_1 f)(x,-v)\,\bar{g}(x,-q^2v)\,h'(x,-v)\,|x|\ . \end{aligned}$$

A bookkeeping exercise based on Eq. (5.24) reveals that

$$q^{-1}\,|\theta|^{\frac{1}{2}}\,C(v) = C_{\infty,0}(v) + C_{\infty,1}(v) + \sum_{i=0}^{4} C_{0,i}(v)\ . \qquad (5.27)$$

In order to show all the nitty gritty work involved, let us calculate $C(-v,-q^2;v,v')$ in the case of $\theta < 0$ (keep Fig. 1 in mind). So in Eq. (5.24) we have to set $a=-v$, $b=-q^2$, $c=v$ and $d=v'$.

(1) If $x \in I_{q^2} \cap [-q^2,-v]$, then $f(x,q^{-2}v')=0$ since $(x,q^{-2}v') \notin K(\theta)$. If $x \in I_{q^2} \cap [-q^2,-v)$ then also $f(q^2x,q^{-2}v')=0$ since $(q^2x,q^{-2}v') \notin K(\theta)$. But note that $(-q^2v,q^{-2}v') \in K(\theta)$. So the first sum of expression (5.24) equals $f(-q^2v,q^{-2}v')\,\bar{g}(-v,v')\,h'(-v,q^{-2}v')$.

(2) For all $y \in I_{q^2} \cap [v,v']$, clearly $\bar{g}(-1,q^2y)=\bar{g}(-1,y)=0$. Thus the second sum in expression (5.24) equals 0.

(3) For all $x \in I_{q^2} \cap [-q^2,-q^{-2}v]$, we have that $f(q^2x,v)-f(x,v) = -|x|\,(\partial_1 f)(x,v)$ and for all $y \in I_{q^2} \cap [v,v']$, we have that $\bar{g}(-v,q^2y)-\bar{g}(-v,y)=|y|\,(\partial_2\bar{g})(-v,y)$.

Putting all these results into Eq. (5.24), we see that

$$\begin{aligned} &-q^{-1}\,|\theta|^{\frac{1}{2}}\,C(-v,-q^2;v,v') \\ &\quad = f(-q^2v,q^{-2}v')\,\bar{g}(-v,v')\,h'(-v,q^{-2}v') + f(-v,v)\,\bar{g}(-v,q^2v)\,h'(-v,v) - f(-q^2v,v)\,\bar{g}(-v,v)\,h'(-v,v) \\ &\qquad + \sum_{x=-q^{-2}v}^{-q^2} (\partial_1 f)(x,v)\,\bar{g}(x,q^2v)\,h'(x,v)\,|x| + \sum_{y=q^{-2}v}^{v'} f(-q^2v,y)\,(\partial_2\bar{g})(-v,y)\ h'(-v,y)\,|y|\ . \end{aligned}$$

Similarly, one calculates $C(v,v';v,v')$, $C(-v,-q^2;-v,-q^2)$ and $C(v,v';-v,-q^2)$. Adding these four results together, one finds Eq. (5.27). The case $\theta < 0$ is treated similarly.

Now we are going to make specific choices for our functions f and g. Therefore take $m,n \in \mathbb{Z}$, $p,t \in I_{q^2}$ such that $\mathrm{sgn}(p)=\mathrm{sgn}(t)=\mathrm{sgn}(\theta)$ and define f and g such that for $(x,y) \in K(\theta)$,

$$f(x,y) = \Psi\left(\begin{matrix} -q^2/p \\ q^{2(1-m)}\,y/|p| \end{matrix}\ ;\ \mathrm{sgn}(\theta)\,q^{2(1-m)}\right)\ \Psi\left(\begin{matrix} -\mathrm{sgn}(\theta)\,q^{2(1+m)}/y \\ q^{2(1+m+k)}x \end{matrix}\ ;\ \mathrm{sgn}(\theta)\,q^{2(1+k)}y\right) \qquad (5.28)$$

if $\mathrm{sgn}(\theta)\,q^{-2m}y \in I_{q^2}$ and $f(x,y)=0$ if $\mathrm{sgn}(\theta)\,q^{-2m}y \notin I_{q^2}$, and on the other hand,

$$g(x,y) = \Psi\left(\begin{matrix} -q^2/t \\ q^{2(1-n)}\,x/|t| \end{matrix}\ ;\ \mathrm{sgn}(\theta)\,q^{2(1-n)}\right)\ \Psi\left(\begin{matrix} -\mathrm{sgn}(\theta)\,q^{2(1+n)}/x \\ q^{2(1+k+n)}y \end{matrix}\ ;\ \mathrm{sgn}(\theta)\,q^{2(1+k)}x\right) \qquad (5.29)$$

if $\operatorname{sgn}(\theta)\, q^{-2n} x \in I_{q^2}$ and $f(x,y) = 0$ if $\operatorname{sgn}(\theta)\, q^{-2n} x \notin I_{q^2}$. By Eq. (5.16) and Result 6.4, $F^k_{m,p}$,$G^k_{-n,t}$ are proportional to f,g respectively. In the next part we will show that for this choice of f and g, $C(v) \to 0$ as $v \to 0$.

Notice that g is obtained from f by interchanging x and y and replacing the parameters m,p by n,t. There is also some symmetry with respect to interchanging x and y in the formulas defining $C_{\infty,0}(v)$ and $C_{\infty,1}(v)$. Similarly, note the symmetry with respect to this variable interchange between $C_{0,1}(v)$, $C_{0,2}(v)$ and $C_{0,3}(v)$, $C_{0,4}(v)$ respectively. This is of course in correspondence with the symmetry observed in Eqs. (5.20) and (5.21). We will make use of this symmetry to reduce our work.

Let $a \in \mathbb{C}$. Then $(\,(aq^2z;q^2)_\infty - (az;q^2)_\infty\,)/z = a\,(aq^2z;q^2)_\infty$ for all $z \in \mathbb{C}$. Using this simple fact, a careful inspection learns that for $(x,y) \in K(\theta)$,

$$(\partial_1 f)(x,y) = q^{2(1+m+k)}\ \Psi\left(\begin{matrix} -q^2/p \\ q^{2(1-m)}\, y/|p| \end{matrix}\ ;\, \operatorname{sgn}(\theta)\, q^{2(1-m)}\right)\ \Psi\left(\begin{matrix} -\operatorname{sgn}(\theta)\, q^{2(1+m)}/y \\ q^{2(2+m+k)} x \end{matrix}\ ;\, \operatorname{sgn}(\theta)\, q^{2(2+k)} y\right)$$

if $\operatorname{sgn}(\theta)\, q^{-2m} y \in I_{q^2}$ and $(\partial_1 f)(x,y) = 0$ if $\operatorname{sgn}(\theta)\, q^{-2m} y \notin I_{q^2}$. On the other hand,

$$(\partial_2 g)(x,y) = q^{2(1+n+k)}\ \Psi\left(\begin{matrix} -q^2/t \\ q^{2(1-n)}\, x/|t| \end{matrix}\ ;\, \operatorname{sgn}(\theta)\, q^{2(1-n)}\right)\ \Psi\left(\begin{matrix} -\operatorname{sgn}(\theta)\, q^{2(1+n)}/x \\ q^{2(2+n+k)} y \end{matrix}\ ;\, \operatorname{sgn}(\theta)\, q^{2(2+k)} x\right) \tag{5.30}$$

if $\operatorname{sgn}(\theta)\, q^{-2n} x \in I_{q^2}$ and $(\partial_2 g)(x,y) = 0$ if $\operatorname{sgn}(\theta)\, q^{-2n} x \notin I_{q^2}$.

Now we will show that the different summands of $C(v)$ as described in (5.27) converge to 0 as $v \to 0$.

(1) First we quickly check that $C_{0,0}(v) \to 0$ as $v \to 0$.

Let $(\,(x_r,y_r)\,)_{r=1}^\infty$ be a sequence in $K(\theta)$ that converges to $(0,0)$. Then $\operatorname{sgn}(\theta)\, q^{-2m} y_r \in I_{q^2}$ and $\operatorname{sgn}(\theta)\, q^{-2n} x_r \in I_{q^2}$ for r big enough. Therefore Eqs. (5.28) and (5.29) imply, as in the proof of Lemma 6.2, that

$$\begin{aligned} f(x_r,y_r) &\to \Psi(-q^2/p;0;\operatorname{sgn}(\theta) q^{2(1-m)})\, \phi(0;-q^{2(2+m+k)}) \qquad \text{as } r \to \infty\ , \\ g(x_r,y_r) &\to \Psi(-q^2/t;0;\operatorname{sgn}(\theta) q^{2(1-n)})\, \phi(0;-q^{2(2+n+k)}) \qquad \text{as } r \to \infty\ , \end{aligned}$$

where we used notation (5.4). We have also that $h'(x_r,y_r) \to 1$ as $r \to \infty$. From this all, we easily conclude that $C_{0,0}(v) \to 0$ as $v \to 0$.

(2) Let us now deal with $C_{\infty,1}(v)$. First consider the case where $\theta < 0$. Assume for the moment that $v \leq q^{-2m}/|\theta|$ and $v \leq q^{-2n}/|\theta|$. Then $q^{-2m}(q^{-2}v') \geq 1$ and $q^{-2n}(q^{-2}v') \geq 1$. Hence, $\operatorname{sgn}(\theta)\, q^{-2m}\,(q^{-2}v')$ and $\operatorname{sgn}(\theta)\, q^{-2n}\,(q^{-2}v')$ do not belong to I_{q^2}. As a consequence, $(\partial_1 f)(x,q^{-2}v') = (\partial_2 g)(q^{-2}v',x) = 0$ for all $x \in I_{q^2} \cap [-q^2,-v]$. We conclude that $C_{\infty,1}(v) = 0$ if $v \leq q^{-2m}/|\theta|$ and $v \leq q^{-2n}/|\theta|$. Thus $C_{\infty,1}(v) \to 0$ as $v \to 0$.

Now we look at the more challenging case where $\theta > 0$. Referring to Definition 3.1, Eq. (5.25) and Result 6.4, we get the existence of a constant $D > 0$, only depending on m, p and θ such that

$$\begin{aligned} &|\,(\partial_1 f)(x,q^{-2}y)|^2\ h'(x,q^{-2}y)\, x\, y \\ &\quad = D\ |a_{\kappa^{-1}(p)}(\kappa^{-1}(q^{-2m})\kappa^{-1}(q^{-2}y),\kappa^{-1}(q^{-2}y))|^2\ |a_{\kappa^{-1}(q^{-2(m+1)}y)}(\kappa^{-1}(\theta y)\,\kappa^{-1}(x),\kappa^{-1}(x))|^2 \end{aligned}$$

for all $x,y \in I^+_{q^2}$. Therefore Proposition 3.2 implies that

$$\begin{aligned} \sum_{y \in I^+_{q^2}} \sum_{x \in I^+_{q^2}} |\,(\partial_1 f)(x,q^{-2}y)|^2\ h'(x,q^{-2}y)\, x\, y &= D \sum_{y \in I^+_{q^2}} |a_{\kappa^{-1}(p)}(\kappa^{-1}(q^{-2m})\kappa^{-1}(q^{-2}y),\kappa^{-1}(q^{-2}y))|^2 \\ &\quad \times \sum_{x \in I^+_{q^2}} |a_{\kappa^{-1}(q^{-2(m+1)}y)}(\kappa^{-1}(\theta y)\,\kappa^{-1}(x),\kappa^{-1}(x))|^2 \\ &\leq D \sum_{y \in I^+_{q^2}} |a_{\kappa^{-1}(p)}(\kappa^{-1}(q^{-2m})\kappa^{-1}(q^{-2}y),\kappa^{-1}(q^{-2}y))|^2\ \leq\ D\ . \end{aligned}$$

Thus, $\sum_{y \in I^+_{q^2}} y \sum_{x \in I^+_{q^2}} |(\partial_1 f)(x,q^{-2}y)|^2\ h'(x,q^{-2}y)\, x < \infty$, which clearly implies that $\sum_{x \in I^+_{q^2}} |\,(\partial_1 f)(x,q^{-2}y)|^2\ h'(x,q^{-2}y)\, x \to 0$ as $y \to \infty$.

On the other hand,

$$\sum_{\substack{y\,\in\, I_{q^2}^+\\ y\ge 1}}\ \sum_{x\in I_{q^2}^+} |g(x,y)|^2\, h'(x,q^{-2}y)\,x = \sum_{\substack{y\,\in\, I_{q^2}^+\\ y\ge 1}}\ \sum_{x\in I_{q^2}^+} |g(x,y)|^2\,\frac{(-x,-y;q^2)_\infty}{(-\theta xy;q^2)_\infty}\,(1+q^{-2}y)\,x$$

$$\le (1+q^{-2}) \sum_{\substack{y\,\in\, I_{q^2}^+\\ y\ge 1}}\ \sum_{x\in I_{q^2}^+} |g(x,y)|^2\,\frac{(-x,-y;q^2)_\infty}{(-\theta xy;q^2)_\infty}\,x\,y \ < \infty\ ,$$

since g belongs to $L^2(K(\theta),\Omega_\theta)$. So we also get that $\sum_{x\in I_{q^2}^+} |g(x,y)|^2\ h'(x,q^{-2}y)\,x \to 0$ as $y\to\infty$. By the Cauchy-Schwarz inequality, we have

$$\Big|\sum_{x=v}^{v'}(\partial_1 f)(x,q^{-2}v')\,\bar g(x,v')\,h'(x,q^{-2}v')\,x\,\Big|^2$$
$$\le \Big(\sum_{x\in I_{q^2}^+} |\,(\partial_1 f)(x,q^{-2}v')|^2\,h'(x,q^{-2}v')\,x\Big)\Big(\sum_{x\in I_{q^2}^+} |g(x,v')|^2\,h'(x,q^{-2}v')\,x\Big)\ .$$

Therefore the considerations above imply that

$$\sum_{x=v}^{v'}(\partial_1 f)(x,q^{-2}v')\,\bar g(x,v')\,h'(x,q^{-2}v')\,x\ \to 0 \qquad \text{as}\ \ v\to 0\ .$$

Hence, the obvious symmetry between f and g guarantees that also

$$\sum_{y=v}^{v'} f(v',y)\,(\partial_2 \bar g)(q^{-2}v',y)\,h'(q^{-2}v',y)\,y\ \to 0 \qquad \text{as}\ \ v\to 0\ .$$

Thus, we conclude that $C_{\infty,1}(v)\to 0$ as $v\to 0$.

(3) In the next step we prove that $C_{0,1}(v)+C_{0,2}(v)\to 0$ as $v\to 0$. (5.31)

Let $v\in I$ such that $v\le q^{2(n+1)}$. Then $|q^{-2n}v|\le q^2$, thus $\pm\mathrm{sgn}(\theta)\,q^{-2n}v\in I_{q^2}$. Therefore Eqs. (5.28), (5.30) and Result 6.4 imply for $y\in I_{q^2}\cap([-q^2,-q^{-2}v]\cup[q^{-2}v,v'])$,

$$\begin{aligned} f(\pm q^2 v,y)\,(\partial_2\bar g)(\pm v,y)\,h'(\pm v,y)\,|y| \ &= q^{2(1+n+k)}\,(\mp v,-y;q^2)_\infty\,(\mp q^2\theta v y;q^2)_\infty^{-1}\,|y| \\ &\Psi(-q^2/p;q^{2(1-m)}y/|p|;\mathrm{sgn}(\theta)q^{2(1-m)})\quad \Psi(\mp 1/v;\mathrm{sgn}(\theta)q^{2(1+k)}y;\pm q^{2(2+m+k)}v) \\ &\Psi(-q^2/t;\pm q^{2(1-n)}v/|t|;\mathrm{sgn}(\theta)\,q^{2(1-n)})\quad \Psi(\mp\mathrm{sgn}(\theta)q^{2(1+n)}/v;q^{2(2+n+k)}y;\pm\mathrm{sgn}(\theta)q^{2(2+k)}v) \end{aligned} \tag{5.32}$$

if $\mathrm{sgn}(\theta)q^{-2m}y\in I_{q^2}$ and $f(\pm q^2v,y)\,(\partial_2\bar g)(\pm v,y)\,h'(\pm v,y)\,|y|\ =0$ if $\mathrm{sgn}(\theta)q^{-2m}y\notin I_{q^2}$ (since $f(\pm q^2v,y)$ $=0$ in this case).

If $v\in I$ and $y\in I_{q^2}\cap([-q^2,-q^{-2}v]\cup[q^{-2}v,v'])$, then $(\pm v,y)\in K(\theta)$, thus $\pm q^2\theta v y\ge -q^2$, so $(\mp q^2\theta vy;q^2)_\infty\ge (q^2;q^2)_\infty$. So we get easily the existence of a constant $C>0$ such that

$$|q^{2(1+n+k)}\,(\mp v;q^2)_\infty\,(\mp q^2\theta v y;q^2)_\infty^{-1}\,\Psi(-q^2/t;\pm q^{2(1-n)}v/|t|;\mathrm{sgn}(\theta)\,q^{2(1-n)})|\le C$$

for all $v\in I$ such that $v\le q^{2(n+1)}$ and $y\in I_{q^2}\cap([-q^2,-q^{-2}v]\cup[q^{-2}v,v'])$.

Moreover, Lemma 6.8 implies the existence of

(a) a number $D>0$ such that $|\Psi(\mp 1/v;\mathrm{sgn}(\theta)q^{2(1+k)}y;\pm q^{2(2+m+k)}v)|\le D$ for all $y\in I_{q^2}$ such that $\mathrm{sgn}(\theta)\,q^{-2m}\,y\in I_{q^2}$ and $v\in I$ such that $v\le q^{2(n+1)}$.

(b) a function $H_1\in l^2(I_{q^2})^+$ such that $|y|^{\frac12}\,|(-y;q^2)_\infty|^{\frac12}\,|\Psi(-q^2/p;q^{2(1-m)}y/|p|;\mathrm{sgn}(\theta)q^{2(1-m)})|\le H_1(y)$ for all $y\in I_{q^2}$

(c) a function $H_2\in l^2(I_{q^2})^+$ such that $|y|^{\frac12}\,|(-y;q^2)_\infty|^{\frac12}\,|\Psi(\mp\mathrm{sgn}(\theta)q^{2(1+n)}/v;q^{2(2+n+k)}y;$ $\pm\mathrm{sgn}(\theta)q^{2(2+k)}v)|\le H_2(y)$ for all $y\in I_{q^2}$ and $v\in I$ such that $v\le q^{2(n+1)}$.

Now define $H \in l^1(I_{q^2})^+$ as $H = C\, D\, H_1\, H_2$. Then Eq. (5.32) implies that

$$|\, f(\pm q^2 v, y)\, (\partial_2 \bar{g})(\pm v, y)\, h'(\pm v, y)\, |y|\, | \leq H(y)$$

for $v \in I$ such that $v \leq q^{2(n+1)}$ and $y \in I_{q^2} \cap ([-q^2, -q^{-2}v] \cup [q^{-2}v, v'])$.
Define the function $G : I_{q^2} \to \mathbb{C}$ such that for $y \in I_{q^2}$,

$$\begin{aligned} G(y) \quad = \quad & q^{2(1+n+k)}\, (-y; q^2)_\infty\, |y|\ \Psi(-q^2/t; 0; \mathrm{sgn}(\theta)\, q^{2(1-n)})\ \Psi(-q^2/p; q^{2(1-m)} y/|p|; \mathrm{sgn}(\theta)\, q^{2(1-m)}) \\ & \phi(\mathrm{sgn}(\theta)\, q^{2(1+k)} y; -q^{2(2+m+k)})\ \phi(q^{2(2+n+k)} y; -q^{2(3+n+k)}) \end{aligned}$$

if $\mathrm{sgn}(\theta)\, q^{-2m} y \in I_{q^2}$ and $G(y) = 0$ if $\mathrm{sgn}(\theta)\, q^{-2m} y \notin I_{q^2}$ (here we use the notation introduced in (5.4)).
If we fix $y \in I_{q^2}$, we see that $y \in [-q^2, -q^{-2}v] \cup [q^{-2}v, v']$ if v is small enough. By Eq. (5.32), we have moreover that $f(\pm q^2 v, y)\, (\partial_2 \bar{g})(\pm v, y)\, h'(\pm v, y)\, |y| \to G(y)$ as $v \to 0$. Therefore the dominated convergence theorem implies that $G \in l^1(I_{q^2})$ and

$$\sum_{y=q^{-2}v}^{v'} f(\pm q^2 v, y)\, (\partial_2 \bar{g})(\pm v, y)\, h'(\pm v, y)\, |y| \ + \sum_{y=-q^2}^{-q^{-2}v} f(\pm q^2 v, y)\, (\partial_2 \bar{g})(\pm v, y)\, h'(\pm v, y)\, |y| \to \sum_{y \in I_{q^2}} G(y)$$

as $v \to 0$. Thus, it follows that $C_{0,1}(v) + C_{0,2}(v) \to 0$ as $v \to 0$. The aforementioned symmetry between f and g then guarantees that also $C_{0,3}(v) + C_{0,4}(v) \to 0$ as $v \to 0$.
(4) In the last step we look at $C_{\infty,0}(v)$. First assume that $\theta < 0$. If $v \in I$ and $v \leq q^{-2m}/|\theta|$ then $\mathrm{sgn}(\theta)\, q^{-2m}\, (q^{-2}v') = -q^{-2m}/v|\theta| \leq -1$. Thus Eq. (5.28) implies that $f(-\mathrm{sgn}(\theta)\, q^2 v, q^{-2}v')$ $\bar{g}(-\mathrm{sgn}(\theta)\, v, v')\, h'(-\mathrm{sgn}(\theta)\, v, q^{-2}v') = 0$. So in this case, $f(-\mathrm{sgn}(\theta)\, q^2 v, q^{-2}v')\, \bar{g}(-\mathrm{sgn}(\theta)\, v, v')$ $h'(-\mathrm{sgn}(\theta)\, v, q^{-2}v') \to 0$ as $v \to 0$.
Now we look at the more challenging case where $\theta > 0$. If $v \in I$ and $v \leq q^{2n+2}$, then $\mathrm{sgn}(\theta) q^{-2m}\, (q^{-2}v')$ and $-q^{-2n}v$ clearly belong to I_{q^2}; thus Eqs. (5.28) and (5.29) and Result 6.4 imply that

$$\begin{aligned} & |f(-\mathrm{sgn}(\theta)\, q^2 v, q^{-2}v')\, \bar{g}(-\mathrm{sgn}(\theta)\, v, v')\, h'(-\mathrm{sgn}(\theta)\, v, q^{-2}v')| = (-v, -1/v\theta; q^2)_\infty \\ & \quad (q^2; q^2)_\infty^{-1}\ |\, \Psi(-q^2/p; q^{2(1-m)}/v\theta p; q^{2(1-m)})\, |\ \ |\, \Psi(1/v; q^{2(1+k)}/v\theta; -q^{2(2+m+k)} v)\, | \\ & \quad |\, \Psi(-q^2/t; -q^{2(1-n)} v/t; q^{2(1-n)})\, |\ \ |\, \Psi(q^{2(1+n)}/v; q^{2(2+n+k)}/v\theta; -q^{2(1+k)} v)\, |\ . \end{aligned}$$

By Lemma 6.7, there exists a number $C > 0$ such that $|\, \Psi(-q^2/p; q^{2(1-m)}/v\theta p; q^{2(1-m)})\, | \leq C$ for all $v \in I$. Therefore the above equality and Lemma 6.9 allow us to conclude that

$$f(-\mathrm{sgn}(\theta)\, q^2 v, q^{-2}v')\, \bar{g}(-\mathrm{sgn}(\theta)\, v, v')\, h'(-\mathrm{sgn}(\theta)\, v, q^{-2}v') \to 0 \qquad \text{as} \ \ v \to 0$$

also in this case. The symmetry between f and g thus guarantees that also the first term of $C_{\infty,0}(v) \to 0$ as $v \to 0$. Therefore $C_{\infty,0}(v) \to 0$ as $v \to 0$.

Together with Eq. (5.27) the convergence results proven in parts (1),(2),(3) and (4) imply that $C(v) \to 0$ as $v \to 0$. By (5.25), this implies that $\langle \gamma_l^\theta f, g \rangle - \langle f, (\gamma_r^\theta)^* g \rangle = 0$. So, by our choice of f and g, we find that $\langle \gamma_l^\theta F_{m,p}^k, G_{-n,t}^k \rangle = \langle F_{m,p}^k, (\gamma_r^\theta)^* G_{-n,t}^k \rangle$.
Therefore the conditions stated in (5.19) imply that $\langle \gamma_l^\theta x', y' \rangle = \langle x', (\gamma_r^\theta)^* y' \rangle$ for all $x' \in D(\gamma_l^\theta)$ and $y' \in D((\gamma_l^\theta)^*)$. Consequently, $(\gamma_r^\theta)^* \subseteq (\gamma_l^\theta)^*$ (*). Since γ_l^θ and γ_r^θ are normal, this implies that $D(\gamma_r^\theta) = D((\gamma_r^\theta)^*) \subseteq D((\gamma_l^\theta)^*) = D(\gamma_l^\theta)$. But taking the adjoint of the inclusion (*), we arrive at the inclusion $\gamma_l^\theta \subseteq \gamma_r^\theta$. Hence, $\gamma_l^\theta = \gamma_r^\theta$. □

Since $U\, ((\Delta \otimes \iota)\Delta(\gamma))\, U^* = M_\zeta \otimes \gamma_l$ and $U\, ((\iota \otimes \Delta)\Delta(\gamma))\, U^* = M_\zeta \otimes \gamma_r$ (see the remarks before lemma 5.7) the previous proposition entails that $(\Delta \otimes \iota)\Delta(\gamma) = (\iota \otimes \Delta)\Delta(\gamma)$ and we have proven Proposition 5.2.

We can finally explain why the comultiplication Δ is defined in such a way that $U_\theta^* \Delta(\gamma^* \gamma) \restriction_{\ell'_\theta} U_\theta = 1 \otimes L_\theta^{\mathrm{sgn}(\theta)}$ (see the discussions surrounding Eq. (3.2) and after Proposition 3.9). Recall that this choice is reflected in the presence of the factor $s(x, y)$ in Definition 3.1.
In the proof above we showed that $C_{0,1}(v) \to 0$ as $v \to 0$, cf. (5.31). Looking at the defining formula (5.26) for $C_{0,1}(v)$, it is not too hard to imagine that for this to be true, we need at least that the functions f and g defined in Eqs. (5.28) and (5.29) have the same limit behavior when crossing the Y-axis, going from

the first to the the second quadrant (for these particular functions, these are just continuous transitions). Note that the relevance of f and g to the coassociativity of Δ is obvious from Lemma 5.5 (and that all potential discontinuous transitions are filtered out by U).
If one changes the comultiplication by leaving out the factor $s(x,y)$ in Definition 3.1, and thus opting for the equality $U_\theta^* \Delta(\gamma^*\gamma)\lceil_{\ell'_\theta} U_\theta = 1 \otimes L_\theta^1$, it turns out that f and g have different limit transitions implying that $C_{0,1}(v)$ does not converge to 0 as $v \to 0$. From this it would follow that $(\Delta \otimes \iota)\Delta(\gamma)$ and $(\iota \otimes \Delta)\Delta(\gamma)$ have different domains in this case.

From Lemma 5.5 we have two orthonormal bases for the same space $L^2(I_{q^2}^3, \Omega)$; $\big(\tilde{F}_{m,p}^k\big)_{p \in I_{q^2}, m,k \in \mathbb{Z}}$ and $\big(\tilde{G}_{m,p}^k\big)_{p \in I_{q^2}, m,k \in \mathbb{Z}}$. Hence, there exists a unitary operator $\mathcal{R}$ on $L^2(I_{q^2}^3, \Omega)$ mapping the first basis on the second. Its matrix elements are given by

$$\mathcal{R}(m_1, k_1, p_1; m_2, k_2, p_2) = \langle \tilde{F}_{m_1,p_1}^{k_1}, \tilde{G}_{m_2,p_2}^{k_2} \rangle_{L^2(I_{q^2}^3, \Omega)}$$

and these matrix elements of $\mathcal{R}$ can be thought of as Racah coefficients. It would be desirable to find an explicit expression for the Racah coefficients and to show that $\mathcal{R}(m_1, k_1, p_1; m_2, k_2, p_2) = 0$ for $p_1 \neq p_2$, since this would immediately imply $\gamma_l = \gamma_r$ and hence yield an alternative proof for Proposition 5.8 and hence of Proposition 5.2. However, we have not been able to carry out this programme.

6. APPENDIX

In this appendix we collect the basic properties of the Ψ-functions defined in Eq. (1) of the Introduction. Most of them stem from special function theory, so the proofs in this section are mainly intended for people that are not too familiar with this theory. As a general reference for q-hypergeometric functions we use [6]. Let us fix a number $0 < q < 1$.

Note first of all that the presence of the factor q^{k^2} in the series (1) implies that if we keep two of the parameters a,b,z fixed, the series converges uniformly on compact subsets in the remaining parameter. As a consequence, $\Psi(a;b;q,z)$ is analytic in the remaining parameter. Also, the function $\mathbb{C}^3 \to \mathbb{C} : (a,b,z) \mapsto \Psi(a;b;q,z)$ is analytic.
The Ψ-functions are closely related to the ${}_1\varphi_1$-functions (see [6, Def. (1.2.22)]) in the sense that $\Psi(a;b;q,z) = (b;q)_\infty \; {}_1\varphi_1(a;b;q,z)$ if $b \notin q^{-\mathbb{N}_0}$ (if $b \in q^{-\mathbb{N}_0}$, ${}_1\varphi_1(a;b;q,z)$ is not defined in general). As a consequence, a lot of the identities that are known for ${}_1\varphi_1$- functions extend by analytic continuation in the parameters to similar identities involving Ψ-functions.

A very basic but useful formula is known as the

Result 6.1 (θ-product identity)**.** *Consider $a \in \mathbb{C} \setminus \{0\}$, $k \in \mathbb{Z}$. Then*

$$(aq^k;q)_\infty \, (q^{1-k}/a;q)_\infty = (-a)^{-k} \, q^{-\frac{1}{2}k(k-1)} \, (a;q)_\infty \, (q/a;q)_\infty \; . \tag{6.1}$$

Proof. If $a \in q^{\mathbb{Z}}$, both sides of the above equation are easily seen to be equal to 0. Now we look at the case where $a \notin q^{\mathbb{Z}}$. Suppose that $k \geq 0$. Then we have for $n \in \mathbb{N}$,

$$\frac{(aq^k;q)_n}{(a;q)_{n+k}} \, \frac{(\,q^{1-k}/a;q)_{n+k}}{(\,q/a;q)_n} = \frac{\prod_{i=0}^{k-1}(1-q^{-i}/a)}{\prod_{i=0}^{k-1}(1-aq^i)} = \frac{(-a)^{-k}\,\prod_{i=0}^{k-1} q^{-i}\,\prod_{i=0}^{k-1}(1-aq^i)}{\prod_{i=0}^{k-1}(1-aq^i)} = (-a)^{-k}\, q^{-\frac{1}{2}k(k-1)}$$

so if we let n tend to infinity, Eq. (6.1) follows. If $k \leq 0$, we apply Eq. (6.1), where we replace k by $-k$ and a by q/a. ☐

The most important low order q-hypergeometric functions are the ${}_2\varphi_1$ functions. If $a,b,c,z \in \mathbb{C}$ satisfy $c \notin q^{-\mathbb{N}_0}$ and $|z| < 1$, these are defined as (see [6, Def. (1.2.22)] and the discussion thereafter)

$${}_2\varphi_1(a,b;c;q,z) = {}_2\varphi_1\left(\begin{array}{cc} a & b \\ \multicolumn{2}{c}{c} \end{array} ; q, z\right) = \sum_{n=0}^{\infty} \frac{(a;q)_n\,(b;q)_n}{(c;q)_n (q;q)_n}\, z^n \; .$$

If we keep a,b and c fixed, the function $z, |z| < 1 \mapsto {}_2\varphi_1(a,b;c;q,z)$ has an analytic extension to the set $\mathbb{C} \setminus [1,\infty)$ (see the beginning of [6, Sec. 4.3]). Of course, the value of this extension at $z \in \mathbb{C} \setminus [1,\infty)$ is also denoted by ${}_2\varphi_1(a,b;c;q,z)$. We will not use this fact directly in this paper but point to this extension because it is used in [3].

A connection between ${}_2\varphi_1$-functions and ${}_1\varphi_1$-functions is obtained by the following basic limit transition.

Lemma 6.2. *Consider $a,b,c,z \in \mathbb{C}$ such that $c \notin q^{-\mathbb{N}_0}$ and $(x_i)_{i=1}^{\infty}$ a sequence in $\mathbb{C} \setminus \{0\}$ that converges to 0 and satisfies $|x_i z| < 1$ for all $i \in \mathbb{N}$. Then $\big({}_2\varphi_1(a, b/x_i; c; q, x_i z)\big)_{i=1}^{\infty}$ converges to ${}_1\varphi_1(a;c;q,bz)$.*

Proof. By definition, we have that ${}_2\varphi_1(a,b/x_i;c;q,x_iz) = \sum_{n=0}^{\infty} \frac{(a;q)_n}{(c;q)_n (q;q)_n}\, z^n\, (b/x_i;q)_n\, x_i^n$ and $(b/x_i;q)_n\, x_i^n = \prod_{k=0}^{n-1}(x_i - bq^k)$. It follows that the n-th term of this series converges to the number $\frac{(a;q)_n}{(c;q)_n(q;q)_n}\,(-1)^n q^{\frac{1}{2}n(n-1)}\,(bz)^n$ as $i \to \infty$. Take $\varepsilon > 0$ such that $|\varepsilon z| < 1$. So there exists $i_0 \in \mathbb{N}$ such that $|x_i| < \varepsilon/2$ for all $i \in \mathbb{Z}_{\geq i_0}$. It is not so difficult to see that there exists $M > 0$ such that $|(b/x_i;q)_n\, x_i^n \varepsilon^{-n}| \leq M$ for all $n \in \mathbb{N}_0$ and $i \in \mathbb{Z}_{\geq i_0}$. Thus, since $\sum_{n=0}^{\infty} \left| \frac{(a;q)_n}{(c;q)_n(q;q)_n}\,(\varepsilon z)^n \right| < \infty$, the dominated convergence theorem implies that the sequence $\big({}_2\varphi_1(a,b/x_i;c;q,x_iz)\big)_{i=1}^{\infty}$ converges to $\sum_{n=0}^{\infty} \frac{(a;q)_n}{(c;q)_n(q;q)_n}\,(-1)^n q^{\frac{1}{2}n(n-1)}\,(bz)^n = {}_1\varphi_1(a;c;q,bz)$. □

This limit transition is used to get a more direct relationship between ${}_2\varphi_1$-functions and Ψ-functions.

Lemma 6.3. *Let $a,b,z \in \mathbb{C}$ such that $|z| < 1$. Then $(z;q)_\infty\, {}_2\varphi_1(a,b;0;q,z) = \Psi(a;az;q,bz)$.*

Proof. First suppose that $a \neq 0$ and $az \notin q^{-\mathbb{N}_0}$. Take a sequence $(x_i)_{i=1}^{\infty}$ in $\mathbb{C} \setminus (\{0\} \cup q^{-\mathbb{N}_0})$ such that $(x_i)_{i=1}^{\infty} \to 0$ and $|x_i/a| < 1$ for all $i \in \mathbb{N}$.
Then Heine's transformation formula [6, Eq. (1.4.5)] implies that ${}_2\varphi_1(a,b;x_i;q,z) = (x_i/a, az;q)_\infty\, (x_i,z;q)_\infty^{-1}\, {}_2\varphi_1(abz/x_i, a; az; q, x_i/a)$. If we let i tend to ∞ in this equality, we obtain by the previous lemma the equality ${}_2\varphi_1(a,b;0;q,z) = (az;q)_\infty\,(z;q)_\infty^{-1}\, {}_1\varphi_1(a;az;q,bz) = (z;q)_\infty^{-1}\,\Psi(a;az;q,bz)$. The general result follows by analytic continuation. □

The following formula will be used throughout the paper.

Result 6.4. *Consider $a,b,z \in \mathbb{C}$ such that $b \neq 0$. Then $\Psi(a;b;q,z) = \Psi(az/b;z;q,b)$.*

Proof. First assume that b and z do not belong to $q^{-\mathbb{N}_0}$. Take a sequence $(x_i)_{i=1}^{\infty}$ in $\mathbb{C} \setminus (\{0\} \cup q^{-\mathbb{N}_0})$ such that $(x_i)_{i=1}^{\infty} \to 0$, $|zx_i| < 1$ and $|bx_i| < 1$ for all $i \in \mathbb{N}$. Let $i \in \mathbb{N}$ and apply Heine's transformation formula [6, Eq. (1.4.5)] for $a \rightsquigarrow a$, $b \rightsquigarrow 1/x_i$, $c \rightsquigarrow b$ and $z \rightsquigarrow x_i z$. Thus, ${}_2\varphi_1(a,1/x_i;b;q,x_iz) = (bx_i, z;q)_\infty\,(b, x_iz;q)_\infty^{-1}\, {}_2\varphi_1(az/b, 1/x_i; z; q, bx_i)$. If we let i tend to ∞ in this equality, Lemma 6.2 implies that ${}_1\varphi_1(a;b;q,z) = (z;q)_\infty\,(b;q)_\infty^{-1}\, {}_1\varphi_1(az/b;z;q,b)$. Thus, $\Psi(a;b;q,z) = \Psi(az/b;z;q,b)$, still under the assumption that $b,z \notin q^{-\mathbb{N}_0}$. The general formula now follows from analytic continuation. □

Roughly speaking, q-contiguous relations are relations between expressions of the form $\Psi(ra;sb;q,tz)$ where $r,s,t \in \{1,q,q^{-1}\}$. We need the following two.

Lemma 6.5. *Consider $a,b,z \in \mathbb{C}$, then*

$$\Psi(a;b;q,z) = (1-a)\ \Psi(qa;b;q,z) + a\ \Psi(a;b;q,qz)\ . \tag{6.2}$$

$$\Psi(a;b;q,z) = (a-b)\ \Psi(a;qb;q,qz) + (1-a)\ \Psi(qa;qb;q,z)\ . \tag{6.3}$$

Proof. By definition, we have

$$\begin{aligned}
(1-a)\ \Psi(qa;b;q,z) &= \sum_{n=0}^{\infty} (1-a)\, \frac{(qa;q)_n\,(q^nb;q)_\infty}{(q;q)_n}\,(-1)^n\, q^{\frac{1}{2}n(n-1)}\, z^n \\
&= \sum_{n=0}^{\infty} \frac{(a;q)_n\,(1-aq^n)\,(q^nb;q)_\infty}{(q;q)_n}\,(-1)^n\, q^{\frac{1}{2}n(n-1)}\, z^n \\
&= \sum_{n=0}^{\infty} \frac{(a;q)_n\,(q^nb;q)_\infty}{(q;q)_n}\,(-1)^n\, q^{\frac{1}{2}n(n-1)}\, z^n - a \sum_{n=0}^{\infty} \frac{(a;q)_n\,(q^nb;q)_\infty}{(q;q)_n}\,(-1)^n\, q^{\frac{1}{2}n(n-1)}\,(qz)^n \\
&= \Psi(a;b;q,z) - a\ \Psi(a;b;q,qz)\ .
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&(a-b)\ \Psi(a;qb;q,qz)+(1-a)\ \Psi(qa;qb;q,z)\\
&\quad=\sum_{n=0}^{\infty}(a-b)\,\frac{(a;q)_n\,(q^{n+1}b;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,(qz)^n\\
&\qquad+\sum_{n=0}^{\infty}(1-a)\,\frac{(qa;q)_n\,(q^{n+1}b;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=\sum_{n=0}^{\infty}(a-b)q^n\,\frac{(a;q)_n\,(q^{n+1}b;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\qquad+\sum_{n=0}^{\infty}(1-q^na)\,\frac{(a;q)_n\,(q^{n+1}b;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=\sum_{n=0}^{\infty}(1-q^nb)\,\frac{(a;q)_n\,(q^{n+1}b;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=\sum_{n=0}^{\infty}\frac{(a;q)_n\,(q^nb;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n=\Psi(a;b;q,z)\ .
\end{aligned}$$

□

We need the following transformation formulas.

Proposition 6.6. *Consider $a,b,z\in\mathbb{C}$ and $k\in\mathbb{Z}$. Then*

(1) $(q^{k+1}/a;q)_\infty\,\Psi(aq^{-k};q^{-k+1};q,z)=(q/a;q)_\infty\,(az/q)^k\ \Psi(a;q^{k+1};q,zq^k)$ *if $a,z\neq0$,*

(2) $(q^{k+1}a/z;q)_\infty\,\Psi(q^{-k};a;q,z)=(z/q)^k\,(q^ka;q)_\infty\,\Psi(q^{-k};qa/z;q,q^2/z)$ *if $z\neq0$ and $k\geq0$.*

Proof. (1) Suppose first that $k\geq0$. Note that $(q^{n+1-k};q)_\infty=0$ for all $n\in\mathbb{Z}_{\leq k-1}$. Hence,

$$\begin{aligned}
&(q^{k+1}/a;q)_\infty\,\Psi(aq^{-k};q^{-k+1};q,z)=(q^{1+k}/a;q)_\infty\sum_{n=k}^{\infty}\frac{(aq^{-k};q)_n(q^{n+1-k};q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=(q^{k+1}/a;q)_\infty\sum_{n=0}^{\infty}\frac{(aq^{-k};q)_{n+k}(q^{n+1};q)_\infty}{(q;q)_{n+k}}\,(-1)^{n+k}\,q^{\frac{1}{2}(n+k)(n+k-1)}\,z^{n+k}\\
&\quad=(-1)^k\,q^{\frac{1}{2}k(k-1)}\ z^k\,(q^{k+1}/a;q)_\infty\,(a/q;q^{-1})_k\sum_{n=0}^{\infty}\frac{(a;q)_n(q^{n+k+1};q)_\infty}{(q;q)_n}\,(-1)^n\,q^{nk}\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=(-1)^k\,q^{\frac{1}{2}k(k-1)}\ z^k\,(q^{k+1}/a;q)_\infty\,(q/a;q)_k\,(-a)^k\,q^{-k}\,q^{-\frac{1}{2}k(k-1)}\ \Psi(a;q^{k+1};q,zq^k)\\
&\quad=(q/a;q)_\infty\,(az/q)^k\ \Psi(a;q^{k+1};q,zq^k)\ .
\end{aligned}$$

If $k<0$, we apply (1) where we replace k by $-k$, a by aq^{-k} and z by zq^k to obtain (1) in this case.

(2) First we assume that $a\neq0$ and $|q^{k+1}a/z|<1$. Since $(q^{-k};q)_n=0$ if $n\geq k+1$, the series terminates and

$$\begin{aligned}
&(q^{k+1}a/z;q)_\infty\,\Psi(q^{-k};a;q,z)=(q^{k+1}a/z;q)_\infty\sum_{n=0}^{k}\frac{(q^{-k};q)_n\,(q^na;q)_\infty}{(q;q)_n}\,(-1)^n\,q^{\frac{1}{2}n(n-1)}\,z^n\\
&\quad=(q^{k+1}a/z;q)_\infty\sum_{n=0}^{k}\frac{(q^{-k};q)_{k-n}\,(q^{k-n}a;q)_\infty}{(q;q)_{k-n}}\,(-1)^{k-n}\,q^{\frac{1}{2}(k-n)(k-n-1)}\,z^{k-n}\\
&\quad=(q^{k+1}a/z;q)_\infty\,(-1)^k\,q^{\frac{1}{2}k(k-1)}\ z^k\sum_{n=0}^{k}\frac{(q^{-k};q)_{k-n}\,(q^{k-n}a;q)_\infty}{(q;q)_{k-n}}\,(-1)^n\,q^{-kn}\,q^{\frac{1}{2}n(n+1)}\,z^{-n}\ . \qquad (6.4)
\end{aligned}$$

Now,

$$\begin{aligned}(q^{k-n}a;q)_\infty &= (q^k a;q)_\infty\,(q^{k-1}a;q^{-1})_n = (q^k a;q)_\infty\; q^{nk}\, q^{-\frac{1}{2}n(n+1)}\,(-a)^n\,(q^{1-k}/a;q)_n\end{aligned}$$

$$\begin{aligned}\frac{(q^{-k};q)_{k-n}}{(q;q)_{k-n}} &= \frac{\prod_{i=-k}^{-1}(1-q^i)}{\prod_{i=-n}^{-1}(1-q^i)}\,\frac{\prod_{i=k-n+1}^{k}(1-q^i)}{\prod_{i=1}^{k}(1-q^i)} = (-1)^k\, q^{-\frac{1}{2}k(k+1)}\,\frac{(-1)^n\, q^{kn}\, q^{-\frac{1}{2}n(n-1)}\,(q^{-k};q)_n}{(-1)^n\, q^{-\frac{1}{2}n(n+1)}\,(q;q)_n}\\ &= (-1)^k\, q^{-\frac{1}{2}k(k+1)}\, q^{(k+1)n}\,\frac{(q^{-k};q)_n}{(q;q)_n}\end{aligned}$$

which upon substitution in Eq. (6.4) implies that

$$\begin{aligned}&(q^{k+1}a/z;q)_\infty\,\Psi(q^{-k};a;q,z) = (q^{k+1}a/z;q)_\infty\,(q^k a;q)_\infty\,(z/q)^k \sum_{n=0}^{k} \frac{(q^{-k};q)_n\,(q^{1-k}/a;q)_n}{(q;q)_n}\,(q^{k+1}a/z)^n\\ &\quad = (q^{k+1}a/z;q)_\infty\,(q^k a;q)_\infty\,(z/q)^k\;{}_2\varphi_1(q^{-k},q^{1-k}/a;0;q,q^{k+1}a/z)\\ &\quad = (q^k a;q)_\infty\,(z/q)^k\;\;\Psi(q^{-k};qa/z;q,q^2/z)\ ,\end{aligned}$$

where we used Lemma 6.3 in the last step. The general result now follows by analytic continuation. □

In most cases it is not necessary to know the precise expression for the Ψ-functions, sometimes only the asymptotic behavior matters. We need the following estimate (see [14, Lem. 2.8]).

Lemma 6.7. *Consider* $a,z \in \mathbb{C}$*,* $k \in \mathbb{Z}_{\geq 0}$*. Then* $|\Psi(a;q^{1-k};q,z)| \leq (-|a|;q)_\infty\,(-|z|;q)_\infty\,|z|^k\,q^{\frac{1}{2}k(k-1)}$*.*

Proof. For $n \in \mathbb{N}_0$, we have $|(a;q)_n| \leq (-|a|;q)_n \leq (-|a|;q)_\infty$. Also notice that $(q^{n-k+1};q)_\infty = 0$ for all $n \in \mathbb{N}_0$ satisfying $n < k$. Hence, by the definition of Ψ, we see that

$$\begin{aligned}&\big|\,\Psi(a;q^{1-k};q,z)\,\big| \leq (-|a|;q)_\infty \sum_{n=0}^{\infty} \frac{(q^{n-k+1};q)_\infty}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\,|z|^n\\ &\quad = (-|a|;q)_\infty \sum_{n=k}^{\infty} \frac{(q^{n-k+1};q)_\infty}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\,|z|^n = (-|a|;q)_\infty \sum_{n=0}^{\infty} \frac{(q^{n+1};q)_\infty}{(q;q)_{n+k}}\, q^{\frac{1}{2}(n+k)(n+k-1)}\,|z|^{n+k}\\ &\quad = (-|a|;q)_\infty\,|z|^k \sum_{n=0}^{\infty} \frac{(q^{n+k+1};q)_\infty}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\, q^{nk}\, q^{\frac{1}{2}k(k-1)}\,|z|^n\\ &\quad \leq (-|a|;q)_\infty\,|z|^k\, q^{\frac{1}{2}k(k-1)} \sum_{n=0}^{\infty} \frac{1}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\,(q^k\,|z|)^n = (-|a|;q)_\infty\,(-|z|q^k;q)_\infty\,|z|^k\, q^{\frac{1}{2}k(k-1)}\end{aligned}$$

where we used [6, Eq. 1.3.16] in the last equality. Now the lemma follows. □

The last two estimates play a vital role in the proof of the coassociativity of the comultiplication.

Lemma 6.8. *Consider* $a,b,c \in \mathbb{C}$*,* $p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ *and* $\beta > 0$*. Let* $m \in [1,\infty]$*,* $M > 0$*,* $\varepsilon > 0$ *and define the q-interval J as* $J = \{\, x \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}} \mid |x| \leq M \text{ or } \operatorname{sgn}(x) = \operatorname{sgn}(p)\,\}$*. Then there exist* $N > 0$ *and* $f \in l^m(J)^+$ *such that*

$$|\Psi(a/\lambda;px;q,b\lambda)| \leq N \qquad \text{and} \qquad |x|^\beta\;|(cx;q)_\infty|^{\frac{1}{2}}\;|\Psi(a/\lambda;px;q,b\lambda)| \leq f(x)$$

for all $x \in J$ *and* $\lambda \in \mathbb{C} \setminus \{0\}$ *such that* $|\lambda| \leq \varepsilon$*.*

Proof. Choose $k \in \mathbb{Z}$ such that $|p| = q^k$. Take $r,s > 0$ such that $r > |b|(\varepsilon + |a|)$ and $s > |c|$. Let $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$ and $x \in J$. Since $(pxq^r;q)_\infty \geq 0$ for all $r \in \mathbb{Z}$, we get that

$$\begin{aligned}&|\,\Psi(a/\lambda;px;q,b\lambda)\,| \leq \sum_{n=0}^{\infty} \frac{|(a/\lambda;q)_n|\,(pxq^n;q)_\infty}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\,|b\lambda|^n\\ &\quad = \sum_{n=0}^{\infty} \frac{(pxq^n;q)_\infty}{(q;q)_n}\,\Big(\prod_{i=0}^{n-1} |\lambda - q^i a|\,\Big) q^{\frac{1}{2}n(n-1)}\,|b|^n \leq \sum_{n=0}^{\infty} \frac{(pxq^n;q)_\infty}{(q;q)_n}\, q^{\frac{1}{2}n(n-1)}\,(|b|\,(|\lambda| + |a|))^n\\ &\quad \leq \Psi(0;px;q,-r)\ .\end{aligned} \tag{6.5}$$

This implies the existence of $C > 0$ such that $|\,\Psi(a/\lambda; px; q, b\lambda)\,| \leq C$ and $|(cx;q)_\infty|^{\frac{1}{2}} \leq C$ for all $x \in J$ such that $|x| \leq \max\{M, q^2/|p|\}$ and $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$. (6.6)

Define the function $f : J \rightarrow \mathbb{R}^+$ such that for $x \in J$, we have that $f(x) = C^2\,|x|^\beta$ if $|x| \leq \max\{M, q^2/|p|\}$ and

$$f(x) = (-r;q)_\infty\,(-s;q)_\infty^{\frac{1}{2}}\,(-q/s;q)_\infty^{\frac{1}{2}}\,q^{\frac{1}{2}k(k-1)}\,q^{(k+\beta)t}\,r^{1-k-t}\,s^{-\frac{t}{2}}\,q^{\frac{1}{4}t(t-1)}$$

if $|x| > \max\{M, q^2/|p|\}$ and $t \in \mathbb{Z}$ is such that $|x| = q^t$. Then it is clear that $f \in l^m(J)^+$.

Take $x \in J$ such that $|x| > \max\{M, q^2/|p|\}$. Choose $t \in \mathbb{Z}$ such that $|x| = q^t$. Since $x \in J$ and $|x| > M$, we have that $\operatorname{sgn}(x) = \operatorname{sgn}(p)$, hence $px = q^{k+t}$. It is also clear that $t \leq 1-k$, thus $1-k-t \geq 0$. By estimate (6.5) and Lemma 6.7, we get for all $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$,

$$\begin{aligned} |\,\Psi(a/\lambda; px; q, b\lambda)\,| &\leq \Psi(0; px; q, -r) = \Psi(0; q^{1-(1-k-t)}; -r) \leq (-r;q)_\infty\,r^{1-k-t}\,q^{\frac{1}{2}(k+t-1)(k+t)} \\ &= (-r;q)_\infty\,r^{1-k-t}\,q^{\frac{1}{2}k(k-1)}\,q^{kt}\,q^{\frac{1}{2}t(t-1)}\ . \end{aligned} \tag{6.7}$$

Also, Lemma 6.1 guarantees that

$$\begin{aligned} |\,(cx;q)_\infty\,|^{\frac{1}{2}} &\leq (-|cx|;q)^{\frac{1}{2}} \leq (-sq^t;q)_\infty^{\frac{1}{2}} = \frac{(-s;q)_\infty^{\frac{1}{2}}\,(-q/s;q)_\infty^{\frac{1}{2}}}{(-q^{1-t}/s;q)_\infty^{\frac{1}{2}}}\,s^{-\frac{t}{2}}\,q^{-\frac{1}{4}t(t-1)} \\ &\leq (-s;q)_\infty^{\frac{1}{2}}\,(-q/s;q)_\infty^{\frac{1}{2}}\,s^{-\frac{t}{2}}\,q^{-\frac{1}{4}t(t-1)}\ . \end{aligned}$$

This estimate, together with the estimates in (6.6) and (6.7) imply that $|x|^\beta\ |(cx;q)_\infty|^{\frac{1}{2}}\ |\Psi(a/\lambda; px; q, b\lambda)| \leq f(x)$ for all $x \in J$ and $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$.

Estimate (6.7) also implies the existence of $D > 0$ such that $|\Psi(a/\lambda; px; q, b\lambda)| \leq D$ for all $x \in J$ such that $|x| > \max\{M, q^2/|p|\}$ and $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$. So if we set $N = \max\{C, D\}$, we find by the estimate in (6.6) that $|\Psi(a/\lambda; px; q, b\lambda)| \leq N$ for all $x \in J$ and $\lambda \in \mathbb{C} \setminus \{0\}$ such that $|\lambda| \leq \varepsilon$. ☐

The next result is an easy consequence of the previous lemma.

Lemma 6.9. *Consider* $a, b, c \in \mathbb{C}$, $p \in -q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$. *Let* $(x_i)_{i\in I}$ *be a net in* $-q^{\mathbb{Z}} \cup q^{\mathbb{Z}}$ *such that* $\operatorname{sgn}(x_i) = \operatorname{sgn}(p)$ *for all* $i \in I$ *and* $(x_i)_{i\in I} \rightarrow \infty$. *Then the net* $\big(|(cx_i;q)_\infty|^{\frac{1}{2}}\ |\Psi(ax_i; px_i; q, b/x_i)|\big)_{i\in I}$ *converges to* 0.

REFERENCES

[1] E. Abe, Hopf algebras. *Cambridge Tracts in Mathematics* **74**, *Cambridge University Press, Cambridge* (1980).

[2] W.A. Al-Salam & L. Carlitz, Some orthogonal q-polynomials. *Math. Nach.* **30** (1965), 47–61.

[3] N. Ciccoli, E. Koelink & T. Koornwinder, q-Laguerre polynomials and big q-Bessel functions and their orthogonality relations. *Methods and Applications of Analysis.* **6, no. 1** (1999), 109–127.

[4] J. Dixmier, Les algèbres d'opérateurs dans l'espace hilbertien. Deuxième édition. *Gauthier-Villars, Paris* (1969).

[5] M. Enock & J.-M. Schwartz, Kac algebras and duality of locally compact groups. *Springer-Verlag, Berlin* (1992).

[6] G. Gasper & M. Rahman, Basic hypergeometric series. *Cambridge University Press* (1990)

[7] R.V. Kadison & J.R. Ringrose, Fundamentals of the theory of operator algebras I. *Graduate Studies in Mathematics, 15. Am. Math. Soc.* (1997).

[8] T. Kakehi, T. Masuda & K. Ueno, Spectral analysis of a q-difference operator which arises from the quantum $SU(1,1)$ group. *Journal of Operator Theory* **33**, 159–196 (1995).

[9] L.I. Korogodsky, Quantum Group $SU(1,1) \rtimes \mathbb{Z}_2$ and super tensor products. *Commun. Math. Phys.* **163** (1994), 433–460.

[10] L.I. Korogodsky & L.L. Vaksman, Spherical functions on the quantum group $SU(1,1)$ and the q-analogue of the Mehler-Fock formula. *(Russian) Funktsional. Anal. i Prilozhen.* **25, no. 1** (1991), 60–62. Translation in *Funct. Anal. Appl.* **25, no. 1** (1991), 48–49.

[11] R. Koekoek and R.F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue. *Report 98-17, Delft Univ. Technology* (1998). `http://aw.twi.tudelft.nl/~koekoek/askey.html`

[12] E. Koelink, Spectral theory and special functions. *Lecture notes for the 2000 Laredo Summer school of the SIAM Activity Group* (2000). `http://aw.twi.tudelft.nl/~koelink/laredo.html`

[13] E. Koelink & J.V. Stokman, The big q-Jacobi function transform. *Preprint Institute de Recherche Mathématique Avancée, Univ. Louis Pasteur & CNRS, Strasbourg* **no. 1999/23** (1999),#math.CA/9904111

[14] E. Koelink & J.V. Stokman, with an appendix by M. Rahman, Fourier transforms on the quantum $SU(1,1)$ group. *Preprint Institut de Mathématiques de Jussieu* (1999), #math.QA/9911163

[15] T.H. Koornwinder, The addition formula for little q-Legendre polynomials and the $SU(2)$ quantum group. *SIAM J. Math. Anal.* **22** (1991), 295–301.

[16] J. Kustermans, KMS-weights on C*-algebras. *Preprint Odense Universitet* (1997) #funct-an/9704008

[17] J. Kustermans, One-parameter representations on C*-algebras. *Preprint Odense Universitet* (1997) #funct-an/9707009

[18] J. Kustermans & S. Vaes, Locally compact quantum groups. *Ann. scient. Éc. Norm. Sup.* **4è série, t.33** (2000), 837–934.

[19] J. Kustermans & S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting. *To appear in Math. Scand.* (2000) #math.QA/0005219

[20] T. Masuda, K. Mimachi, Y. Nakagami, M. Noumi, Y. Saburi & K. Ueno, Unitary representations of the quantum group $SU_q(1,1)$: structure of the dual space of $\mathcal{U}_q(\mathfrak{sl}(2))$. *Letters in Math. Phys.* **19**, 187-194 (1990).

[21] T. Masuda, K. Mimachi, Y. Nakagami, M. Noumi, Y. Saburi & K. Ueno, Unitary representations of the quantum group $SU_q(1,1)$: II - matrix elements of unitary representations and the basic hypergeometric functions. *Letters in Math. Phys.* **19**, 195-204 (1990).

[22] G.K. Pedersen & M. Takesaki, The Radon-Nikodym theorem for von Neumann algebras. *Acta Math.* **130** (1973), 53–87.

[23] K. Schmüdgen, Unbounded operator algebras and representation theory. *Operator Theory* **37**, Birkhäuser (1990).

[24] D. Shklyarov, S. Sinel'shchikov & L.L. Vaksman, On function theory in quantum discs: invariant kernels. *Preprint Institute for Low Temperature Physics & Engineering* (1998). #math.QA/9808047

[25] S. Stratila, Modular Theory in Operator Algebras. *Abacus Press, Turnbridge Wells, England* (1981).

[26] S. Vaes, A Radon-Nikodym theorem for von Neumann algebras. *To appear in J. Operator Theory* (1998). #math.OA/9811122

[27] S.L. Woronowicz, Unbounded elements affiliated with C*-algebras and non-compact quantum groups. *Commun. Math. Phys.* **136** (1991), 399–432.

[28] S.L. Woronowicz, C*-algebras generated by unbounded elements. *Rev. Math. Phys.* **7, no. 3** (1995), 481–521.

[29] S.L. Woronowicz, From multiplicative unitaries to quantum groups. *Int. J. Math.*, **7, no. 1** (1996), 127–149.

[30] S.L. Woronowicz, Extended $SU(1,1)$ quantum group. Hilbert space level. *Preprint KMMF, not completely finished yet.* (2000)

[31] S.L. Woronowicz & S. Zakrzewski, Quantum $ax+b$ group. Preprint KMMF (1999)